

# Classification, reduction and stability of toric principal bundles



Jyoti Dasgupta

(Joint work with Indranil Biswas, Arijit Dey, Bivas Khan  
and Mainak Poddar)


IISER Pune

IISc-IISER Joint Math Symposium 2021

September 18, 2021



vector bundle



Principal bundle

- Let  $G$  denote a complex linear algebraic group.
- A principal  $G$ -bundle  $\pi : \mathcal{E} \rightarrow X$  is a variety  $\mathcal{E}$  with a right  $G$ -action, the action being free, such that  $\pi$  is  $G$ -equivariant, where  $X$  is being given the trivial  $G$ -action.

# Principal bundles

- Let  $G$  denote a complex linear algebraic group.
- A principal  $G$ -bundle  $\pi : \mathcal{E} \rightarrow X$  is a variety  $\mathcal{E}$  with a right  $G$ -action, the action being free, such that  $\pi$  is  $G$ -equivariant, where  $X$  is being given the trivial  $G$ -action.
- Further, the bundle is assumed to be locally trivial in the étale topology. This means that, for every point  $x \in X$ , there exists a neighbourhood  $U$  and an étale morphism  $U' \rightarrow U$  such that when  $\mathcal{E}$  is pulled back to  $U'$ , it is trivial as a  $G$ -bundle.

## Examples of Principal Bundles

- Let  $X$  be a variety. Then for an algebraic group  $G$ , we have the trivial principal  $G$ -bundle  $X \times G \rightarrow X$  with the  $G$ -action on the second factor by right multiplication.

## Examples of Principal Bundles

- Let  $X$  be a variety. Then for an algebraic group  $G$ , we have the trivial principal  $G$ -bundle  $X \times G \rightarrow X$  with the  $G$ -action on the second factor by right multiplication.
- Let  $Y$  be an algebraic variety with free right  $G$ -action by a reductive group  $G$  and let the geometric quotient  $Y \rightarrow Y/G$  exists. Then  $Y \rightarrow Y/G$  is a principal  $G$ -bundle.

## Examples of Principal Bundles

- Let  $X$  be a variety. Then for an algebraic group  $G$ , we have the trivial principal  $G$ -bundle  $X \times G \rightarrow X$  with the  $G$ -action on the second factor by right multiplication.
- Let  $Y$  be an algebraic variety with free right  $G$ -action by a reductive group  $G$  and let the geometric quotient  $Y \rightarrow Y/G$  exists. Then  $Y \rightarrow Y/G$  is a principal  $G$ -bundle.
- Let  $\mathcal{P} \rightarrow X$  be a principal  $G$ -bundle and let  $f : Y \rightarrow X$  be any morphism. Then the pullback bundle  $f^*(\mathcal{P}) := \mathcal{P} \times_X Y \rightarrow Y$  is a principal  $G$ -bundle over  $Y$ .

## DEFINITION 1

A *toric variety* is a normal variety  $X$  such that

- (1) a torus  $T \cong (\mathbb{C}^*)^n$  is a Zariski dense open subset of  $X$ , and
- (2) the natural action of  $T$  on itself extends to an action of  $T$  on  $X$ .




## DEFINITION 1

A *toric variety* is a normal variety  $X$  such that

- (1) a torus  $T \cong (\mathbb{C}^*)^n$  is a Zariski dense open subset of  $X$ , and
- (2) the natural action of  $T$  on itself extends to an action of  $T$  on  $X$ .

## EXAMPLE 2

- $(\mathbb{C}^*)^n, \mathbb{C}^n, \mathbb{P}^n$ .
- Projectivization of direct sum of line bundles on a toric variety.
- Blow up of a toric variety at an invariant subvariety.



$$\Phi: (\mathbb{C}^*)^2 \longrightarrow \mathbb{P}^2$$

$$(t_1, t_2) \mapsto [t_1^0 t_2^0 : t_1^1 t_2^0 : t_1^0 t_2^1] = [1 : t_1 : t_2]$$

Closure of the "image" of  $\Phi$  is  $\mathbb{P}^2$ .

- For simplicity, we will assume that  $X$  is complete and nonsingular.

- For simplicity, we will assume that  $X$  is complete and nonsingular.
- Then  $X$  can be written as the union of affine toric varieties  
 $X = \cup_{\sigma \in \Xi^*} X_\sigma$ , where  $X_\sigma \cong \mathbb{C}^n$ .

- For simplicity, we will assume that  $X$  is complete and nonsingular.
- Then  $X$  can be written as the union of affine toric varieties  
 $X = \cup_{\sigma \in \Xi^*} X_\sigma$ , where  $X_\sigma \cong \mathbb{C}^n$ .
- Each  $X_\sigma$  contains a unique  $T$ -fixed point denoted by  $x_\sigma$ .  $\Xi^*$  may be identified with the set of  $T$ -fixed points of  $X$ .

- For simplicity, we will assume that  $X$  is complete and nonsingular.
- Then  $X$  can be written as the union of affine toric varieties  $X = \cup_{\sigma \in \Xi^*} X_\sigma$ , where  $X_\sigma \cong \mathbb{C}^n$ .
- Each  $X_\sigma$  contains a unique  $T$ -fixed point denoted by  $x_\sigma$ .  $\Xi^*$  may be identified with the set of  $T$ -fixed points of  $X$ .
- We identify the open orbit  $O$  in  $X$  with  $T$ . Let  $x_0 \in O$  correspond to  $1_T \in T$ .

- The study of torus equivariant vector bundles on toric varieties was initiated by Kaneyama (1975,'88) and Klyachko (1989).

## Toric vector bundles

- The study of torus equivariant vector bundles on toric varieties was initiated by Kaneyama (1975,'88) and Klyachko (1989).
- Kaneyama proved the existence of an equivariant splitting for any torus equivariant vector bundle over  $\mathbb{P}^n$  of rank  $r < n$ . This is closely related to the Hartshorne conjecture on the splitting of any rank two vector bundle over  $\mathbb{P}^n$  for  $n \geq 7$ .



## Toric vector bundles

- The study of torus equivariant vector bundles on toric varieties was initiated by Kaneyama (1975,'88) and Klyachko (1989).
- Kaneyama proved the existence of an equivariant splitting for any torus equivariant vector bundle over  $\mathbb{P}^n$  of rank  $r < n$ . This is closely related to the Hartshorne conjecture on the splitting of any rank two vector bundle over  $\mathbb{P}^n$  for  $n \geq 7$ .
- Klyachko used the stability of toric vector bundles in his proof of Horn's conjecture on eigenvalues of sums of Hermitian matrices.

## Toric vector bundles


- The study of torus equivariant vector bundles on toric varieties was initiated by Kaneyama (1975,'88) and Klyachko (1989).
- Kaneyama proved the existence of an equivariant splitting for any torus equivariant vector bundle over  $\mathbb{P}^n$  of rank  $r < n$ . This is closely related to the Hartshorne conjecture on the splitting of any rank two vector bundle over  $\mathbb{P}^n$  for  $n \geq 7$ .
- Klyachko used the stability of toric vector bundles in his proof of Horn's conjecture on eigenvalues of sums of Hermitian matrices.

**Goal:** Study “toric principal bundles” over toric varieties.


## Toric principal bundles

Let  $G$  denote a complex linear algebraic group. Let  $X$  be a toric variety with dense torus  $T$ , then a principal  $G$ -bundle  $\pi : \mathcal{E} \rightarrow X$  is said to be a **toric principal bundle** if  $\mathcal{E}$  is endowed with a lift of  $T$ -action on  $X$ .

Moreover, the  $T$ -action on  $\mathcal{E}$  must commute with the  $G$ -action.



## Equivariant trivialization over affine toric variety:



For any  $\sigma \in \Xi^*$ , set  $\mathcal{E}_\sigma := \mathcal{E}|_{X_\sigma}$ .

A section  $s_\sigma : X_\sigma \rightarrow \mathcal{E}_\sigma$  is called a **distinguished section** if

$$t s_\sigma(x) = s_\sigma(tx) \cdot \rho_\sigma(t), \quad \forall x \in X_\sigma, \quad \forall t \in T$$

where  $\rho_\sigma : T \rightarrow G$  is a homomorphism of algebraic groups.

## Equivariant trivialization over affine toric variety:



For any  $\sigma \in \Xi^*$ , set  $\mathcal{E}_\sigma := \mathcal{E}|_{X_\sigma}$ .

A section  $s_\sigma : X_\sigma \rightarrow \mathcal{E}_\sigma$  is called a **distinguished section** if

$$t s_\sigma(x) = s_\sigma(tx) \cdot \rho_\sigma(t), \quad \forall x \in X_\sigma, \quad \forall t \in T$$

where  $\rho_\sigma : T \rightarrow G$  is a homomorphism of algebraic groups.

$\mathcal{E}_\sigma$  is trivial and admits a distinguished section.

## Classification of distinguished sections


Let  $s_\sigma$  be a distinguished section with associated homomorphism  $\rho_\sigma$ .  
Then for any  $g \in G$ ,  $s_\sigma \cdot g$  is a distinguished section with associated homomorphism  $g^{-1}\rho_\sigma g$ .

## Classification of distinguished sections


Let  $s_\sigma$  be a distinguished section with associated homomorphism  $\rho_\sigma$ . Then for any  $g \in G$ ,  $s_\sigma \cdot g$  is a distinguished section with associated homomorphism  $g^{-1}\rho_\sigma g$ .

Suppose  $s_\sigma$  and  $s'_\sigma$  are two arbitrary distinguished sections for  $\mathcal{E}_\sigma$  with homomorphisms  $\rho_\sigma, \rho'_\sigma$  respectively. Then,

- 1  $\rho'_\sigma(t) = \alpha_\sigma^{-1} \rho_\sigma(t) \alpha_\sigma$ .
- 2  $\rho_\sigma(t) \beta_\sigma \alpha_\sigma^{-1} \rho_\sigma(t)^{-1}$  extends to a  $G$ -valued function over  $X_\sigma$ .



## Admissible collections



An **admissible collection**  $\{\rho_\sigma, P(\tau, \sigma)\}$  consists of a collection of homomorphisms

$$\{\rho_\sigma : T \longrightarrow G \mid \sigma \in \Xi^*\}$$

and a collection of elements  $\{P(\tau, \sigma) \in G \mid \tau, \sigma \in \Xi^*\}$ .



## Classification of toric principal bundles

$\{\rho_\sigma, P(\tau, \sigma)\}$  satisfies the following conditions:

## Classification of toric principal bundles

$\{\rho_\sigma, P(\tau, \sigma)\}$  satisfies the following conditions:

- 1 For  $\tau, \sigma \in \Xi^*$ , the function  $t \mapsto \rho_\tau(t)P(\tau, \sigma)\rho_\sigma(t)^{-1}$  extends to a  $G$ -valued function over  $X_\sigma \cap X_\tau \supset T$ .

## Classification of toric principal bundles

$\{\rho_\sigma, P(\tau, \sigma)\}$  satisfies the following conditions:

- 1 For  $\tau, \sigma \in \Xi^*$ , the function  $t \mapsto \rho_\tau(t)P(\tau, \sigma)\rho_\sigma(t)^{-1}$  extends to a  $G$ -valued function over  $X_\sigma \cap X_\tau \supset T$ .
- 2  $P(\sigma, \sigma) = 1_G$ .

## Classification of toric principal bundles

$\{\rho_\sigma, P(\tau, \sigma)\}$  satisfies the following conditions:

- 1 For  $\tau, \sigma \in \Xi^*$ , the function  $t \mapsto \rho_\tau(t)P(\tau, \sigma)\rho_\sigma(t)^{-1}$  extends to a  $G$ -valued function over  $X_\sigma \cap X_\tau \supset T$ .
- 2  $P(\sigma, \sigma) = 1_G$ .
- 3 For  $\tau, \sigma, \delta \in \Xi^*$ , the cocycle condition  $P(\tau, \sigma)P(\sigma, \delta)P(\delta, \tau) = 1_G$  holds.

## Classification of toric principal bundles

$\{\rho_\sigma, P(\tau, \sigma)\}$  satisfies the following conditions:

- 1 For  $\tau, \sigma \in \Xi^*$ , the function  $t \mapsto \rho_\tau(t)P(\tau, \sigma)\rho_\sigma(t)^{-1}$  extends to a  $G$ -valued function over  $X_\sigma \cap X_\tau \supset T$ .
- 2  $P(\sigma, \sigma) = 1_G$ .
- 3 For  $\tau, \sigma, \delta \in \Xi^*$ , the cocycle condition  $P(\tau, \sigma)P(\sigma, \delta)P(\delta, \tau) = 1_G$  holds.

Two admissible collections are said to be equivalent if they satisfy certain conjugacy relations and some extension conditions.

## Classification of toric principal bundles

$\{\rho_\sigma, P(\tau, \sigma)\}$  satisfies the following conditions:

- 1 For  $\tau, \sigma \in \Xi^*$ , the function  $t \mapsto \rho_\tau(t)P(\tau, \sigma)\rho_\sigma(t)^{-1}$  extends to a  $G$ -valued function over  $X_\sigma \cap X_\tau \supset T$ .
- 2  $P(\sigma, \sigma) = 1_G$ .
- 3 For  $\tau, \sigma, \delta \in \Xi^*$ , the cocycle condition  $P(\tau, \sigma)P(\sigma, \delta)P(\delta, \tau) = 1_G$  holds.

Two admissible collections are said to be equivalent if they satisfy certain conjugacy relations and some extension conditions.

**Theorem 1** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*The isomorphism classes of  $T$ -equivariant principal  $G$ -bundles on  $X$  are in one-to-one correspondence with the “equivalence classes” of admissible collections  $[\{\rho_\sigma, P(\tau, \sigma)\}]$ .*

## Automorphisms of $\mathcal{E}$

- Let  $\mathcal{E}$  be a  $T$ -equivariant principal  $G$ -bundle over  $X$ .

## Automorphisms of $\mathcal{E}$

- Let  $\mathcal{E}$  be a  $T$ -equivariant principal  $G$ -bundle over  $X$ .
- Let  $e \in \mathcal{E}_{x_0}$ , the fiber of  $\mathcal{E}$  at  $x_0$ .




## Automorphisms of $\mathcal{E}$

- Let  $\mathcal{E}$  be a  $T$ -equivariant principal  $G$ -bundle over  $X$ .
- Let  $e \in \mathcal{E}_{x_0}$ , the fiber of  $\mathcal{E}$  at  $x_0$ .
- For each  $\sigma \in \Xi^*$ , take a distinguished section  $s_\sigma : X_\sigma \rightarrow \mathcal{E}$  such that  $s_\sigma(x_0) = e$ .

## Automorphisms of $\mathcal{E}$


- Let  $\mathcal{E}$  be a  $T$ -equivariant principal  $G$ -bundle over  $X$ .
- Let  $e \in \mathcal{E}_{x_0}$ , the fiber of  $\mathcal{E}$  at  $x_0$ .
- For each  $\sigma \in \Xi^*$ , take a distinguished section  $s_\sigma : X_\sigma \rightarrow \mathcal{E}$  such that  $s_\sigma(x_0) = e$ .



Let  $\Phi \in \text{Aut}_T(\mathcal{E})$  and  $\Phi(e) = e \cdot g$  for some  $g \in G$ .

## Automorphisms of $\mathcal{E}$

- Let  $\mathcal{E}$  be a  $T$ -equivariant principal  $G$ -bundle over  $X$ .
- Let  $e \in \mathcal{E}_{x_0}$ , the fiber of  $\mathcal{E}$  at  $x_0$ .
- For each  $\sigma \in \Xi^*$ , take a distinguished section  $s_\sigma : X_\sigma \rightarrow \mathcal{E}$  such that  $s_\sigma(x_0) = e$ .




Let  $\Phi \in \text{Aut}_T(\mathcal{E})$  and  $\Phi(e) = e \cdot g$  for some  $g \in G$ .

- By continuity of  $\Phi$  and density of  $O$  in  $X$ ,  $\Phi$  is determined by  $\Phi|_{\mathcal{E}_O}$ .

## Automorphisms of $\mathcal{E}$

- Let  $\mathcal{E}$  be a  $T$ -equivariant principal  $G$ -bundle over  $X$ .
- Let  $e \in \mathcal{E}_{x_0}$ , the fiber of  $\mathcal{E}$  at  $x_0$ .
- For each  $\sigma \in \Xi^*$ , take a distinguished section  $s_\sigma : X_\sigma \rightarrow \mathcal{E}$  such that  $s_\sigma(x_0) = e$ .




Let  $\Phi \in \text{Aut}_T(\mathcal{E})$  and  $\Phi(e) = e \cdot g$  for some  $g \in G$ .

- By continuity of  $\Phi$  and density of  $O$  in  $X$ ,  $\Phi$  is determined by  $\Phi|_{\mathcal{E}_O}$ .
- By  $T$ -equivariance,  $\Phi|_{\mathcal{E}_O}$  is determined by  $\Phi|_{\mathcal{E}_{x_0}}$ .

## Automorphisms of $\mathcal{E}$

- Let  $\mathcal{E}$  be a  $T$ -equivariant principal  $G$ -bundle over  $X$ .
- Let  $e \in \mathcal{E}_{x_0}$ , the fiber of  $\mathcal{E}$  at  $x_0$ .
- For each  $\sigma \in \Xi^*$ , take a distinguished section  $s_\sigma : X_\sigma \rightarrow \mathcal{E}$  such that  $s_\sigma(x_0) = e$ .



Let  $\Phi \in \text{Aut}_T(\mathcal{E})$  and  $\Phi(e) = e \cdot g$  for some  $g \in G$ .

- By continuity of  $\Phi$  and density of  $O$  in  $X$ ,  $\Phi$  is determined by  $\Phi|_{\mathcal{E}_O}$ .
- By  $T$ -equivariance,  $\Phi|_{\mathcal{E}_O}$  is determined by  $\Phi|_{\mathcal{E}_{x_0}}$ .
- $\Phi|_{\mathcal{E}_{x_0}}$  is determined by  $\Phi(e)$  using  $G$ -equivariance.

Consider the map  $\xi : \text{Aut}_T(\mathcal{E}) \rightarrow G$ , uniquely determined by the relation

$$\Phi(e) = e \cdot \xi(\Phi), \text{ for all } \Phi \in \text{Aut}_T(\mathcal{E}).$$

This map is an injective homomorphism.

Consider the map  $\xi : \text{Aut}_T(\mathcal{E}) \rightarrow G$ , uniquely determined by the relation

$$\Phi(e) = e \cdot \xi(\Phi), \text{ for all } \Phi \in \text{Aut}_T(\mathcal{E}).$$

This map is an injective homomorphism.

**Theorem 2** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*$\text{Aut}_T(\mathcal{E})$ , the group of  $T$ -equivariant automorphisms of  $\mathcal{E}$ , is given by*

$$\text{Aut}_T(\mathcal{E}) = \{g \in G \mid \text{for all } \sigma \in \Xi^*, \rho_\sigma(t)g\rho_\sigma(t)^{-1} \text{ extends to } X_\sigma\}.$$

Consider the map  $\xi : \text{Aut}_T(\mathcal{E}) \rightarrow G$ , uniquely determined by the relation

$$\Phi(e) = e \cdot \xi(\Phi), \text{ for all } \Phi \in \text{Aut}_T(\mathcal{E}).$$

This map is an injective homomorphism.

**Theorem 2** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*$\text{Aut}_T(\mathcal{E})$ , the group of  $T$ -equivariant automorphisms of  $\mathcal{E}$ , is given by*

$$\text{Aut}_T(\mathcal{E}) = \{g \in G \mid \text{for all } \sigma \in \Xi^*, \rho_\sigma(t)g\rho_\sigma(t)^{-1} \text{ extends to } X_\sigma\}.$$

In general, we have the inclusions

$$Z(G) \subseteq \text{Aut}_T(\mathcal{E}) \subseteq G.$$

For  $X = \mathbb{P}^m \times \mathbb{P}^n$  and  $G = GL(m+n, \mathbb{C})$ , we have  $Z(G) \neq \text{Aut}_T(\mathcal{E})$ , where  $\mathcal{E}$  is the tangent frame bundle of  $X$ .



## Reduction of structure group

- Let  $H$  be a closed subgroup of the complex linear algebraic group  $G$  and let  $\phi : H \hookrightarrow G$  denote the inclusion map.

## Reduction of structure group

- Let  $H$  be a closed subgroup of the complex linear algebraic group  $G$  and let  $\phi : H \hookrightarrow G$  denote the inclusion map.
- Let  $\mathcal{E}_H \rightarrow X$  be a toric principal  $H$ -bundle with admissible collection  $\{\rho_\sigma, P(\tau, \sigma)\}$ .

## Reduction of structure group

- Let  $H$  be a closed subgroup of the complex linear algebraic group  $G$  and let  $\phi : H \hookrightarrow G$  denote the inclusion map.
- Let  $\mathcal{E}_H \rightarrow X$  be a toric principal  $H$ -bundle with admissible collection  $\{\rho_\sigma, P(\tau, \sigma)\}$ .
- Then the associated bundle  $\mathcal{E}_H \times_H G$  is a toric principal  $G$ -bundle with admissible collection  $\{\phi \circ \rho_\sigma, \phi(P(\tau, \sigma))\}$ .

## Reduction of structure group

- Let  $H$  be a closed subgroup of the complex linear algebraic group  $G$  and let  $\phi : H \hookrightarrow G$  denote the inclusion map.
- Let  $\mathcal{E}_H \rightarrow X$  be a toric principal  $H$ -bundle with admissible collection  $\{\rho_\sigma, P(\tau, \sigma)\}$ .
- Then the associated bundle  $\mathcal{E}_H \times_H G$  is a toric principal  $G$ -bundle with admissible collection  $\{\phi \circ \rho_\sigma, \phi(P(\tau, \sigma))\}$ .

Let  $\mathcal{E}_G$  be a toric principal  $G$ -bundle over  $X$ .  $\mathcal{E}_G$  is said to admit an **equivariant reduction of structure group** to  $H \leq G$  if there exists a toric principal  $H$ -bundle  $\mathcal{E}_H$  such that the toric principal  $G$ -bundle  $\mathcal{E}_H \times_H G$  is equivariantly isomorphic to  $\mathcal{E}_G$ .

## Equivariant Levi reduction

Let  $G$  be reductive. A Levi subgroup of  $G$  is the centralizer in  $G$  of some torus in  $G$ .

## Equivariant Levi reduction

Let  $G$  be reductive. A Levi subgroup of  $G$  is the centralizer in  $G$  of some torus in  $G$ .

Let  $G = \mathrm{GL}(n, \mathbb{C})$ ,  $m \leq n$ .

$$T = \left\{ \begin{pmatrix} * & & & & \\ & \ddots & & & \\ & & * & & \\ & & & 1 & \\ & & & & \ddots \\ & & & & & 1 \end{pmatrix} \right\}, \quad C_G(T) = \left\{ \left( \begin{array}{ccc|c} * & & & 0 \\ & \ddots & & \vdots \\ & & * & 0 \\ \hline 0 & \dots & 0 & A \end{array} \right) \right\}.$$

## Equivariant Levi reduction

Let  $G$  be reductive. A Levi subgroup of  $G$  is the centralizer in  $G$  of some torus in  $G$ .

Let  $G = \mathrm{GL}(n, \mathbb{C})$ ,  $m \leq n$ .

$$T = \left\{ \begin{pmatrix} * & & & & \\ & \ddots & & & \\ & & * & & \\ & & & 1 & \\ & & & & \ddots \\ & & & & & 1 \end{pmatrix} \right\}, \quad C_G(T) = \left\{ \left( \begin{array}{ccc|c} * & & & 0 \\ & \ddots & & \vdots \\ & & * & 0 \\ \hline 0 & \dots & 0 & A \end{array} \right) \right\}.$$

**Theorem 3** (D, KHAN, BISWAS, DEY, PODDAR (2021))

$\mathcal{E}_G$  has an equivariant reduction of structure group to a Levi subgroup  $H$  of  $G$  if and only if

$$Z^0(H) \subseteq \mathrm{Aut}_T(\mathcal{E}_G).$$

## Equivariant splitting of a toric principal bundle

We say that  $\mathcal{E}_G$  splits equivariantly if it admits an equivariant reduction of structure group to a torus subgroup of  $G$ .



## Equivariant splitting of a toric principal bundle

We say that  $\mathcal{E}_G$  splits equivariantly if it admits an equivariant reduction of structure group to a torus subgroup of  $G$ .

**Theorem 4** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*Let  $\phi : G \rightarrow G'$  be an injective homomorphism of reductive algebraic groups. Let  $\mathcal{E}_{G'} := \mathcal{E}_G \times_G G'$ , where  $\mathcal{E}_G$  is a toric principal  $G$ -bundle on  $X$ . Suppose that  $\mathcal{E}_{G'}$  is equivariantly split, then  $\mathcal{E}_G$  itself splits equivariantly.*

## Principal bundle analogue of Kaneyama's result

**Theorem 5** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*Let  $G$  be a reductive subgroup of  $\mathrm{GL}(r, \mathbb{C})$ . Any toric principal  $G$ -bundle on  $\mathbb{P}^n$  splits equivariantly if  $r < n$ .*

## Principal bundle analogue of Kaneyama's result

**Theorem 5** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*Let  $G$  be a reductive subgroup of  $\mathrm{GL}(r, \mathbb{C})$ . Any toric principal  $G$ -bundle on  $\mathbb{P}^n$  splits equivariantly if  $r < n$ .*

**PROOF.**

Let  $\mathcal{E}_G$  be a  $T$ -equivariant principal  $G$ -bundle on  $\mathbb{P}^n$ .

## Principal bundle analogue of Kaneyama's result

**Theorem 5** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*Let  $G$  be a reductive subgroup of  $GL(r, \mathbb{C})$ . Any toric principal  $G$ -bundle on  $\mathbb{P}^n$  splits equivariantly if  $r < n$ .*

**PROOF.**

Let  $\mathcal{E}_G$  be a  $T$ -equivariant principal  $G$ -bundle on  $\mathbb{P}^n$ . Then we have that  $\mathcal{E}_G \times_G GL(r, \mathbb{C})$  splits equivariantly since  $r < n$  by Kaneyama's result.

## Principal bundle analogue of Kaneyama's result

**Theorem 5** (D, KHAN, BISWAS, DEY, PODDAR (2021))

*Let  $G$  be a reductive subgroup of  $GL(r, \mathbb{C})$ . Any toric principal  $G$ -bundle on  $\mathbb{P}^n$  splits equivariantly if  $r < n$ .*

**PROOF.**

Let  $\mathcal{E}_G$  be a  $T$ -equivariant principal  $G$ -bundle on  $\mathbb{P}^n$ . Then we have that  $\mathcal{E}_G \times_G GL(r, \mathbb{C})$  splits equivariantly since  $r < n$  by Kaneyama's result. Then by Theorem 4,  $\mathcal{E}_G$  splits equivariantly. □

## Principal bundle analogue of Kaneyama's result

**Theorem 5** (D, KHAN, BISWAS, DEY, PODDAR (2021))


*Let  $G$  be a reductive subgroup of  $GL(r, \mathbb{C})$ . Any toric principal  $G$ -bundle on  $\mathbb{P}^n$  splits equivariantly if  $r < n$ .*

**PROOF.**

Let  $\mathcal{E}_G$  be a  $T$ -equivariant principal  $G$ -bundle on  $\mathbb{P}^n$ . Then we have that  $\mathcal{E}_G \times_G GL(r, \mathbb{C})$  splits equivariantly since  $r < n$  by Kaneyama's result. Then by Theorem 4,  $\mathcal{E}_G$  splits equivariantly. □

This theorem has an alternative proof using certain results of Biswas and Parameswaran.

# Why study (semi)stability



- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .



## Stability of vector bundles

- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .
- $\mathcal{E}$  vector bundle on  $X$ .

## Stability of vector bundles

- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .
- $\mathcal{E}$  vector bundle on  $X$ .
- $\deg \mathcal{E} := c_1(\mathcal{E}) \cdot H^{n-1}$ , slope  $\mu(\mathcal{E}) := \frac{\deg \mathcal{E}}{\text{rank } \mathcal{E}}$ .

## Stability of vector bundles

- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .
- $\mathcal{E}$  vector bundle on  $X$ .
- $\deg \mathcal{E} := c_1(\mathcal{E}) \cdot H^{n-1}$ , slope  $\mu(\mathcal{E}) := \frac{\deg \mathcal{E}}{\text{rank } \mathcal{E}}$ .

### DEFINITION 3 ( $\mu$ -STABILITY OR MUMFORD-TAKEMOTO STABILITY)

$\mathcal{E}$  is said to be **(semi)stable** if for any coherent subsheaf  $\mathcal{F}$  of  $\mathcal{E}$  with  $0 < \text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ , we have  $\mu(\mathcal{F})(\leq) < \mu(\mathcal{E})$ .

## Stability of vector bundles

- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .
- $\mathcal{E}$  vector bundle on  $X$ .
- $\deg \mathcal{E} := c_1(\mathcal{E}) \cdot H^{n-1}$ , slope  $\mu(\mathcal{E}) := \frac{\deg \mathcal{E}}{\text{rank } \mathcal{E}}$ .

### DEFINITION 3 ( $\mu$ -STABILITY OR MUMFORD-TAKEMOTO STABILITY)

$\mathcal{E}$  is said to be **(semi)stable** if for any coherent subsheaf  $\mathcal{F}$  of  $\mathcal{E}$  with  $0 < \text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ , we have  $\mu(\mathcal{F})(\leq) < \mu(\mathcal{E})$ .

### EXAMPLE 4

- *Line bundles are stable.*

## Stability of vector bundles

- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .
- $\mathcal{E}$  vector bundle on  $X$ .
- $\deg \mathcal{E} := c_1(\mathcal{E}) \cdot H^{n-1}$ , slope  $\mu(\mathcal{E}) := \frac{\deg \mathcal{E}}{\text{rank } \mathcal{E}}$ .

### DEFINITION 3 ( $\mu$ -STABILITY OR MUMFORD-TAKEMOTO STABILITY)

$\mathcal{E}$  is said to be **(semi)stable** if for any coherent subsheaf  $\mathcal{F}$  of  $\mathcal{E}$  with  $0 < \text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ , we have  $\mu(\mathcal{F})(\leq) < \mu(\mathcal{E})$ .

### EXAMPLE 4

- *Line bundles are stable.*
- *Tensor product of semistable bundles is again semistable.*

## Stability of vector bundles

- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .
- $\mathcal{E}$  vector bundle on  $X$ .
- $\deg \mathcal{E} := c_1(\mathcal{E}) \cdot H^{n-1}$ , slope  $\mu(\mathcal{E}) := \frac{\deg \mathcal{E}}{\text{rank } \mathcal{E}}$ .

### DEFINITION 3 ( $\mu$ -STABILITY OR MUMFORD-TAKEMOTO STABILITY)

$\mathcal{E}$  is said to be **(semi)stable** if for any coherent subsheaf  $\mathcal{F}$  of  $\mathcal{E}$  with  $0 < \text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ , we have  $\mu(\mathcal{F})(\leq) < \mu(\mathcal{E})$ .

### EXAMPLE 4

- *Line bundles are stable.*
- *Tensor product of semistable bundles is again semistable.*
- *Dual of a semistable bundle is semistable.*

## Stability of vector bundles

- $(X, H)$  be a polarized nonsingular projective variety of dimension  $n$ .
- $\mathcal{E}$  vector bundle on  $X$ .
- $\deg \mathcal{E} := c_1(\mathcal{E}) \cdot H^{n-1}$ , slope  $\mu(\mathcal{E}) := \frac{\deg \mathcal{E}}{\text{rank } \mathcal{E}}$ .

### DEFINITION 3 ( $\mu$ -STABILITY OR MUMFORD-TAKEMOTO STABILITY)

$\mathcal{E}$  is said to be **(semi)stable** if for any coherent subsheaf  $\mathcal{F}$  of  $\mathcal{E}$  with  $0 < \text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ , we have  $\mu(\mathcal{F})(\leq) < \mu(\mathcal{E})$ .

### EXAMPLE 4

- *Line bundles are stable.*
- *Tensor product of semistable bundles is again semistable.*
- *Dual of a semistable bundle is semistable.*
- *Tangent bundle of projective space is stable.*

## Equivariant stability of toric vector bundles

### DEFINITION 5

An equivariant vector bundle  $\mathcal{E}$  on a toric variety  $X$  is said to be **equivariantly (semi)stable** with respect to an equivariant ample line bundle  $H$  if for any equivariant coherent subsheaf  $\mathcal{F}$  of  $\mathcal{E}$  with  $0 < \text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ , we have  $\mu(\mathcal{F})(\leq) < \mu(\mathcal{E})$ .



## Equivariant stability of toric vector bundles

### DEFINITION 5

An equivariant vector bundle  $\mathcal{E}$  on a toric variety  $X$  is said to be **equivariantly (semi)stable** with respect to an equivariant ample line bundle  $H$  if for any equivariant coherent subsheaf  $\mathcal{F}$  of  $\mathcal{E}$  with  $0 < \text{rank } \mathcal{F} < \text{rank } \mathcal{E}$ , we have  $\mu(\mathcal{F})(\leq) < \mu(\mathcal{E})$ .

### Theorem 6 ( KOOL (2011), BISWAS, DEY, GENÇ, AND PODDAR (2018))

Let  $\mathcal{E}$  be an equivariant vector bundle on a nonsingular projective toric variety  $X$ . Then  $\mathcal{E}$  is (semi)stable if and only if it is equivariantly (semi)stable.

## Stability of toric principal bundles

### DEFINITION 6

An equivariant principal  $G$ -bundle  $\mathcal{E}_G$  over a toric variety  $X$  is called **equivariantly semistable** (respectively, **equivariantly stable**) if

## Stability of toric principal bundles

### DEFINITION 6

An equivariant principal  $G$ -bundle  $\mathcal{E}_G$  over a toric variety  $X$  is called **equivariantly semistable** (respectively, **equivariantly stable**) if

- for every proper parabolic subgroup  $P \subsetneq G$  and

## Stability of toric principal bundles

### DEFINITION 6

An equivariant principal  $G$ -bundle  $\mathcal{E}_G$  over a toric variety  $X$  is called **equivariantly semistable** (respectively, **equivariantly stable**) if

- for every proper parabolic subgroup  $P \subsetneq G$  and
- every equivariant reduction  $\mathcal{E}_P$  of  $\mathcal{E}_G$  over any  $T$ -invariant open subset  $U \subset X$  with  $\text{codim}(X \setminus U) \geq 2$ , and

## Stability of toric principal bundles

### DEFINITION 6

An equivariant principal  $G$ -bundle  $\mathcal{E}_G$  over a toric variety  $X$  is called **equivariantly semistable** (respectively, **equivariantly stable**) if

- for every proper parabolic subgroup  $P \subsetneq G$  and
- every equivariant reduction  $\mathcal{E}_P$  of  $\mathcal{E}_G$  over any  $T$ -invariant open subset  $U \subset X$  with  $\text{codim}(X \setminus U) \geq 2$ , and
- for every nontrivial character  $\chi : P \longrightarrow \mathbb{C}^*$  which is dominant with respect to some Borel subgroup of  $G$  contained in  $P$ ,

## Stability of toric principal bundles

### DEFINITION 6

An equivariant principal  $G$ -bundle  $\mathcal{E}_G$  over a toric variety  $X$  is called **equivariantly semistable** (respectively, **equivariantly stable**) if

- for every proper parabolic subgroup  $P \subsetneq G$  and
- every equivariant reduction  $\mathcal{E}_P$  of  $\mathcal{E}_G$  over any  $T$ -invariant open subset  $U \subset X$  with  $\text{codim}(X \setminus U) \geq 2$ , and
- for every nontrivial character  $\chi : P \longrightarrow \mathbb{C}^*$  which is dominant with respect to some Borel subgroup of  $G$  contained in  $P$ ,

the associated line bundle  $\mathcal{E}_P(\chi)$  has non-positive (respectively, negative) degree.

## Stability of toric principal bundles

### DEFINITION 6

An equivariant principal  $G$ -bundle  $\mathcal{E}_G$  over a toric variety  $X$  is called **equivariantly semistable** (respectively, **equivariantly stable**) if

- for every proper parabolic subgroup  $P \subsetneq G$  and
- every equivariant reduction  $\mathcal{E}_P$  of  $\mathcal{E}_G$  over any  $T$ -invariant open subset  $U \subset X$  with  $\text{codim}(X \setminus U) \geq 2$ , and
- for every nontrivial character  $\chi : P \longrightarrow \mathbb{C}^*$  which is dominant with respect to some Borel subgroup of  $G$  contained in  $P$ ,

the associated line bundle  $\mathcal{E}_P(\chi)$  has non-positive (respectively, negative) degree.

### Theorem 7 (D, KHAN, BISWAS, DEY, PODDAR (2021))

Let  $\mathcal{E}_G$  be a  $T$ -equivariant principal  $G$ -bundle on a projective toric variety  $X$ . Then  $\mathcal{E}_G$  is stable if and only if  $\mathcal{E}_G$  is equivariantly stable.



Thank  
You