

Triangle Singularities in a Hilbert's House

CD24, Bochum

Ajay S. Sakthivasan ¹

M. Döring ^{2,3} M. Mai ^{2,4} A. Rusetsky ^{1,5}

¹HISKP & BCTP, University of Bonn

²INS & Department of Physics, GWU

³Thomas Jefferson National Accelerator Facility

⁴AEC, University of Bern

⁵Tbilisi State University


August 26, 2024

Presentation Outline

- 1 Introduction
- 2 Analytic Aspects
- 3 Infinite Volume Unitarity Formalism
- 4 Conclusions

Introduction

- Hadronic resonances correspond to poles of the S -matrix.




Pole position of the $a_1(1260)$.¹

¹Figure from: 10.1103/PhysRevD.105.054020

Introduction

- Hadronic resonances correspond to poles of the S -matrix.
- Experimentally — peaks in invariant mass distributions.




Pole position of the $a_1(1260)$.¹

¹Figure from: [10.1103/PhysRevD.105.054020](https://doi.org/10.1103/PhysRevD.105.054020)

Introduction

- Hadronic resonances correspond to poles of the S -matrix.
- Experimentally — peaks in invariant mass distributions.
- All observed peaks correspond to hadronic states? **No.**



Pole position of the $a_1(1260)$.¹

¹Figure from: 10.1103/PhysRevD.105.054020

- **Landau Singularities** may not correspond to poles of the S -matrix, but can *mimic* a resonance.


- **Landau Singularities** may not correspond to poles of the S -matrix, but can *mimic* a resonance.
- Originally proposed by Landau. Recently used to explain a lot of observed peaks (*one-loop level, rescattering effects not taken into account*).

- **Landau Singularities** may not correspond to poles of the S -matrix, but can *mimic* a resonance.
- Originally proposed by Landau. Recently used to explain a lot of observed peaks (*one-loop level, rescattering effects not taken into account*).
- We try to achieve the following:

- Landau Singularities may not correspond to poles of the S -matrix, but can *mimic* a resonance.
- Originally proposed by Landau. Recently used to explain a lot of observed peaks (*one-loop level, rescattering effects not taken into account*).
- We try to achieve the following:
 - ① To systematically study the effect of final-state rescattering.


- Landau Singularities may not correspond to poles of the S -matrix, but can *mimic* a resonance.
- Originally proposed by Landau. Recently used to explain a lot of observed peaks (*one-loop level, rescattering effects not taken into account*).
- We try to achieve the following:
 - ① To systematically study the effect of final-state rescattering.
 - ② To consistently describe the unstable particles.

- Landau Singularities may not correspond to poles of the S -matrix, but can *mimic* a resonance.
- Originally proposed by Landau. Recently used to explain a lot of observed peaks (*one-loop level, rescattering effects not taken into account*).
- We try to achieve the following:
 - ① To systematically study the effect of final-state rescattering.
 - ② To consistently describe the unstable particles.
 - ③ Apply this to an $a_1(1420)$ (no spin-)system, and eventually the $a_1(1420)$ system with spin.²




²Figure from: 10.1103/physrevlett.127.082501

- Landau Singularities may not correspond to poles of the S -matrix, but can *mimic* a resonance.
- Originally proposed by Landau. Recently used to explain a lot of observed peaks (*one-loop level, rescattering effects not taken into account*).
- We try to achieve the following:
 - ① To systematically study the effect of final-state rescattering.
 - ② To consistently describe the unstable particles.
 - ③ **Apply this to an $a_1(1420)$ (no spin-)system**, and eventually the $a_1(1420)$ system with spin.²



²Figure from: 10.1103/physrevlett.127.082501

- Landau Singularities may not correspond to poles of the S -matrix, but can *mimic* a resonance.
- Originally proposed by Landau. Recently used to explain a lot of observed peaks (*one-loop level, rescattering effects not taken into account*).
- We try to achieve the following:
 - ① To systematically study the effect of final-state rescattering.
 - ② To consistently describe the unstable particles.
 - ③ **Apply this to an $a_1(1420)$ (no spin-)system**, and eventually the $a_1(1420)$ system with spin.²



²Figure from: 10.1103/physrevlett.127.082501

- Effects of Final State Interactions on Landau singularities
(arXiv:2407.17969 [hep-ph]) — under review.

- Effects of Final State Interactions on Landau singularities
(arXiv:2407.17969 [hep-ph]) — under review.
- $a_1(1420)$: resonance like structure in COMPASS experiment.
(10.1103/PhysRevLett.115.082001)
- Interpretation as a kinematical singularity — Mikhasenko et al.
(10.1103/PhysRevD.91.094015), Bayar et al.
(10.1103/PhysRevD.94.074039)
- Relevance to lattice QCD and ChPT — Korpa et al.
(10.1103/PhysRevD.107.L031505), Isken et al.
(10.1103/PhysRevD.109.034032), Mai et al.
(10.1103/PhysRevLett.127.222001)
- Review — Guo et al. (10.1016/j.ppnp.2020.103757)

The Landau Equations

- **The Landau Equations:** singularities corresponding to Feynman integrals (*Landau singularities*).

The Landau Equations

- **The Landau Equations:** singularities corresponding to Feynman integrals (*Landau singularities*).
- For the Feynman integral with N internal propagators and l loop momenta,

$$I_1 = \int \prod_{i=1}^l d^4 k_i \frac{1}{\prod_{r=1}^N (q_r(k_i)^2 - m_r^2)}.$$

The Landau Equations

- **The Landau Equations:** singularities corresponding to Feynman integrals (*Landau singularities*).
- For the Feynman integral with N internal propagators and l loop momenta,

$$I_l = \int \prod_{i=1}^l d^4 k_i \frac{1}{\prod_{r=1}^N (q_r(k_i)^2 - m_r^2)}.$$

- For some real parameters α_i , the Landau equations read,


$$\alpha_i(q_i^2 - m_i^2) = 0 \implies \alpha_i = 0 \text{ or } q_i^2 = m_i^2,$$

$$\sum_{i \in \text{loop}} \alpha_i q_i^\mu(k_j) = 0.$$


- $\alpha_i \neq 0 \implies$ all propagators go on-shell for the **leading singularity**.
One also needs $\alpha_i > 0$.

- $\alpha_i \neq 0 \implies$ all propagators go on-shell for the **leading singularity**.
One also needs $\alpha_i > 0$.
- $\alpha_i = 0$ case is called the **subleading singularity** — equivalent to contracting the propagator.

- The leading Landau singularity associated with the graph:




- The leading Landau singularity associated with the graph:




- Landau singularity for $p_3 = m_\pi$: $p_2/\text{GeV} \equiv \sqrt{\sigma}/\text{GeV} \in [0.986, 1.024]$ and $p_1/\text{GeV} \equiv \sqrt{s}/\text{GeV} \in [1.385, 1.436]$.

- What happens to singularities after rescattering? **Ladder diagrams.**

- What happens to singularities after rescattering? **Ladder diagrams.**
- Triangle +1 box diagram:



- What happens to singularities after rescattering? **Ladder diagrams.**
- Triangle +1 box diagram:




- Check for leading and subleading singularities. . .


- Leading singularity **is not present.**

- Leading singularity **is not present**.
- For subleading singularities go through all contractions and check for singular graphs...


- Leading singularity is not present.
- For subleading singularities go through all contractions and check for singular graphs . . .



- Leading singularity is not present.
- For subleading singularities go through all contractions and check for singular graphs...




- Leading singularity is not present.
- For subleading singularities go through all contractions and check for singular graphs...



- Accommodating infinite rescatterings.

- Accommodating infinite rescatterings.
- The triangle singularity is present and no other singularities.



- Landau equations give the conditions for the singularity to be present.

- Landau equations give the conditions for the singularity to be present.
- One still needs to evaluate the corresponding Feynman integrals.


- Study the system using **Infinite Volume 3-Body Unitary Formalism**, due to Mai et al.³

³[10.1140/epja/i2017-12368-4](https://doi.org/10.1140/epja/i2017-12368-4)

- Study the system using **Infinite Volume 3-Body Unitary Formalism**, due to Mai et al.³
- **In brief:** the system is split into a two-body subsystem — the **isobar**, and the **spectator**, which is on-shell.

³[10.1140/epja/i2017-12368-4](https://doi.org/10.1140/epja/i2017-12368-4)


- Study the system using **Infinite Volume 3-Body Unitary Formalism**, due to Mai et al.³
- **In brief:** the system is split into a two-body subsystem — the **isobar**, and the **spectator**, which is on-shell.
- The full three-body amplitude is split into a **connected** and a **disconnected** amplitude.⁴




³[10.1140/epja/i2017-12368-4](https://doi.org/10.1140/epja/i2017-12368-4)

⁴Figure from: [10.1140/epja/i2017-12368-4](https://doi.org/10.1140/epja/i2017-12368-4)

- Study the system using **Infinite Volume 3-Body Unitary Formalism**, due to Mai et al.³
- **In brief:** the system is split into a two-body subsystem — the **isobar**, and the **spectator**, which is on-shell.
- The full three-body amplitude is split into a **connected** and a **disconnected** amplitude.⁴



- One considers the **Bethe-Salpeter ansatz**, $T = B + BGT$, to determine T .⁴



³[10.1140/epja/i2017-12368-4](https://doi.org/10.1140/epja/i2017-12368-4)

⁴Figure from: [10.1140/epja/i2017-12368-4](https://doi.org/10.1140/epja/i2017-12368-4)

- The equation to be solved,

$$T(p, q) = B(p, q) + \int \frac{d^3 l}{(2\pi)^3} \frac{1}{2E_l} B(p, l) \tau(\sigma(l)) T(l, q).$$

- The equation to be solved,

$$T(p, q) = B(p, q) + \int \frac{d^3 l}{(2\pi)^3} \frac{1}{2E_l} B(p, l) \tau(\sigma(l)) T(l, q).$$


- We have the explicit forms of B and τ .

- The equation to be solved,


$$T(p, q) = B(p, q) + \int \frac{d^3 l}{(2\pi)^3} \frac{1}{2E_l} B(p, l) \tau(\sigma(l)) T(l, q).$$

- We have the explicit forms of B and τ .
- In this work, we consider scalar propagators, in relative s -wave.

- Need to carry out a coupled channel analysis.




- Need to carry out a coupled channel analysis.




- B is parametrised through the spectator masses.

- Need to carry out a coupled channel analysis.




- B is parametrised through the spectator masses.
- There are two unknowns corresponding to each isobar — coupling and bare mass.

- Need to carry out a coupled channel analysis.



- B is parametrised through the spectator masses.
- There are two unknowns corresponding to each isobar — coupling and bare mass.
- These are fixed through PDG values of physical mass and width.

- Need to carry out a coupled channel analysis.




- B is parametrised through the spectator masses.
- There are two unknowns corresponding to each isobar — coupling and bare mass.
- These are fixed through PDG values of physical mass and width.
- For f_0 we also make use of g_K/g_ρ ratio from BaBar collaboration.


- The integral is solved numerically, using Gaussian quadrature.

- The integral is solved numerically, using Gaussian quadrature.
- But, singularities on the integration contour may prevent numerical implementation.

- The integral is solved numerically, using Gaussian quadrature.
- But, singularities on the integration contour may prevent numerical implementation.
- **Contour deformation:** Evaluate the integral along a complex contour, that is far from the singularities \rightarrow obtain $T(\sqrt{s}, q \in \mathbb{C})$.



- The integral is solved numerically, using Gaussian quadrature.
- But, singularities on the integration contour may prevent numerical implementation.
- **Contour deformation:** Evaluate the integral along a complex contour, that is far from the singularities \rightarrow obtain $T(\sqrt{s}, q \in \mathbb{C})$.



- But, we need the amplitudes for $q \in \mathbb{R} \rightarrow$ can be done in different ways!

- The solution of the Bethe-Salpeter equation can be given in form of the **Born series**.

$$\text{---} \circ T \text{---} = \text{---} \backslash B \text{---} + \text{---} \backslash B / B \text{---} + \dots$$


- The solution of the Bethe-Salpeter equation can be given in form of the **Born series**.

$$\text{Diagram } T = \text{Diagram } B + \text{Diagram } B \text{ (with loop)} + \dots$$


- This defines the **loop expansion** for the invariant mass distributions.

- **Rational Analytic Continuation:** We use a **continued fraction** to obtain the isobar-spectator amplitude for $q \in \mathbb{R}$.

- **Rational Analytic Continuation:** We use a **continued fraction** to obtain the isobar-spectator amplitude for $q \in \mathbb{R}$.
- At $1 + \infty$ -loop level,




Amplitude squared vs. $\sqrt{\sigma}$, for
 $\sqrt{s} = 1.42$ GeV.




Amplitude squared vs. \sqrt{s} , for
 $\sqrt{\sigma} = 0.99$ GeV.

- **Rational Analytic Continuation:** We use a **continued fraction** to obtain the isobar-spectator amplitude for $q \in \mathbb{R}$.
- At $1 + \infty$ -loop level,



Amplitude squared vs. $\sqrt{\sigma}$, for
 $\sqrt{s} = 1.42$ GeV.




Amplitude squared vs. \sqrt{s} , for
 $\sqrt{\sigma} = 0.99$ GeV.


- We see the Landau singularity in both the plots.

- **Cahill & Sloan Method:** We make use **Cauchy's integral theorem** instead.

- **Cahill & Sloan Method:** We make use **Cauchy's integral theorem** instead.
- Matches with the other method.




Amplitude squared vs. $\sqrt{\sigma}$, for
 $\sqrt{s} = 1.42$ GeV.




Amplitude squared vs. \sqrt{s} , for
 $\sqrt{\sigma} = 0.99$ GeV.

- Triangle diagram contributes the most to the amplitude.

- Triangle diagram contributes the most to the amplitude.
- With decreasing contributions from subsequent rescatterings.




Ratios of amplitudes to triangle amplitudes vs. $\sqrt{\sigma}$, for $\sqrt{s} = 1.42$ GeV.




Ratios of amplitudes to triangle amplitudes vs. $\sqrt{\sigma}$, for $\sqrt{s} = 0.99$ GeV.

- Triangle diagram contributes the most to the amplitude.
- With decreasing contributions from subsequent rescatterings.



Ratios of amplitudes to triangle amplitudes vs. $\sqrt{\sigma}$, for $\sqrt{s} = 1.42 \text{ GeV}$.



Ratios of amplitudes to triangle amplitudes vs. $\sqrt{\sigma}$, for $\sqrt{s} = 0.99 \text{ GeV}$.

- The amplitudes at triangle +1 box are about 1.1x the triangle diagram amplitudes.

Conclusions

- Triangle singularity at 1-loop level and at higher loops as **subleading singularity**.

Conclusions

- Triangle singularity at 1-loop level and at higher loops as **subleading singularity**.
- The isobar-spectator model numerically implemented and the triangle singularity observed at $1 + \infty$ -loop level.

Conclusions

- Triangle singularity at 1-loop level and at higher loops as **subleading singularity**.
- The isobar-spectator model numerically implemented and the triangle singularity observed at $1 + \infty$ -loop level.
- The singularity is of **logarithmic** nature at all levels.

Conclusions

- Triangle singularity at 1-loop level and at higher loops as **subleading singularity**.
- The isobar-spectator model numerically implemented and the triangle singularity observed at $1 + \infty$ -loop level.
- The singularity is of **logarithmic** nature at all levels.
- The contributions of the higher order diagrams depend on the system considered.

Conclusions

- Triangle singularity at 1-loop level and at higher loops as **subleading singularity**.
- The isobar-spectator model numerically implemented and the triangle singularity observed at $1 + \infty$ -loop level.
- The singularity is of **logarithmic** nature at all levels.
- The contributions of the higher order diagrams depend on the system considered.
- Outlook:
 - ① Realistic model, i.e., spin with pseudovector source — in progress.

Conclusions

- Triangle singularity at 1-loop level and at higher loops as **subleading singularity**.
- The isobar-spectator model numerically implemented and the triangle singularity observed at $1 + \infty$ -loop level.
- The singularity is of **logarithmic** nature at all levels.
- The contributions of the higher order diagrams depend on the system considered.
- Outlook:
 - ➊ Realistic model, i.e., spin with pseudovector source — in progress.
 - ➋ $a_1(1420)$ on the lattice? Implement finite volume unitarity⁵— in progress.


⁵[10.1140/epja/i2017-12440-1](https://doi.org/10.1140/epja/i2017-12440-1)

Thanks for listening!

Landau singularities


Endpoint singularities: singularities in g , hitting the contour C . E.g.,

$$f(z) = \int_1^2 dw \frac{1}{w-z}. \quad (1)$$




Pinch singularities: Contour C , gets trapped between two singularities in g . E.g.,

$$f(z) = \int_1^2 dw \frac{1}{(w-z)(w-5/2)}. \quad (2)$$



Pinch singularity avoided when the singularities approach from the same side of the contour.



The Other Method


- **Cahill & Sloan Method⁶:** Instead of a continued fraction, use the knowledge of the **location of the branch cuts** to analytically continue the amplitude for $q \in \mathbb{R}$.⁷

⁶[10.1016/0375-9474\(71\)90156-4](https://doi.org/10.1016/0375-9474(71)90156-4)


⁷For a comprehensive review: *The Quantum Mechanical Three-Body Problem*, Schmid & Ziegelmann.

The Other Method

- **Cahill & Sloan Method**⁶: Instead of a continued fraction, use the knowledge of the **location of the branch cuts** to analytically continue the amplitude for $q \in \mathbb{R}$.⁷



Branch cuts of B for q outside the triangle singularity region.



Branch cuts of B for q in the triangle singularity region.

⁶10.1016/0375-9474(71)90156-4

⁷For a comprehensive review: *The Quantum Mechanical Three-Body Problem*, Schmid & Ziegelmann.