Electromagnetic Form Factors from Lattice QCD

Ajay S. Sakthivasan[1], A. Rusetsky[1], J. J. Wu[2], U. G. Meißner[1], G. Schierholz[3]

[1]HISKP, Universität Bonn [2]CAS, Beijing [3]DESY, Hamburg

Based on JHEP 06 (2026) 164 [1]

1. Motivation

The calculation of hadronic form factors on the lattice provides important information about the internal structure of the corresponding states. However, **calculating the form factor for unstable states**, viz. resonances, is not as straightforward, both from a conceptual point of view and in terms of technical implementation. In this work, we specifically consider **the electromagnetic form factor of the ρ-meson**. We make use of **the external field method (the Feynman-Hellmann method)**, where the system is placed in an external field coupling to a spatially periodic current. The structure of the finite-volume energy levels is described by **the modified Lüscher approach**, which enables one to directly extract the short-range part of the interaction with the external field from the fit to the shifted energy levels. This method has the advantage that the separation of **the triangle diagram**, which has an irregular dependence on L, is automatically taken into account, and the short-range interaction has a regular dependence on L and exponentially suppressed for large L. Through this work, we provide **a complete framework for the extraction of the form factor from raw lattice data** when available. We, in addition, provide **a crude estimate of the short-range contribution to the form factor at the lowest order in ChPT**.

2. The Covariant Formulation of NREFT

We work with the the covariant formulation of the pion non-relativistic effective field theory, whose Lagrangian given by

$$\mathcal{L} = \overbrace{\sum_{a=+,0} \phi_a^\dagger 2W(i(v\cdot\partial) - W)\phi_a}^{\text{Pion Lagrangian}} - \overbrace{\Psi_\mu^\dagger \Psi^\mu}^{\text{Rho Lagrangian}}$$

$$W = \sqrt{M^2 + \partial^2 - (v\cdot\partial)^2} \quad \text{quantisation axis}$$

$$\Psi^\mu = \phi_+ \overset{\leftrightarrow}{W}{}^\mu f(\overset{\leftrightarrow}{W}{}^2)\phi_0,$$
$$W^\mu = v^\mu W + i(\partial^\mu - v^\mu(v\cdot\partial)),$$
$$\overset{\leftrightarrow}{W}{}^\mu = \frac{1}{2}(\vec{W}{}^\mu - \overset{\leftarrow}{W}{}^\mu)$$
$$f(\overset{\leftrightarrow}{W}{}^2) = f_0 + f_1 \overset{\leftrightarrow}{W}{}^2 + \dots$$

Note that the ρ Lagrangian is constructed using a dipion field. The quantity f is related to experimental data through

$$q\cot\delta = -\frac{24\pi\sqrt{P^2}}{q^2 f^2(q^2)}$$

2.1 Including the Electromagnetic Interaction

Gauging the Lagrangian is no longer straightforward due to the presence of non-linear operators. It can be seen that the gauging the quadratic operator leads to

$$W^2 \to W^2 - ie(\partial\cdot A + A\cdot\partial) + ie((v\cdot\partial)(v\cdot A) + (v\cdot A)(v\cdot\partial)) + \mathcal{O}(e^2)$$
$$\equiv W^2 + B + \mathcal{O}(e^2)$$

Then the non-linear operator can be gauged as


$$W \to W + D, \quad D = \int_0^\infty dt\, e^{-tW} B e^{tW}$$

However, the contribution of the 4-pion-photon vertex to the EM form factor of the ρ-meson from this procedure is not apparent. We therefore explicitly include the operators that lead to the relevant form factor structures.

2.2 Invariant Form Factors

We can now calculate the relevant form factors in the NREFT, and relate them to standard ChPT form factors:


$$\begin{aligned} G_1(k^2) &= -eZg_1k^2 + eZ\langle F_1\rangle \\ G_2(k^2) &= -eZg_2 + eZ\langle F_2\rangle \\ \frac{1}{2s_R}G_3(k^2) &= eZg_3 + eZ\langle F_3\rangle \end{aligned}$$



where Z is the sum of all the pion bubbles. All the other quantities above are also well-defined in the NREFT.

2.3 Estimates of Contact Term Contributions

The contributions arising from the contact terms can be estimated crudely using next-to-leading order chiral perturbation theory.



The numerical stability of this approach is tested with two different parametrisations for the two pion scattering amplitudes. The error bands show the estimated contributions from the contact terms.

3. The Modified Lüscher Approach

The two-pion is system is placed in a periodic electromagnetic field on a cubic lattice of size L. The two-point function is appropriately modified and it defines the energy spectrum of the two-pion system in a background field. The quantisation condition can also be derived:

$$\det(\Pi^{-1}_{\mu\nu} - D^0_{\mu\nu}) = 0$$

"self-energy" the two-point function of the interpolating field

which determines the position of the poles. The interpolating field is related to the dipion field. Now projecting this on to the relevant irreps and specific background field configurations, leads to relations involving g_1, g_2 and g_3, which can be fit to lattice data. This lets us extract the electromagnetic form factors.

4. Conclusions

The formalism developed here establishes a **consistent way to extract the electromagnetic structure of unstable resonances directly from finite-volume spectra**. In the particular case of the ρ-meson, by combining **the external-field approach** with **the modified Lüscher equation**, we isolate the short-range contribution to the ρ-meson form factors and remove the pathological finite-volume effects originating from the triangle diagram. Additionally, this framework also gives an **estimate of the size of the couplings starting from ChPT**, which is useful before running simulations. Therefore, this framework lays the groundwork for future lattice studies where the extraction of resonance form factors can be performed without resorting to model assumptions or ad hoc subtractions, enabling a clean comparison with phenomenology once lattice data become available.

References

[1] Ulf-G. Meißner et al. "Form factors of the ρ meson from effective field theory and the lattice". In: *JHEP* 06 (2026), p. 164. DOI: 10.1007/JHEP06(2026)164. arXiv: 2602.23044 [hep-lat].

[2] J. Gasser et al. "Hadronic atoms in QCD + QED". In: *Phys. Rept.* 456 (2008), pp. 167–251. DOI: 10.1016/j.physrep.2007.09.006. arXiv: 0711.3522 [hep-ph].

[3] D. Hoja et al. "Resonances in an external field: The 1+1 dimensional case". In: *JHEP* 04 (2010), p. 050. DOI: 10.1007/JHEP04(2010)050. arXiv: 1001.1641 [hep-lat].

[4] V. Bernard et al. "Matrix elements of unstable states". In: *JHEP* 09 (2012), p. 023. DOI: 10.1007/JHEP09(2012)023. arXiv: 1205.4642 [hep-lat].

[5] Fabian Müller et al. "Relativistic-invariant formulation of the NREFT three-particle quantization condition". In: *JHEP* 02 (2022), p. 158. DOI: 10.1007/JHEP02(2022)158. arXiv: 2110.09351 [hep-lat].

[6] Jonathan Lozano et al. "Resonance form factors from finite-volume correlation functions with the external field method". In: *JHEP* 10 (2022), p. 106. DOI: 10.1007/JHEP10(2022)106. arXiv: 2205.11316 [hep-lat].

[7] Fabian Müller et al. "Three-particle Lellouch-Lüscher formalism in moving frames". In: *JHEP* 02 (2023), p. 214. DOI: 10.1007/JHEP02(2023)214. arXiv: 2211.10126 [hep-lat].