

The **AMMIS**tb**P** package: A brief introduction

Ajay B. C.¹, J. Aravind² and R. Abdul Fiyaz³

2018-07-12

1. RRS, ICAR-Directorate of Groundnut Research, Anantapur.
 2. ICAR-National Bureau of Plant Genetic Resources, New Delhi.
 3. ICAR-Indian Institute of Rice Research, Hyderabad.
-

Contents

Installation	1
AMMI	1
AMMI stability parameters	2
Yield Stability Index	6
Citing AMMIS tb P	6
Session Info	6
References	7

The package AMMIS**tb**P is

Installation

The package can be installed using the following functions:

```
# Install from CRAN
install.packages('AMMIStbP', dependencies=TRUE)

# Install development version from Github
devtools::install_github("ajaygpb/AMMIStbP")
```

Then the package can be loaded using the function

```
library(AMMIStbP) # change eval
```

AMMI

The AMMI equation

$$Y_{ij} = \mu + \alpha_i + \beta_j + \sum_{n=1}^N \lambda_n \gamma_{in} \delta_{jn} + \rho_{ij}$$

Where, Y_{ij} is the yield of i th genotype in j th environment, μ is the grand mean, α_i is the genotype deviation from the grand mean, β_j is the environment deviation, N is the total number of interaction principal components (IPCs), λ_n is the singular value for IPC n and correspondingly λ_n^2 is its eigen value, γ_{in} is the eigenvector value for i th genotype, δ_{jn} is the eigenvector value for j th environment and ρ_{ij} is the residual.

AMMI stability parameters

The details about AMMI stability parameters/indices implemented in **AMMIS**tbP**** are described in Table 1.

Table 1 : AMMI stability parameters/indices implemented in AMMISbP.

AMMI stability parameter	function	Details	Reference
Sums of the absolute value of the IPC scores (<i>SIPC</i>)	<code>SIPC.AMMI</code>	$SIPC = \sum_{n=1}^{N'} \lambda_n^{0.5} \gamma_{in} $ $SIPC = \sum_{n=1}^{N'} PC_n $	Sneller et al. (1997)
Averages of the squared eigenvector values <i>EV</i>	<code>EV.AMMI</code>	$EV = \sum_{n=1}^{N'} \frac{\gamma_{in}^2}{N'}$	Zobel (1994)
Sum across environments of GEI modelled by AMMI (<i>AMGE</i>)	<code>AMGE.AMMI</code>	$AMGE = \sum_{j=1}^E \sum_{n=1}^{N'} \lambda_n \gamma_{in} \delta_{jn}$	Sneller et al. (1997)
$AV_{(AMGE)}$	<code>AVAMGE.AMMI</code>	$AV_{(AMGE)} = \sum_{j=1}^E \sum_{n=1}^{N'} \lambda_n \gamma_{in} \delta_{jn} $	Zali et al. (2012)
Annicchiarico's D parameter (D_a)	<code>DA.AMMI</code>	The unsquared Euclidean distance from the origin of significant IPC axes in the AMMI model.	Annicchiarico (1997)
		$D_a = \sqrt{\sum_{n=1}^{N'} (\lambda_n \gamma_{in})^2}$	
Zhang's D parameter or AMMI statistic coefficient or AMMI distance or AMMI stability index (D_z)	<code>DZ.AMMI</code>	The distance of IPC point from origin in space.	Zhang et al. (1998)
		$D_z = \sqrt{\sum_{n=1}^{N'} \gamma_{in}^2}$	

AMMI stability parameter	function	Details	Reference
AMMI stability value (ASV)	<code>agricolae::index.AMMI</code>	Distance from the coordinate point to the origin in a two dimensional scattergram generated by plotting of IPC1 score against IPC2 score.	Purchase (1997); Purchase et al. (1999); Purchase et al. (2000)
Modified AMMI stability value (ASV)	<code>MASV.AMMI</code>	$MASV = \sqrt{\left(\frac{SSIPC_1}{SSIPC_2} \times PC_1\right)^2 + (PC_2)^2}$	Zali et al. (2012)
Absolute value of the relative contribution of IPCs to the interaction Za	<code>ZA.AMMI</code>	$Za = \sum_{i=1}^{N'} \theta_n \gamma_{in} $	Zali et al. (2012)
Stability measure based on fitted AMMI model FA	<code>FA.AMMI</code>	$FA = \sum_{n=1}^{N'} \lambda_n^2 \gamma_{in}^2$	Raju (2002); Zali et al. (2012)
FP	NA	Equivalent to FA , when only the first IPC axis is considered for computation.	Raju (2002); Zali et al. (2012)

$$FP = \lambda_1^2 \gamma_{i1}^2$$

As λ_1^2 will be same for all the genotypes, the absolute value of γ_{i1} alone is sufficient for comparison. So this is also equivalent to the comparison based on biplot with first IPC axis.

AMMI stability parameter	function	Details	Reference
B	NA	Equivalent to FA , when the first two IPC axes are considered for computation. $B = \sum_{n=1}^2 \lambda_n^2 \gamma_{in}^2$ Stability comparisons based on this measure will be equivalent to the comparisons based on biplot with first two IPC axes.	Raju (2002); Zali et al. (2012)
$W_{(AMMI)}$	NA	Equivalent to FA , when all the IPC axes in the AMMI model are considered for computation. $W_{(AMMI)} = \sum_{n=1}^N \lambda_n^2 \gamma_{in}^2$ Equivalent to Wricke's ecovalence.	Wricke (1962); Raju (2002); Zali et al. (2012)
AMMI Stability Index (ASI)	ASI.AMMI	$ASI = \sqrt{[PC_1^2 \times \theta_1^2] + [PC_2^2 \times \theta_2^2]}$	Jambhulkar et al. (2014); Jambhulkar et al. (2015); Jambhulkar et al. (2017)

Where, N is the total number of interaction principal components (IPCs); N' is the number of significant IPCAs (number of IPC that were retained in the AMMI model via F tests); λ_n is the singular value for IPC n and correspondingly λ_n^2 is its eigen value; γ_{in} is the eigenvector value for i th genotype; δ_{jn} is the eigenvector value for j th environment; $SSIPC_1, SSIPC_2, \dots, SSIPC_n$ are the sum of squares of the 1st, 2th, \dots , and n th IPC; PC_1, PC_2, \dots, PC_n are the scores of 1st, 2th, \dots , and n th IPC; θ_n is the percentage sum of squares explained by n th principal component interaction effect; and E is the number of environments.

Yield Stability Index

The most stable genotype need not necessarily be the high yielding genotype. As we need to select the the most stable and highest yielding genotypes, Yield stability index (*YSI*) was proposed (Farshadfar et al. (2011), Jambhulkar et al. (2017)). *YSI* is a simultaneous selection index for yield and yield stability which is computed by summation of the ranks of the stability index/parameter and the ranks of the mean yields. *YSI* is computed for all the stability parameters/indices implemented in this package.

$$YSI = R_{SP} + R_Y$$

Where, R_{SP} is the stability parameter/index rank of the genotype and R_Y is the mean yield rank of the genotype.

Examples

Citing AMMISbP

Session Info

```
sessionInfo()
```

```
R version 3.5.1 (2018-07-02)
```

```
Platform: x86_64-w64-mingw32/x64 (64-bit)
```

```
Running under: Windows >= 8 x64 (build 9200)
```

```
Matrix products: default
```

```
locale:
```

```
[1] LC_COLLATE=English_India.1252 LC_CTYPE=English_India.1252
```

```
[3] LC_MONETARY=English_India.1252 LC_NUMERIC=C
```

```
[5] LC_TIME=English_India.1252
```

```
attached base packages:
```

```
[1] stats      graphics  grDevices  utils      datasets  methods    base
```

```
other attached packages:
```

```
[1] readxl_1.1.0      AMMISbP_0.0.0.9000
```

```
loaded via a namespace (and not attached):
```

```
[1] gtools_3.5.0      pander_0.6.1      splines_3.5.1
[4] lattice_0.20-35   agricolae_1.2-8    expm_0.999-2
[7] miniUI_0.1.1      htmltools_0.3.6    yaml_2.1.19
[10] rlang_0.2.1        AlgDesign_1.1-7.3  pillar_1.2.2
[13] later_0.7.1        withr_2.1.2        sp_1.2-7
[16] questionr_0.6.2    stringr_1.3.0      combinat_0.0-8
[19] commonmark_1.5     cellranger_1.1.0   devtools_1.13.5
[22] coda_0.19-1        memoise_1.1.0      evaluate_0.10.1
[25] knitr_1.20         httpuv_1.4.1       gbRd_0.4-11
[28] spdep_0.7-7        highr_0.6          Rcpp_0.12.16
[31] xtable_1.8-2       promises_1.0.1     backports_1.1.2
[34] gdata_2.18.0       desc_1.2.0         mime_0.5
[37] deldir_0.1-15      klaR_0.6-14        digest_0.6.15
```

[40] stringi_1.1.7	gmodels_2.16.2	shiny_1.0.5
[43] grid_3.5.1	rprojroot_1.3-2	bibtex_0.4.2
[46] Rdpack_0.7-0	tools_3.5.1	LearnBayes_2.15.1
[49] magrittr_1.5	tibble_1.4.2	cluster_2.0.7-1
[52] crayon_1.3.4	MASS_7.3-50	Matrix_1.2-14
[55] xml2_1.2.0	spData_0.2.8.3	assertthat_0.2.0
[58] rmarkdown_1.10	roxygen2_6.0.1	rstudioapi_0.7.0-9000
[61] R6_2.2.2	boot_1.3-20	nlme_3.1-137
[64] compiler_3.5.1		

References

- Annicchiarico, P. (1997). Joint regression vs AMMI analysis of genotype-environment interactions for cereals in Italy. *Euphytica* 94, 53–62. doi:[10.1023/A:1002954824178](https://doi.org/10.1023/A:1002954824178).
- Farshadfar, E., Mahmodi, N., and Yaghotipoor, A. (2011). AMMI stability value and simultaneous estimation of yield and yield stability in bread wheat (*Triticum aestivum* L.). *Australian Journal of Crop Science* 5, 1837–1844.
- Jambhulkar, N. N., Bose, L. K., Pande, K., and Singh, O. N. (2015). Genotype by environment interaction and stability analysis in rice genotypes. *Ecology, Environment and Conservation* 21, 1427–1430. Available at: http://www.envirobiotechjournals.com/article_abstract.php?aid=6346&iid=200&jid=3.
- Jambhulkar, N. N., Bose, L. K., and Singh, O. N. (2014). “AMMI Stability Index for Stability Analysis,” in *CRRI Newsletter, January-March 2014*, ed. T. Mohapatra (Cuttack, Orissa: Central Rice Research Institute), 15. Available at: http://www.crrri.nic.in/CRRI_newsletter/crnl_jan_mar_14_web.pdf.
- Jambhulkar, N., Rath, N., Bose, L., Subudhi, H., Biswajit, M., Lipi, D., et al. (2017). Stability analysis for grain yield in rice in demonstrations conducted during rabi season in India. *Oryza* 54, 236–240. doi:[10.5958/2249-5266.2017.00030.3](https://doi.org/10.5958/2249-5266.2017.00030.3).
- Purchase, J. (1997). Parametric Analysis to Describe Genotype \times Environment Interaction and Yield Stability in Winter Wheat. Available at: <http://hdl.handle.net/11660/1966>.
- Purchase, J., Hatting, H., and Van Deventer, C. (1999). “The use of the AMMI model and AMMI stability value to describe genotype \times environment interaction and yield stability in winter wheat (*Triticum aestivum* L.),” in *Proceedings of the Tenth Regional Wheat Workshop for Eastern, Central and Southern Africa, 14-18 September 1998* (South Africa: University of Stellenbosch).
- Purchase, J. L., Hatting, H., and Deventer, C. S. van (2000). Genotype \times environment interaction of winter wheat (*Triticum aestivum* L.) In South Africa: II. Stability analysis of yield performance. *South African Journal of Plant and Soil* 17, 101–107. doi:[10.1080/02571862.2000.10634878](https://doi.org/10.1080/02571862.2000.10634878).
- Raju, B. M. K. (2002). A Study on AMMI Model and its Biplots. *Journal of the Indian Society of Agricultural Statistics* 55, 297–322.
- Sneller, C. H., Kilgore-Norquest, L., and Dombek, D. (1997). Repeatability of Yield Stability Statistics in Soybean. *Crop Science* 37, 383–390. doi:[10.2135/cropsci1997.0011183X003700020013x](https://doi.org/10.2135/cropsci1997.0011183X003700020013x).
- Wricke, G. (1962). On a method of understanding the biological diversity in field research. *Zeitschrift für Pflanzenzüchtung* 47, 92–146.
- Zali, H., Farshadfar, E., Sabaghpour, S. H., and Karimizadeh, R. (2012). Evaluation of genotype \times environment interaction in chickpea using measures of stability from AMMI model. *Annals of Biological Research* 3, 3126–3136. Available at: http://www.ijabbr.com/article_7777_620ea1a0c1fd04868f60bd23c6dda48b.pdf.
- Zhang, Z., Lu, C., and Xiang, Z. (1998). Analysis of variety stability based on AMMI model. *Acta Agronomica Sinica* 24, 304–309. Available at: <http://zwxb.chinacrops.org/EN/Y1998/V24/I03/304>.

Zobel, R. (1994). "Stress resistance and root systems," in *Proceedings of the Workshop on Adaptation of Plants to Soil Stress. 1-4 August, 1993. INTSORMIL Publication 94-2* (Institute of Agriculture; Natural Resources, University of Nebraska-Lincoln), 80-99.