

f. TRICHO NOURISH EVA

fluence

TOPICAL FINASTERIDE GEL WITH VITAMIN D3, AND ARACHIDONIC ACID

Finasteride

0.25%

Vitamin D3

5000 IU per gm

Arachidonic acid

0.01% w/w

A topical gel for Hirsutism



PRECAUTIONS:

- Watch for side effects like headache, itching or irritation of the scalp.
- In case of irritation, flush with water immediately and contact to your doctor.
- Avoid contact with eyes or broken skin. In such event, rinse thoroughly with plenty of water contact your doctor.
- Keep out of reach of children.
- F-Tricho Nourish Eva should not be used by pregnant women, lactating mothers & those planning a pregnancy.

F-TRICHONOURISH EVA GEL IS FOR EXTERNAL USE ONLY.

FOR HIRSUTISM

Finasteride IP0.25%w/w

In women with hirsutism, topically applied finasteride significantly decreases hair growth and thickness, with no adverse effects.

- Azziz R, Carmina E, Sawaya ME. Idiopathic hirsutism.
- Tahvilian R, Ebrahimi A, Beiki O, Nemati H, Masoud S. Preparation and clinical evaluation of Finasteride gel in the treatment of idiopathic Hirsutism. *J Drug Assess.*
- Farshi S, Mansouri P, Rafie F. A randomized double blind, vehicle controlled bilateral comparison study of the efficacy and safety of finasteride 0.5% solution in combination with intense pulsed light in the treatment of facial hirsutism. *J Cosmet Laser Ther*
- K. Jean Lucas, MD, FACP, FACE (2001) FINASTERIDE CREAM IN HIRSUTISM. *Endocrine Practice.*

APPLICATION ON SKIN:

1. Clean and dry the area before every application.
2. Take a small coin sized gel and spread gently in upward and circular motion on affected area of skin.
3. Apply once in a day, allow gel to remain on least for two hours.
4. Wash your hands after application.



STORAGE:

Store it in dry place below 30 degree temperature.
Protect from light and moisture.

Fluence Pharma Pvt. Ltd.

910, B Wing, Samartha Aishwarya, Off New Link Road, Opp. Highland Park Co-op Society, Andheri (W),
Mumbai 400 053. T: 022 2630 0106 / 0107, F: 022 2630 0108. Web: www.fluencepharma.com