

f-TRICHO NOURISH

FINASTERIDE GEL



Topical Finasteride Gel with D3, Arachidonic acid, Ketoconazole & Caffeine

Finasteride	0.25%
Vitamin D3	5000 IU per gm
Caffeine	0.01%
Arachidonic acid	0.01%
Ketoconazole	2%

**Topical benefits
without systemic
side effects**



PRECAUTIONS:

- Apply F-Tricho Nourish gel on the roots of hair.
- Watch for side effects, like headache, itching or irritation of the scalp.
- In case of irritation, flush with water immediately and contact to your doctor.
- Avoid contact with eyes or broken skin. In such a event, rinse thoroughly with plenty of water & contact your doctor.
- Keep out of reach of children
- F-Tricho Nourish gel should not be used by pregnant women, lactating mothers & those planning a pregnancy.

STORAGE:

Store it in dry place below 30 degree temperature.
Protect from light and moisture.

F-TRICHONOURISH GEL IS FOR EXTERNAL USE ONLY.

Fluence Pharma Pvt. Ltd.


910, B Wing, Samartha Aishwarya, Off New Link Road, Opp. Highland Park Co-op Society, Andheri (W),
Mumbai 400 053. T: 022 2630 0106 / 0107, F: 022 2630 0108. Web: www.fluencepharma.com

FOR ACTIVE HAIR GROWTH

1. Finasteride promotes active anagen hair growth cycle.
 - Sawaya ME, Blume-Peytavi U, et.al. Effects of finasteride on apoptosis and regulation of the human hair cycle. *J Cutan Med Surg*.
2. Topical application of Vit D3 has direct growth induction effect on hair root, dermal papilla cells.
 - Aoi N, Inoue K, et. al. 1 α , 25-dihydroxyvitamin D3 modulates the hair-inductive capacity of dermal papilla cells: therapeutic potential for hair regeneration. *Stem Cells Transl Med*.
3. Topical application of vit D3 helps hair regrowth in Alopecia Areata.
 - Kim DH, Lee JW, Kim IS, et al. Successful treatment of alopecia areata with topical calcipotriol. *Ann Dermatol*.
4. Topical application of Arachidonic acid stimulates cell proliferation in the hair matrix, improves circulation & action of growth factors.
 - Munkhbayar S, Jang S, et.al. Role of Arachidonic Acid in Promoting Hair Growth. *Ann Dermatol*. Promotes mitosis, matrix keratinocytes, angiogenesis, hair elongation & expression of Growth Factors in dermal papilla.
5. Topical Caffeine stimulates anagen growth phase, improves follicular penetration & prevents transepidermal loss.
 - Bansal M, Manchanda K, Pandey SS. Role of caffeine in the management of androgenetic alopecia. *Int J Trichology*.
6. Ketoconazole 2% regulates sebum secretion, dandruff and stimulates hair growth.
 - Inui S, Itami S. Reversal of androgenetic alopecia by topical ketoconazole: Relevance of anti-androgenic activity. *Journal of Dermatological Science*.

Application on scalp:

1. Application is recommended on a clean dry scalp.
2. Take a small coin sized gel & spread gently in circular motion over the affected area.
3. Apply once in a day, allow gel to remain for at least two hours or leave it overnight.
4. After two hours one may wash the gel and can apply any styling gel or oil as required.
5. If the application interferes with looks of the front hairline or styling, apply F-Tricho Nourish gel 1-2 inches behind the hair line, over top of the head and crown area. It will still reach all the hair through circulation & deliver benefit all over the scalp.
6. Wash your hands after application.



FOR REDUCING HIRSUTISM (EXCESS BODY HAIR)

Topical application of finasteride decreases facial hirsutism & hair thickness significantly, with no adverse effects..

- Azziz R, Carmina E, Sawaya ME. Idiopathic hirsutism.
- Tahvilian R, Ebrahimi A, Beiki O, Nemati H, Masoud S. Preparation and clinical evaluation of Finasteride gel in the treatment of idiopathic Hirsutism. *J Drug Assess*.

Application on skin:

1. Clean and dry the area before every application
2. Take a small coin sized gel and spread gently in upward motion on affected area of skin.
3. Apply once in a day, allow gel to remain on least for two hours.
4. Wash your hands after application.

