

# JEE EXPERT

## JEE MAIN EXCLUSIVE TOPICS

(SPECIAL TEST - 01)

PHYSICS, CHEMISTRY & MATHEMATICS

Batch : 12th (X01 & X02) [19.12.2019]

### Answer Key

#### PHYSICS

1	(A)	2	(C)	3	(A)	4	(C)	5	(A)
6	(A)	7	(D)	8	(C)	9	(D)	10	(B)
11	(C)	12	(A)	13	(A)	14	(C)	15	(C)
16	(C)	17	(C)	18	(B)	19	(A)	20	(B)
21	(C)	22	(B)	23	(B)	24	(A)	25	(A)
26	(C)	27	(A)	28	(A)	29	(D)	30	(A)

#### CHEMISTRY

31	(D)	32	(C)	33	(C)	34	(A)	35	(B)
36	(C)	37	(D)	38	(B)	39	(C)	40	(D)
41	(A)	42	(C)	43	(D)	44	(C)	45	(C)
46	(B)	47	(A)	48	(C)	49	(A)	50	(A)
51	(D)	52	(B)	53	(C)	54	(A)	55	(D)
56	(A)	57	(A)	58	(C)	59	(A)	60	(A)

#### MATHEMATICS

61	(C)	62	(A)	63	(D)	64	(C)	65	(D)
66	(D)	67	(C)	68	(C)	69	(C)	70	(C)
71	(B)	72	(B)	73	(A)	74	(B)	75	(C)
76	(A)	77	(B)	78	(A)	79	(B)	80	(D)
81	(D)	82	(C)	83	(B)	84	(C)	85	(B)
86	(C)	87	(B)	88	(A)	89	(B)	90	(A)

# JEE EXPERT

## JEE MAIN EXCLUSIVE TOPICS

(SPECIAL TEST - 01)

PHYSICS, CHEMISTRY & MATHEMATICS

Batch : 12th (X01 & X02) [19.12.2019]

### Solutions

#### PHYSICS

1. A lift is tied .....

**Sol.** The tension  $T$  in the rope of the lift when it goes upward is given by

$$T = (m)(g + a) = 1000(9.8 + 1.2) = 11000 \text{ N}$$

Let  $r$  be the radius of the wire, then the maximum stress will be  $\frac{T}{\pi r^2}$

$$\text{Hence, } \frac{T}{\pi r^2} = 1.4 \times 10^8$$

or

$$r^2 = \frac{T}{\pi \times 1.4 \times 10^8} = \frac{11000}{3.14 \times (1.4 \times 10^8)} = \frac{1}{4 \times 10^4}$$

$$\text{Now } r = \frac{1}{2 \times 10^2} = \frac{1}{200} = 0.005 \text{ m}$$

Therefore, diameter of the wire =  $2r = 0.01 \text{ m}$ .

2. The current .....

**Sol.** Given that,

$$V_1 = 0.01 \text{ volt}$$

$$R_1 = 1 \text{ k}\Omega = 10^3 \Omega$$

$$\therefore I_b = \frac{V_1}{R_1} = \frac{0.01}{1 \times 10^3} = 0.01 \times 10^{-3} \Omega = 0.01 \text{ mA}$$

$$\text{Further, } I_c = \beta I_b = 50 \times 0.01 \text{ mA}$$

$$= 0.5 \text{ mA} = 500 \mu\text{A}$$

3. What is the .....

**Sol.**  $m \frac{g}{2} = mg - F_B - F_V$

$$F_V + F_B = \frac{1}{2} mg$$

$$6\pi\eta r v + \sigma \frac{4}{3} \pi r^3 g = \frac{1}{2} \rho \frac{4}{3} \pi r^3 g$$

$$v = \frac{r^2 g}{9\eta} (\rho - 2\sigma)$$

4. An air bubble .....

**Sol.** Excess pressure inside an air bubble in soap solution

$$\frac{2T}{r} = \frac{2 \times 2.50 \times 10^{-2}}{5.00 \times 10^{-3}} = 10 \text{ pa}$$

Total pressure inside the air bubble

$$= P_0 + h\rho g + \frac{2T}{r}$$

$$= 1.01 \times 10^5 + 0.4 \times 1.2 \times 10^3 \times 9.8 + 10$$

$$= 1.06 \times 10^5 \text{ Pa}$$

5. The excess .....

**Sol.**  $\frac{P_1}{P_2} = 4$  or  $P_1 = 4P_2$

$$\frac{4s}{r^1} = \frac{4 \times 4s}{r_2} \Rightarrow \frac{r_1}{r_2} = \frac{1}{4}$$

$$\frac{V_1}{V_2} = \frac{r_1^3}{r_2^3} = \left(\frac{1}{4}\right)^3 = \frac{1}{64}$$

6. A mercury drop .....

**Sol.** Energy expended will be the work done against the increase in surface area i.e.

$$\Delta S = n(4\pi r^2) - 4\pi R^2$$

$$\Delta E = W = T\Delta S = T.4\pi(nr^2 - R^2)$$

But the total volume remains constant.

$$\text{i.e. } \frac{4}{3} \pi R^3 = n \frac{4}{3} \pi r^3$$

$$\text{or } r = \frac{R}{(n)^{1/3}} \text{ and hence, } E = 4\pi R^2 T (n^{1/3} - 1)$$

7. Water rises in .....

**Sol.**  $L = \frac{h'}{\cos \phi} = \frac{\left( \frac{2T \cos \theta}{\rho g r'} \right)}{\cos \phi} = \frac{3h}{\cos \phi}$

8. (C)

9. For a faulty .....

**Sol.** Zero error = MSR + (n)(L.C)  
 $= -0.1 + (7)(0.01) = -0.03 \text{ cm}$   
 Actual value = Measured value - Zero error  
 $= 8.65 - (-0.03) = 8.68 \text{ cm}$

10. (B)

11. If the length .....

**Sol.**  $B - A = (b \pm \Delta b) - (a \pm \Delta a) = (b - a) \pm (\Delta b + \Delta a)$   
 $= (4.19 - 3.25) \pm (0.01 + 0.01) = 0.94 \pm 0.02 \text{ cm}$

12. The vernier .....

**Sol.** L.C = 1 MSD - 1 VSD

13. A block of .....

**Sol.**  $A = \frac{F_0}{\sqrt{m^2 (\omega^2 - \omega_d^2)^2 + \omega_d^2 b^2}}$

In this case  $\omega_d = \omega = \sqrt{\frac{k}{m}} = 20 \text{ rad/s}$

so,  $A = \frac{F_0}{\omega_d b} \Rightarrow b = \frac{F_0}{\omega_d A} = \frac{0.1}{20 \times 5} = 10^{-3}$

14. In a Searle's .....

**Sol.** Young's modulus of elasticity is given by

$$Y = \frac{\text{stress}}{\text{strain}} = \frac{F/A}{\ell/L} = \frac{FL}{\ell A} = \frac{FL}{\ell \left( \frac{\pi d^2}{4} \right)}$$

Substituting the values, we get

$$y = \frac{50 \times 11 \times 4}{(1.25 \times 10^{-3}) \times \pi (5.0 \times 10^{-4})^2} = 2.24 \times 10^{11}$$

N/m<sup>2</sup>

Now,  $\frac{\Delta Y}{Y} = \frac{\Delta L}{L} + \frac{\Delta \ell}{\ell} + 2 \frac{\Delta d}{d}$

$$= \left( \frac{0.1}{110} \right) + \left( \frac{0.001}{0.125} \right) + 2 \left( \frac{0.001}{0.05} \right) = 0.0489$$

$$\Delta Y = (0.0489)Y = (0.0489) \times (2.24 \times 10^{11}) \text{ N/m}^2$$

$$= 1.09 \times 10^{10} \text{ N/m}^2$$

15. A carnot's .....

**Sol.** Efficiency =  $\eta = 1 - \frac{T_L}{T_H} = 1 - \frac{300}{900} = \frac{2}{3}$

$\therefore$  If heat rejected is E

$\Rightarrow$  heat supplied is 3E and work done is 2E

$$\text{So, } 2E = \frac{1}{2} \left( \frac{mR^2}{2} \right) \omega^2 \Rightarrow \omega = \sqrt{\frac{8E}{mR^2}}$$

16. A monochromatic .....

**Sol.**  $\frac{1}{2} = \frac{1}{2} \epsilon_0 E_{\text{rms}}^2 \Rightarrow 1 = \frac{1}{2} \epsilon_0 E_0^2$

$$\frac{P}{4\pi r^2} = \frac{1}{2} \epsilon_0 E_0^2 C \Rightarrow E_0 = \frac{1}{r} \sqrt{\frac{P}{2\pi \epsilon_0 C}}$$

$$\text{and } C = \frac{1}{\sqrt{\mu_0 \epsilon_0}} \Rightarrow \frac{1}{\epsilon_0} = \mu_0 C^2$$

$$\text{so, } E_0 = \frac{1}{r} \sqrt{\frac{P}{2\pi C} \times \mu_0 C^2} = \frac{1}{r} \sqrt{\frac{\mu_0 P C}{2\pi}}$$

16. A simple microscope .....

**Sol.** R.P. =  $\frac{2\mu \sin 30^\circ}{\lambda}$

17. A paramagnetic .....

**Sol.** If the temperature is decreased, the thermal vibration will be reduced. So there would not be negative effect on magnetisation.

19. A Ge specimen .....

**Sol.** When Ge specimen is doped with A  $\ell$ , then concentration of acceptor atoms is also called concentration of holes.

Using formula  $n_i^2 = n_e n_h$

$n_i$  = concentration of electron hole pair =  $10^{19}/\text{m}^3$

$n_e$  = concentration of electrons

$n_h$  = concentration of holes =  $10^{21} \text{ atoms/m}^3$

$$\therefore (10^{19})^2 = 10^{21} \times n_e, n_e = 10^{17} / \text{m}^3$$

20. The following .....

**Sol.** Out of upper ABD gate =  $A\bar{B}$

Output of lower AND gate =  $B\bar{A}$

Output  $Y = \overline{AB} + \overline{BA}$

This is Boolean expression for XOR gate.

21. In a given .....

Sol. Given at saturation

$$V_{CE} = 0V, V_{BE} = 0.8V$$

$$V_{CE} = V_{CC} - I_C R_C$$

$$I_C = V_{CC}/R_C = 5.0V/1.0k\Omega = 5.0mA$$

$$\text{Therefore } I_B = I_C/\beta = 5.0mA/250 = 20\mu A$$

The input voltage at which the transistor will go into saturation is given by

$$V_{IH} = V_{BB} = I_B R_B + V_{BE} = 20\mu A \times 100k\Omega + 0.8V = 2.8V$$

22. The current gain .....

$$\text{Sol. } \alpha = \frac{I_C}{I_E} = \frac{I_E - I_B}{I_E} \Rightarrow 0.95 = \frac{10 - I_B}{10}$$

$$\Rightarrow I_B = 0.5mA$$

23. In an npn .....

Sol. Collector current,

$$I_C = 10mA; \alpha = \frac{90}{100} = 0.9$$

$$\alpha = \frac{I_C}{I_E} \Rightarrow \text{Emitter current, } I_E = \frac{I_C}{\alpha}$$

$$I_E = \frac{10}{0.9} = 11mA$$

$$\text{Base current, } I_B = I_E - I_C$$

$$I_E = \frac{10}{0.9} = 11mA$$

$$\text{Base current, } I_B = I_E - I_C$$

$$I_B = 11 - 10 = 1mA$$

24. Consider an .....

Sol. Optical source frequency

$$f = \frac{c}{\lambda} = 3 \times 10^8 / (800 \times 10^{-9})$$

$$= 3.8 \times 10^{14} \text{ Hz}$$

$$\text{Bandwidth of channel (1\% of above)} = 3.8 \times 10^{12} \text{ Hz}$$

$$\text{Number of channel} = (\text{Total bandwidth of channel}) / (\text{Bandwidth needed per channel})$$

$$\text{Number of channels for audio signal}$$

$$= (3.8 \times 10^{12}) / (8 \times 10^3) = 4.8 \times 10^8$$

25. The diameter .....

$$\text{Sol. } d\theta = \frac{1.22\lambda}{a}$$

27. Linearly polarised .....

$$\text{Sol. In incident light, } \frac{E_z}{E_y} = \sqrt{3} \Rightarrow \vec{E}$$

is at angle  $60^\circ$  from y-axis

whereas transmission axis ( $y = z$ ) is at  $45^\circ$  from y-axis

$$\text{So, } E_{\text{transmitted}} = E_{\text{incident}} \cos(60^\circ - 45^\circ) \\ = 2E_0 \cos 15^\circ$$

28. In Amplitude .....

$$\text{Sol. Modulation index} = \frac{A_m}{A_c}$$

29. Following diagram .....

Sol. We know that combination of AND gate and NOT gate is known as NAND gate since the given symbol contains AND gate and NOT, therefore it is a NAND gate.

30. If  $\rho$  is the .....

$$\text{Sol. Stress} = \text{weight/area} = A\ell\rho g / A = \ell\rho g \\ \Rightarrow \sigma = \ell\rho g$$

## CHEMISTRY

31. (1)  $\Delta G = \Delta H - T\Delta S < 0$  as  $\Delta S < 0$  so  $\Delta H$  has to be negative.

(2) micelles formation will take place above  $T_k$  and above CMC.

(4)  $\text{Fe}^{3+}$  ions will have greater flocculability power so smaller flocculating value.

38. The phenomenon of scattering of light from the surface of colloidal particles is called tyndall effect. The minimum conc. required for micelle formation is called critical micelle concentration.

For soap solution above CMC tyndall effect is observed.

39. Higher the gold number, lesser will be the protective power of colloid.

40. Colloidal solution shows brownian movement. The Brownian movement does not allow the collidal particles to settle down to gravity & thus responsible for their stability.

41. Easily liquefiable gases like  $\text{CO}_2$  are adsorbed to a greater extent than gases like  $\text{O}_2$ ,  $\text{N}_2$  and  $\text{H}_2$

42. Mass adsorbed =  $\frac{100(0.3-0.125)60}{1000}$   
 $= (0.3-0.125)6$   
 Mass adsorbed per gram of charcoal  
 $= \frac{(0.3-0.125)6}{0.8} = 1.31\text{g}$

## MATHEMATICS

61. If 'R' is an equivalence .....

$$\forall a \in A, (a, a) \in R \Rightarrow (a, a) \in R^{-1}$$

$$[(a, b) \in R \Rightarrow (b, a) \in R]$$

$$\Rightarrow \{(a, b) \in R^{-1}\} \wedge \{(b, a) \in R^{-1}\}$$

$$[(a, b) \in R \wedge (b, c) \in R] \rightarrow (a, c) \in R$$

$$[(c, b) \in R^{-1} \wedge (b, a) \in R^{-1}] \rightarrow (c, a) \in R^{-1}$$

$\therefore$  It is an equivalence relation

62. Find cardinal number .....

Sol. Let A be the set of positive integers not exceeding 1000 that are divisible by 7 and B  $\rightarrow$  divisible by 11

$$n(A \cup B) = n(A) + n(B) - n(A \cap B)$$

$$= \left[ \frac{1000}{7} \right] + \left[ \frac{1000}{11} \right] - \left[ \frac{1000}{77} \right] = 142 + 90 - 12 = 220$$

63. Let  $P = \left\{ \theta : \sin^2 \theta + \cos^4 \theta = \frac{1 - \sin \theta}{\cos \theta} \right\}$  .....

Sol.  $\sin^2 \theta + \cos^4 \theta = \cos^2 \theta + \sin^4 \theta$  and

$$\frac{1 - \sin \theta}{\cos \theta} = \cot \left( \frac{\pi}{4} + \frac{\theta}{2} \right)$$

64. In a survey of .....

Sol.  $n(M' \cap C' \cap P') = n(\cup) - n(M \cap P \cap C)$

$$\Rightarrow n(M \cup P \cup C) = 450$$

$$n(M \cap P \cap C) = 20$$

$$\text{who liked exactly one} = 95 + 40 + 190 = 325$$

65. Which of the following .....

Sol. Obvious

66. The relation R .....

Sol. Obvious


67. Number of elements .....

Sol.  $P = \frac{2n^5 + 3n^4 + 4n^3 + 4n^2 + 6}{n}$  is integer

$$\Rightarrow n = 1, 2, 3, 6$$

68. If A, B, C are 3 .....

Sol.



69. Which of the following .....

Sol. Obvious

70. The relation T .....

Sol.  $a \geq a^{\frac{2}{3}} \Rightarrow a^3 \geq a^2$  not true for  $\bar{\mathbb{Z}}$  so not reflexive

$$27 \geq 8^{\frac{2}{3}} \text{ but } 8 \not\geq 27^{\frac{2}{3}} \text{ so not symmetric}$$

$$a \geq b^{\frac{2}{3}} \text{ and } b^3 \geq C^{\frac{2}{3}} \quad (\because a \geq 0, b \geq 0)$$

$$\Rightarrow a \geq (C^{\frac{2}{3}})^2 \Rightarrow a^3 \geq C^{\frac{2}{3}}$$

$$\left( \because (C^{\frac{2}{3}})^2 \geq C^{\frac{2}{3}} \right)$$

$\Rightarrow$  transitive

71. Consider relation .....

Sol. (a, b) is not related to (a, b) as  $\text{LCM}(a, a) \neq \text{LCM}(b, b)$

72. The number of .....

Sol.  $R_1 = \{(a, a), (b, b), (c, c)\}$

$R_2 = (a, b), (b, a), (a, a), (b, b), (c, c)$   
 $R_3 = (a, c), (c, a), (a, a), (c, c), (b, b)$   
 $R_4 = (b, c), (c, b), (b, b), (c, c), (a, a)$   
 $R_5 = \{(a, a), (b, b), (c, c), (a, b), (b, a), (a, c), (c, a), (b, c), (c, b)\}$

**73.** Let R and S be .....

**Sol.** Let  $A = \{1, 2, 3\}$   
 $R = \{(1, 1), (1, 2)\}$   
 $S = \{(2, 2), (2, 3)\}$  be transitive relations on A.  
Then  $R \cup S = \{(1, 1), (1, 2), (2, 2), (2, 3)\}$   
Which shows that  $R \cup S$  is not transitive  
Since  $(1, 2) \in R \cup S$  and  $(2, 3) \in R \cup S$  but  $(1, 3) \notin R \cup S$

**74.** Consider relation .....

**Sol.** Obvious

**75.** Let  $A = \{2, 3, 4, \dots, 17, 18\}$  .....

**Sol.** (21 to 22)

$$(3, 2) \sim (c, d) \Rightarrow 3d = 2c \Rightarrow c = \frac{3d}{2}$$

$$\Rightarrow d = 2, 4, 6, 8, 10, 12$$

**76.** Let A and B be .....

**Sol.**  ${}^6C_3 + {}^6C_4 + {}^6C_5 + {}^6C_6 = 2^6 - {}^6C_0 - {}^6C_1 - {}^6C_2$   
 $= 64 - 1 - 6 - 15 = 64 - 22 = 42$

**77.** Let  $A = \{z : |z|^2\}$  .....

**Sol.** The given equation are the equations of two circles touching, each other externally.

**78.** Let A and B be .....

**Sol.** B:  $\frac{(x-4)^2}{9} + \frac{(y-3)^2}{4} \leq 1$   
A:  $2 < x < 4$  and  $2 < y < 4$

**79.** Consider the following .....

**Sol.** The given relation is only symmetric

**80.** Let  $A = \{(x, y) : \dots\}$

**Sol.** APA  $\Rightarrow$  reflexive  
area  $(A - B) = 0$  and area  $(B - A) = 4\pi - \pi = 3\pi$   
 $\Rightarrow (A, B) \notin P \Rightarrow$  symmetric and transitive

**81.** Consider relations .....


**Sol.**  $R_1$  is reflexive, not symmetric and transitive  
 $R_2$  is not reflexive, symmetric and not transitive

**82.** The intersection of .....

**Sol.** Obvious

**83.** Let A and B be .....

**Sol.**  $(A \cap B = \phi) \Rightarrow A \subset (X - B)$



**84.** Let A consist of .....

**Sol.** The subsets of A which contain at least one of  $a_1, a_2, \dots, a_r$  and all of  $a_{s+1}, a_{s+2}, \dots, a_n$  are obtained, once each, as the unions of  $\{a_{s+1}, a_{s+2}, \dots, a_n\}$  with the subsets of  $\{a_1, a_2, \dots, a_s\}$  which contains at least one of  $a_1, a_2, a_3, \dots, a_r$  from the subsets of  $\{a_1, a_2, \dots, a_s\}$  which contains at least  
 $= 2^{s-r}(2^r - 1)$

**85.** Relation  $\otimes$  is defined .....

**Sol.** Obvious

**86.** Relation C is defined .....

**Sol.** Obvious

**87.** The number of .....

**Sol.** Total no. of relations  $= 2^{4 \times 4} = 2^{16}$   
In cartesian product we have 16 ordered pairs in which 4 are compulsory of reflexive relation, out of 12 ordered pairs.  
We can take  
 ${}^{12}C_0 + {}^{12}C_1 + {}^{12}C_2 + \dots + {}^{12}C_{12} = 2^{12}$

**88.** Consider the two .....

**Sol.** Obvious

**89.** Consider relation .....

**Sol.**  $(a, b) \in T$  and  $(b, a) \in T$  but  $(a, a) \notin T$  so not transitive

**90.** If an examination .....

**Sol.** Only failed in physics  $= 37\% - 19\% = 18\%$   
 $=$  alone pass in maths. let total  $= x$   
 $x = 1700$