

Active sites of nitrogen-doped carbon materials for oxygen reduction reaction clarified using model catalysts

Donghui Guo, Riku Shibuya, Chisato Akiba, Shunsuke Saji, Takahiro Kondo and Junji Nakamura

Science **351** (6271), 361-365.
DOI: 10.1126/science.aad0832

The right kind of dopant

The oxygen reduction reaction is an important step in fuel cells and other electrochemical processes but is still largely dependent on precious metal-containing catalysts. Recently explored alternatives include carbon materials that are doped with different, preferably non-precious metal, atoms. Guo *et al.* studied model graphite catalysts to try to understand the role of nitrogen doping and to elucidate the active catalytic sites. A nitrogen atom bound to two carbons formed an active catalyst site with an activity rivaling that of N-doped graphene catalysts.

Science, this issue p. 361

ARTICLE TOOLS

<http://science.sciencemag.org/content/351/6271/361>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2016/01/20/351.6271.361.DC1>

REFERENCES

This article cites 21 articles, 2 of which you can access for free
<http://science.sciencemag.org/content/351/6271/361#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science* is a registered trademark of AAAS.