

ORIGINAL ARTICLE

Anti-tumour necrosis factor agents reduce non-steroidal anti-inflammatory drug-induced small bowel injury in rheumatoid arthritis patients

Toshio Watanabe,¹ Tetsuya Tanigawa,¹ Masatsugu Shiba,¹ Yuji Nadatani,¹ Yasuaki Nagami,¹ Satoshi Sugimori,¹ Hirokazu Yamagami,¹ Kenji Watanabe,¹ Kazunari Tominaga,¹ Yasuhiro Fujiwara,¹ Tatsuya Koike,² Tetsuo Arakawa¹

¹Department of Gastroenterology, Osaka City University Graduate School of Medicine, Osaka, Japan
²Department of Rheumatology, Osaka City University Graduate School of Medicine, Osaka, Japan

Correspondence to

Dr Toshio Watanabe,
Department of Gastroenterology, Osaka City University Graduate School of Medicine, 1-4-3 Asahimachi, Abeno-ku, Osaka 545-8585, Japan;
watanabet@med.osaka-cu.ac.jp

Received 13 February 2013

Revised 23 April 2013

Accepted 23 April 2013

ABSTRACT

Objective The role of tumour necrosis factor α (TNF α) in the pathogenesis of non-steroidal anti-inflammatory drug (NSAID)-induced small intestinal damage remains unclear. We evaluated the preventive effect of anti-TNF therapy against NSAID-induced enteropathy in rheumatoid arthritis (RA) patients.

Design Capsule endoscopy was performed in 95 consecutive RA patients who received NSAID for more than 3 months, with or without anti-TNF therapy over a period of 3 months. The findings were scored from 0 to 4: 0, normal; 1, red spots; 2, one to four erosions; 3, more than four erosions; and 4, large erosions/ulcers. The relationship between the use of anti-TNF therapy and the risk of severe damage (scores 3 or 4) or the most severe damage (score 4) was assessed using multiple logistic regression analysis. Furthermore, a propensity score matching analysis was performed to reduce the effects of TNF selection bias.

Results By stratifying the patients on the basis of anti-TNF therapy, we obtained crude OR of 0.23 for severe damage (95% CI 0.09 to 0.65) and 0.37 for the most severe damage (95% CI 0.16 to 0.86). This protective effect of anti-TNF therapy remained robust to adjustments for baseline characteristics, with the adjusted OR for severe damage and the most severe damage ranging from 0.23 to 0.26 and 0.06 to 0.41, respectively. Propensity score matching yielded similar results and showed the protective effects of anti-TNF therapy against severe and most severe damage.

Conclusions Anti-TNF therapy may protect against NSAID-induced small intestinal damage in RA patients.

INTRODUCTION

Although disease-modifying antirheumatic drugs (DMARD) such as methotrexate have gained increasing importance in the treatment of rheumatoid arthritis (RA), patients with RA still frequently take non-steroidal anti-inflammatory drugs (NSAID) to reduce joint pain, stiffness and swelling. The most common and serious adverse effect of NSAID therapy is gastrointestinal toxicity that can cause ulceration, bleeding and perforation. Until quite recently, most studies had focused on NSAID-induced upper gastrointestinal complications. However, since the revolutionary advance in the detection of small bowel diseases that was brought about by the development of video capsule

Significance of this study

What is already known on this subject?

- ▶ NSAID frequently injure the small intestine as well as the upper gastrointestinal tract.
- ▶ Experimental studies have demonstrated that TNF α plays an important role in the development of NSAID-induced damage to the small intestine.
- ▶ Anti-TNF biological agents are effective in the treatment of gastrointestinal lesions caused by several inflammatory disorders in which TNF α plays an important pathogenetic role.

What are the new findings?

- ▶ In the anti-TNF therapy-naïve group, about half of the patients had NSAID-induced severe small intestinal damage.
- ▶ RA patients who received anti-TNF therapy had less severe NSAID-induced small bowel damage than those who did not receive such therapy.
- ▶ The protective effect of anti-TNF therapy against severe enteropathy remained robust to propensity score adjustment.

How might it impact on clinical practice in the foreseeable future?

- ▶ We demonstrate a pathogenic role of TNF α in NSAID-induced enteropathy in humans.
- ▶ Therapies targeting TNF α , and particularly anti-TNF biological therapy, could be a novel treatment for NSAID-induced severe small intestinal pathologies.

endoscopy (VCE)¹ in the beginning of the 21st century, NSAID-induced small bowel injury has received close attention. Several studies using VCE have demonstrated the high potential of NSAID to injure the small intestine, regardless of the length of NSAID therapy. It has been demonstrated that 40–90% of NSAID users have mucosal breaks (ulcers and/or erosions) in the small intestine.^{2–5} A high prevalence of NSAID-induced enteropathy was also observed in our previous study, in which 81% of RA patients who received NSAID for more than 3 months had mucosal breaks in the small intestine.⁶

To cite: Watanabe T, Tanigawa T, Shiba M, *et al.* Gut Published Online First: [please include Day Month Year] doi:10.1136/gutjnl-2013-304713

Small bowel

Tumour necrosis factor α (TNF α) plays a key role in the pathogenesis of many inflammatory diseases, including RA, inflammatory bowel diseases and psoriasis. Therefore, anti-TNF agents have been used to treat these inflammatory pathologies.⁷ In RA, anti-TNF biological therapies, which are used when conventional DMARD are unable to control RA synovitis, have been shown to be highly effective in reducing disease activity, disability and radiological damage.^{8–9} Recent experimental studies have indicated that TNF α plays an important role in the development of NSAID-induced damage in the small intestine, similar to its role in the stomach.^{10–11} In rats, the in-vivo blocking of TNF α by neutralising antibodies led to a significant reduction in indomethacin-induced small intestinal damage,¹² and TNF α knockout mice exhibited less severe damage, with reduction in neutrophil infiltration and epithelial cell apoptosis.¹³ Therefore, we hypothesised that RA patients who chronically used NSAID and also received anti-TNF therapy would exhibit less severe enteropathy than those not receiving the therapy.

METHODS

Patients

Between January 2008 and December 2011, 103 consecutive RA patients (19 men and 84 women; mean age 61 years; age range 24–80 years) who received NSAID for more than 3 months, with or without anti-TNF therapy over a period of 3 months, were enrolled in this study. The main exclusion criteria included swallowing disorders, active malignancy, pregnancy; history of abdominal surgery and any serious central nervous system, psychiatric, cardiovascular, respiratory, or intestinal disease. Patients on misoprostol were excluded from the study because of the preventive effects of this drug on NSAID-induced damage to the small intestine that were confirmed by a VCE study.⁵ As we specifically focused on the effects of long-term anti-TNF biological administration on NSAID-induced enteropathy in the present study, we excluded patients who received anti-TNF therapy for less than 3 months or patients who received biological agents targeting molecules other than TNF α , such as tocilizumab and abatacept. Written informed consent was obtained from all the patients. The protocol for the study was approved by the ethics committee of the Osaka City University Graduate School of Medicine.

VCE procedure and data interpretation

VCE was performed using a PillCam SB1 or SB2 (Given Imaging, Ltd, Yoqneam, Israel). Patients fasted for 12 h before swallowing the capsule. Fluids were allowed 2 h later, followed by a light meal after another 2 h. Data were collected for up to 8 h after capsule ingestion. After 8 h, the sensor array and recording device were removed.

The VCE digital image stream was reviewed and interpreted independently by two endoscopists (YN and SS) who were blind to the treatments of the patients. Capsule endoscopic findings of NSAID-induced enteropathy were scored according to the following method of Graham *et al.*:¹⁴ 0, normal; 1, red spots; 2, one to four small erosions; 3, more than four small erosions; and 4, large erosions/ulcers. A red spot was defined as a mucosal disruption that was denuded of villi, with red areas but no clear mucosal breaks. A small erosion was defined as a circumscribed area of mucosal disruption that was denuded of villi, with or without exudates or red colour and that involved, at most, a diameter that was equivalent to those of valvulae conniventes. Large erosions were defined as circumscribed breaks in the mucosa that were larger than the equivalent diameter of valvulae conniventes. Ulcers were defined as large erosions with a

central area with exudates. Typical examples of small bowel pathologies are shown in figure 1. Endoscopic scores of 3 and 4 indicated severe damage.¹⁴ If the endoscopic findings were different between the two interpreters, the findings were reviewed together by both blinded interpreters to reach a consensus.

Statistical analysis

Data are presented as mean \pm SD for continuous variables and as numbers (percentages) for categorical variables. For categorical data, comparisons between groups were performed using the χ^2 test (or Fisher's exact test when necessary because of small sample sizes), whereas continuous data were compared using Student's t test. The relationship between the use of anti-TNF therapy and the risk of severe damage (endoscopic scores 3 or 4) or the most severe damage (endoscopic score 4) was estimated by calculating OR and 95% CI by using multiple logistic regression.

Furthermore, a propensity score matching analysis was performed to reduce the effects of TNF selection bias and potential confounding factors. The propensity score matching method was proposed to evaluate statistically causal effects free from confounding effects, by mathematically refashioning an observational study into a randomised study; therefore, studies using the propensity score approach are occasionally considered pseudo-randomised studies.¹⁵ We created a propensity score matched cohort by attempting to match each RA patient receiving anti-TNF therapy with a patient not receiving anti-TNF therapy (a 1:1 match) without replacement by using an optimal matching technique. A total of 11 variables that could possibly influence the outcomes (severe or most severe damage) was used to generate a propensity score ranging from 0 to 1 by logistic regression.¹⁶ These variables included age, gender, history of peptic ulcers, drinking and smoking habits, dose of NSAID, type of NSAID (non-selective cyclooxygenase (COX) inhibitor or selective COX-2 inhibitor (celecoxib)), concomitant use of salazosulapyridine, bisphosphonate, and acid suppressants (proton pump inhibitors (PPI) or histamine-2-receptor




Figure 1 Capsule endoscopic images of small bowel pathologies induced by non-steroidal anti-inflammatory drugs. (A) red spot (arrow), (B) small erosion, (C) large erosion, (D) ulcer.

antagonists (H₂RA)) and disease activity score in 28 joints (DAS28, a disease activity index for RA). After matching, crude comparisons of the matched cohorts were performed using the McNemar's test and paired t tests; OR with 95% CI were calculated using conditional logistic regression. The propensity score matched sets were then used in logistic regression models to evaluate the relationship between the use of anti-TNF therapy and the risk of severe damage or the most severe damage. Absolute standardised differences (ASD) were computed to evaluate matching effectiveness. Statistical analyses were performed using SPSS version software 11.0 for Windows and the R statistical package (free download from <http://www.r-project.org>) V2.13.0. All statistical tests were two-sided, and p values less than 0.05 were considered to be statistically significant.

RESULTS

Baseline characteristics of patients

Comparative data were analysed for 95 patients; eight patients (all women; mean age 58 years) were excluded because the entire small bowel could not be visualised. A total of 65 patients received the standard doses of NSAID, 30 patients were being treated with lower doses of NSAID, and no patients received higher doses of NSAID. Of the 95 patients, 59 patients did not receive anti-TNF therapy, while 36 patients received anti-TNF therapy (etanercept (n=19), infliximab (n=15) and adalimumab (n=2)).

Table 1 shows the baseline demographic information, as well as the clinical variables. Compared to the patients not receiving anti-TNF therapy, the patients receiving anti-TNF therapy had a higher rate of concomitant use of methotrexate, with a tendency to be younger and to use selective COX-2 inhibitors more frequently. The two treatment groups were similar with respect to gender, history of peptic ulcers, drinking and smoking habits, NSAID doses, rates of concomitant use of steroids, salazosulfapyridine, bisphosphonate and acid suppressants, and DAS28 score.

After propensity score matching, there were 36 matched pairs of patients who did or did not receive anti-TNF therapy. The baseline characteristics of the two groups were comparable. The C statistic, which is equivalent to the receiver operating

characteristic curve, was 0.72 for the propensity model. The ASD values ranged from 6% to 35%.

Effect of anti-TNF therapy on NSAID-induced severe small intestinal damage

Of the 59 patients in the anti-TNF therapy-naïve group, 27 (45.8%) had severe damage (score of 3 (n=9) or 4 (n=18)), while the remaining 32 patients had non-severe (score<3) damage (scores of 1 (n=20) and 2 (n=12)). Of the 36 patients in the anti-TNF therapy group, six (16.7%) had severe damage (score of 3 (n=5) or 4 (n=1)), while the remaining 30 patients had non-severe damage (scores of 0 (n=6), 1 (n=15) and 2 (n=9)).

By categorising the patients on the basis of anti-TNF therapy, we obtained OR of 0.23 for severe damage (95% CI 0.09 to 0.65) and 0.37 for the most severe damage (95% CI 0.16 to 0.86). This effect of anti-TNF therapy on NSAID-induced enteropathy remained robust to adjustments for age, drinking and smoking habits, use of salazosulfapyridine, bisphosphonate, or acid suppressants, and DAS28 score, with the adjusted OR for severe damage and the most severe damage ranging from 0.23 to 0.26 and 0.06 to 0.41, respectively (table 2).

Propensity score matching yielded similar results in terms of the protective effects of anti-TNF therapy, with the adjusted OR for severe damage and the most severe damage ranging from 0.23 to 0.33 and 0.09 to 0.11, respectively (table 2).

DISCUSSION

In this study, we demonstrated that RA patients who received anti-TNF therapy had less severe NSAID-induced small bowel damage than those who did not receive such therapy. Given that the observed association between the use of anti-TNF biological agents and the risk of severe or most severe NSAID-induced enteropathy could be affected by confounding factors such as gender, age, dose of NSAID and clinical activity of RA, we conducted a sensitivity analysis by using the propensity score matching method. The results of this analysis showed a similar reduction in the risk of NSAID-induced severe damage to the small intestine in RA patients who received anti-TNF therapy. These results strongly suggest that anti-TNF biological agents

Table 1 Baseline characteristics and propensity score matched baseline characteristics

	Baseline			Propensity score matched baseline			
	TNF- (n=59)	TNF+ (n=36)	p Value	TNF- (n=36)	TNF+ (n=36)	p Value	ASD (%)
Age (years)*							
Mean (SD)	63 (10)	59 (12)	0.09	60 (11)	59 (13)	0.56	9
Female, * n (%)	46 (78.0)	30 (83.3)	0.6	34 (94.4)	30 (83.3)	0.29	35
History of peptic ulcer, * n (%)	20 (3.9)	12 (33.3)	1.00	11 (30.6)	12 (33.3)	1.00	6
Alcohol intake, * n (%)	8 (13.6)	5 (13.9)	1.00	4 (11.1)	5 (13.9)	1.00	8
Smoker, * n (%)	8 (13.6)	6 (16.7)	0.77	5 (13.9)	6 (16.7)	1.00	8
Standard dose of NSAID, * n (%)	38 (64.4)	27 (75.0)	0.36	23 (63.9)	27 (75.0)	0.45	24
Selective COX-2 inhibitors, * n (%)	9 (15.3)	12 (33.3)	0.05	9 (25.0)	12 (33.3)	0.55	18
Methotrexate use, n (%)	42 (71.2)	33 (91.7)	0.02	31 (86.1)	33 (91.7)	0.68	–
Steroid use, n (%)	18 (30.5)	17 (47.2)	0.13	12 (33.3)	17 (47.2)	0.38	–
Salazosulfapyridine use, * n (%)	9 (15.3)	3 (8.3)	0.53	4 (11.1)	3 (8.3)	1.00	9
Bisphosphonate use, * n (%)	13 (22.0)	4 (11.1)	0.27	6 (16.7)	4 (11.1)	0.68	16
Acid suppressants use, * n (%)	30 (50.8)	19 (52.8)	1.00	18 (50.0)	19 (52.8)	1.00	6
DAS28*	3.3±1.1	3.1±0.9	0.48	3.2±1.0	3.1±0.9	0.71	11

*Variable used to compute propensity scores.

ASD, absolute standardised difference; COX, cyclooxygenase; DAS28, disease activity score in 28 joints; NSAID, non-steroidal anti-inflammatory drug; TNF, tumour necrosis factor.

Table 2 The relationship between the use of anti-TNF therapy and the risk of NSAID-induced severe damage or most severe damage

	Severe damage		Most severe damage	
	OR (95% CI)	p Value	OR (95% CI)	p Value
Before propensity score matching				
Crude	0.23 (0.09 to 0.65)	<0.01	0.37 (0.16 to 0.86)	0.02
Adjusted for age	0.26 (0.09 to 0.74)	0.01	0.41 (0.17 to 0.97)	0.04
Adjusted for alcohol intake and smoking	0.23 (0.08 to 0.66)	<0.01	0.07 (0.01 to 0.52)	0.01
Adjusted for salazosulfapyridine use	0.23 (0.08 to 0.65)	<0.01	0.06 (0.01 to 0.47)	<0.01
Adjusted for bisphosphonate use	0.25 (0.09 to 0.71)	<0.01	0.07 (0.01 to 0.55)	0.01
Adjusted for acid suppressant use	0.23 (0.08 to 0.64)	<0.01	0.06 (0.01 to 0.50)	<0.01
Adjusted for DAS28	0.24 (0.09 to 0.65)	<0.01	0.37 (0.16 to 0.88)	0.02
Adjusted for age, alcohol intake, smoking and use of salazosulfapyridine and acid suppressants	0.23 (0.08 to 0.67)	<0.01	0.06 (0.01 to 0.47)	<0.01
After propensity score matching				
Crude	0.29 (0.09 to 0.87)	0.03	0.09 (0.01 to 0.70)	0.02
Adjusted for age	0.26 (0.08 to 0.86)	0.03	0.07 (0.01 to 0.86)	0.04
Adjusted for alcohol intake and smoking	0.26 (0.07 to 0.87)	0.03	0.11 (0.01 to 0.88)	0.04
Adjusted for salazosulfapyridine use	0.31 (0.10 to 0.94)	0.04	0.10 (0.01 to 0.78)	0.03
Adjusted for bisphosphonate use	0.30 (0.10 to 0.93)	0.04	0.11 (0.01 to 0.88)	0.04
Adjusted for acid suppressant use	0.27 (0.08 to 0.86)	0.03	0.09 (0.01 to 0.72)	0.02
Adjusted for DAS28	0.33 (0.10 to 1.04)	0.06	0.11 (0.01 to 1.18)	0.07
Adjusted for age, alcohol intake, smoking and use of salazosulfapyridine and acid suppressants	0.23 (0.05 to 1.01)	0.05	0.09 (0.01 to 1.32)	0.08

Capsule endoscopic findings of NSAID-induced enteropathy were scored from 0 to 4: 0, normal; 1, red spots; 2, one to four small erosions; 3, more than four small erosions; 4, large erosions/ulcers. The relationship between the use of anti-TNF therapy and the risk of severe damage (scores 3 or 4) or severest damage (score 4) was evaluated by logistic regression analysis.

DAS28, disease activity score in 28 joints; NSAID, non-steroidal anti-inflammatory drug; TNF, tumour necrosis factor.

may have a protective effect against NSAID-induced enteropathy. As our study was a cross-sectional study, the possibility that the anti-TNF therapy accelerated the healing of such enteropathy should be considered. However, further studies are required to determine whether the anti-TNF biological agents exert these two effects (a protective effect and a wound-healing effect) in NSAID-induced enteropathy.

Recently, anti-TNF biological agents have been reported to be effective in the treatment of gastrointestinal lesions caused by several inflammatory disorders in which TNF α plays an important pathogenetic role. For example, a meta-analysis by Peyrin-Biroulet *et al*¹⁷ demonstrated that anti-TNF biological agents such as infliximab and adalimumab were effective in inducing clinical remission and fistula closure in patients with Crohn's disease. Another meta-analysis demonstrated that infliximab was superior to placebo in inducing the remission of moderate to severely active ulcerative colitis.¹⁸ Furthermore, Melikoglu *et al*¹⁹ reported a significant improvement in the oral lesions associated with Behçet's syndrome with etanercept therapy. Multiple lines of evidence from experimental studies suggest that the overexpression of TNF α occurs in the early phase of NSAID-induced small intestine damage and that TNF α may play a crucial role in the subsequent induction of inflammatory responses, such as neutrophil infiltration, resulting in intestinal injuries.^{12 13 20 21} These animal data have led to the hypothesis that anti-TNF therapy could also prevent NSAID-induced damage to the small intestine. As expected, we found that anti-TNF therapy had a protective effect against the damage. Moreover, we demonstrated for the first time a pathogenic role of TNF α in NSAID-induced enteropathy in humans.

In addition to TNF α , various factors such as enteric bacteria, deficiency of prostaglandins and bile have been shown to be involved in the pathogenesis of NSAID-induced small intestinal damage, and in experimental studies, many agents have shown the potential to prevent the damage.^{3 12 20 21} However, only a small

number of these agents have been demonstrated to be effective for NSAID-induced enteropathy in clinical studies. Misoprostol²² and rebamipide²³ (a muco-protective drug) exerted protective effects against diclofenac-induced small-intestinal mucosal injuries in humans. Importantly, these studies were conducted in young, healthy volunteers with low risks, and the efficacy of these agents was assessed after the short-term administration of NSAID. Furthermore, no drugs have been proved to be effective in the treatment of NSAID-induced severe enteropathy, which is thought to be a clinically significant condition. Therefore, therapies targeting TNF α , and particularly anti-TNF biological therapy, should be considered as a novel treatment for NSAID-induced severe small intestinal pathologies such as bleeding ulcers and large ulcers. Considering that anti-TNF therapy is expensive and NSAID-induced damage in the small intestine, similar to that observed in the upper gastrointestinal tract, is believed to heal spontaneously after the cessation of NSAID administration, patients with severe enteropathy for whom NSAID administration cannot be discontinued (eg, low-dose aspirin) may be candidates for anti-TNF biological therapies.

In the present study, the patients receiving anti-TNF therapy had a tendency to use a selective COX-2 inhibitor, celecoxib, more frequently compared to those who did not receive such therapy, although the difference between these two groups did not reach statistical significance. Previous studies have demonstrated that the short-term administration of selective COX-2 inhibitors had less ulcerogenic effects on the small intestine than that observed by the administration of non-selective COX inhibitors.^{4 24} No difference in the incidence of injury to the small intestine was observed between chronic users of the two types of NSAID.²⁵ Therefore, long-term use of selective COX-2 inhibitors may abolish their beneficial effect on the small intestine, and the chronic use of both types of NSAID may result in similar intestinal ulcerogenicity. Furthermore, propensity score matching resulted in a similar rate of celecoxib use between the

anti-TNF therapy and anti-TNF therapy-naïve groups and yielded results similar to those obtained before propensity score matching in terms of the protective effects of anti-TNF therapy, suggesting that this prescription bias did not affect our main results.

In addition to NSAID, several anti-RA drugs such as steroids and methotrexate have been reported to induce upper gastrointestinal injuries.^{26 27} Although the small intestinal toxicity of these drugs remains unclear, concomitant use of these drugs and NSAID might increase the severity of enteropathy. However, in the present study, compared to the anti-TNF therapy-naïve group, the rates of the concomitant use of NSAID and steroids or methotrexate were higher in the anti-TNF therapy group, in which severe NSAID-induced small bowel damage was reduced. Therefore, the bias in the prescription of these anti-RA drugs would also not affect our main result that anti-TNF therapy reduced severe NSAID-induced enteropathy.

Recent animal studies indicated that both PPI and H₂RA exacerbated NSAID-induced small intestinal injury^{28 29} and our clinical study demonstrated that the use of PPI and H₂RA was associated with an increased risk of severe NSAID-induced small intestinal damage with OR of 5.22 and 3.95, respectively.³⁰ Therefore, in the present study, we included the use of acid suppressants (PPI or H₂RA) in the confounding variables and found that the preventive effect of anti-TNF therapy on NSAID-induced severe small intestinal damage remained robust to adjustment for the use of acid suppressants, thus further confirming the reduction of the risk for NSAID-induced severe enteropathy through anti-TNF therapy.

This study had several limitations. First, as the number of patients was relatively small, the OR 95% CI of the relationship between the use of anti-TNF therapy and the development of severe damage or most severe damage was wide, allowing for the possibility that the protective effect of anti-TNF therapy on severe damage, although significant, may be of minor clinical importance. Second, in this study, some patients received non-selective COX inhibitors, whereas others were being treated with a selective COX-2 inhibitor, celecoxib. Although small intestinal ulcerogenicity seems to be similar between non-selective NSAID and selective COX-2 inhibitors in chronic users, data obtained from a study using a single non-selective NSAID would be more convincing. Third, in the propensity score matching analysis, we employed a 1:1 matching without replacement by using an optimal matching technique. Therefore, of the 59 patients in the anti-TNF therapy-naïve group, 23 patients (39.9%) were unmatched and not accounted for in the conditional logistic regression analysis. A better matching technique could have improved the current propensity score matching. Fourth, in the present study, the ASD values ranged from 6% to 35%. In general, ASD values less than 10% indicate a balancing of baseline variables between treated and untreated cohorts;³¹ therefore, the ASD values noted in this study indicate that our propensity score matching could not completely remove the imbalance of variables, possibly because of the relatively small number of patients in both groups. To resolve these limitations, prospective, randomised, large-scale studies using a single NSAID to evaluate the effect of anti-TNF therapy on NSAID-induced severe enteropathy are needed.

In conclusion, these results suggest that anti-TNF therapy may protect against NSAID-induced small intestinal damage in RA patients. Drugs targeting TNF α may be effective in the treatment of NSAID-induced enteropathy. Prospective studies with repeated VCE before and after anti-TNF therapy are needed to validate our findings.

Contributors Conception and design of the study: TW, TT, TK and TA; statistical analysis: MS; acquisition of the data: YN and SS; analysis and interpretation of the data: TW, YN, KT and YF; obtaining informed consent: TT, HY and KW; recruiting of patients: TK; drafting of the manuscript: TW and TT; critical revision of the manuscript for important intellectual content: TW and TA.

Competing interests None.

Ethics approval The protocol for the study was approved by the ethics committee of the Osaka City University Graduate School of Medicine.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Iddan G, Meron G, Glukhovskiy A, *et al.* Wireless capsule endoscopy. *Nature* 2000;405:417.
- Maiden L, Thjodleifsson B, Theodor A, *et al.* A quantitative analysis of NSAID-induced small bowel pathology by capsule enteroscopy. *Gastroenterology* 2005;128:1172–8.
- Watanabe T, Tanigawa T, Nadatani Y, *et al.* Mitochondrial disorders in NSAIDs-induced small bowel injury. *J Clin Biochem Nutr* 2011;48:117–21.
- Goldstein JL, Eisen GM, Lewis B, *et al.* Video capsule endoscopy to prospectively assess small bowel injury with celecoxib, naproxen plus omeprazole, and placebo. *Clin Gastroenterol Hepatol* 2005;3:133–41.
- Watanabe T, Sugimori S, Kameda N, *et al.* Small bowel injury by low-dose enteric-coated aspirin and treatment with misoprostol: a pilot study. *Clin Gastroenterol Hepatol* 2008;6:1279–82.
- Sugimori S, Watanabe T, Tabuchi M, *et al.* Evaluation of small bowel injury in patients with rheumatoid arthritis by capsule endoscopy: effects of anti-rheumatoid arthritis drugs. *Digestion* 2008;78:208–13.
- Braun J, Sieper J. Overview of the use of the anti-TNF agent infliximab in chronic inflammatory diseases. *Expert Opin Biol Ther* 2003;3:141–68.
- Elliott MJ, Maini RN, Feldmann M, *et al.* Randomised double-blind comparison of chimeric monoclonal antibody to tumour necrosis factor alpha (cA2) versus placebo in rheumatoid arthritis. *Lancet* 1994;344:1105–10.
- Moreland LW, Schiff MH, Baumgartner SW, *et al.* Etanercept therapy in rheumatoid arthritis. A randomized, controlled trial. *Ann Intern Med* 1999;130:478–86.
- Santucci L, Fiorucci S, Giansanti M, *et al.* Pentoxifylline prevents indomethacin induced acute gastric mucosal damage in rats: role of tumour necrosis factor alpha. *Gut* 1994;35:909–15.
- Appleyard CB, McCafferty DM, Tigley AW, *et al.* Tumor necrosis factor mediation of NSAID-induced gastric damage: role of leukocyte adherence. *Am J Physiol* 1996;270:G42–8.
- Watanabe T, Higuchi K, Kobata A, *et al.* Non-steroidal anti-inflammatory drug-induced small intestinal damage is Toll-like receptor 4 dependent. *Gut* 2008;57:181–7.
- Fukumoto K, Naito Y, Takagi T, *et al.* Role of tumor necrosis factor-alpha in the pathogenesis of indomethacin-induced small intestinal injury in mice. *Int J Mol Med* 2011;27:353–9.
- Graham DY, Opekun AR, Willingham FF, *et al.* Visible small-intestinal mucosal injury in chronic NSAID users. *Clin Gastroenterol Hepatol* 2005;3:55–9.
- Rubin DB. Estimating causal effects from large data sets using propensity scores. *Ann Intern Med* 1997;127:757–63.
- D'Agostino RB Jr. Propensity score methods for bias reduction in the comparison of a treatment to a non-randomized control group. *Stat Med* 1998;17:2265–81.
- Peyrin-Birolet L, Deltenre P, de Suray N, *et al.* Efficacy and safety of tumor necrosis factor antagonists in Crohn's disease: meta-analysis of placebo-controlled trials. *Clin Gastroenterol Hepatol* 2008;6:644–53.
- Ford AC, Sandborn WJ, Khan KJ, *et al.* Efficacy of biological therapies in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol* 2011;106:644–59; quiz 60.
- Melikoglu M, Fresko I, Mat C, *et al.* Short-term trial of etanercept in Behcet's disease: a double blind, placebo controlled study. *J Rheumatol* 2005;32:98–105.
- Arakawa T, Watanabe T, Tanigawa T, *et al.* Small intestinal injury caused by NSAIDs/aspirin: finding new from old. *Curr Med Chem* 2012;19:77–81.
- Watanabe T, Nishio H, Tanigawa T, *et al.* Probiotic *Lactobacillus casei* strain Shirota prevents indomethacin-induced small intestinal injury: involvement of lactic acid. *Am J Physiol Gastrointest Liver Physiol* 2009;297:G506–13.
- Fujimori S, Seo T, Gudis K, *et al.* Prevention of nonsteroidal anti-inflammatory drug-induced small-intestinal injury by prostaglandin: a pilot randomized controlled trial evaluated by capsule endoscopy. *Gastrointest Endosc* 2009;69:1339–46.
- Niwa Y, Nakamura M, Ohmiya N, *et al.* Efficacy of rebamipide for diclofenac-induced small-intestinal mucosal injuries in healthy subjects: a prospective, randomized, double-blinded, placebo-controlled, cross-over study. *J Gastroenterol* 2008;43:270–6.

Small bowel

- 24 Hawkey CJ, Ell C, Simon B, *et al.* Less small-bowel injury with lumiracoxib compared with naproxen plus omeprazole. *Clin Gastroenterol Hepatol* 2008;6:536–44.
- 25 Maiden L, Thjodleifsson B, Seigal A, *et al.* Long-term effects of nonsteroidal anti-inflammatory drugs and cyclooxygenase-2 selective agents on the small bowel: a cross-sectional capsule enteroscopy study. *Clin Gastroenterol Hepatol* 2007;5:1040–5.
- 26 Laine L, Bombardier C, Hawkey CJ, *et al.* Stratifying the risk of NSAID-related upper gastrointestinal clinical events: results of a double-blind outcomes study in patients with rheumatoid arthritis. *Gastroenterology* 2002;123:1006–12.
- 27 Sartori S, Trevisani L, Nielsen I, *et al.* Randomized trial of omeprazole or ranitidine versus placebo in the prevention of chemotherapy-induced gastroduodenal injury. *J Clin Oncol* 2000;18:463–7.
- 28 Wallace JL, Syer S, Denou E, *et al.* Proton pump inhibitors exacerbate NSAID-induced small intestinal injury by inducing dysbiosis. *Gastroenterology* 2011;141:1314–22; 22 e1–5.
- 29 Satoh H, Amagase K, Takeuchi K. Exacerbation of nonsteroidal anti-inflammatory drug-induced small intestinal lesions by antisecretory drugs in rats: the role of intestinal motility. *J Pharmacol Exp Ther* 2012;343:270–7.
- 30 Watanabe T, Tanigawa T, Nadatani Y, *et al.* Risk factors for severe nonsteroidal anti-inflammatory drug-induced small intestinal damage. *Dig Liver Dis* 2013;45:390–5.
- 31 Austin PC. Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples. *Stat Med* 2009;28:3083–107.



Anti-tumour necrosis factor agents reduce non-steroidal anti-inflammatory drug-induced small bowel injury in rheumatoid arthritis patients

Toshio Watanabe, Tetsuya Tanigawa, Masatsugu Shiba, et al.

Gut published online May 22, 2013
doi: 10.1136/gutjnl-2013-304713

Updated information and services can be found at:
<http://gut.bmj.com/content/early/2013/05/21/gutjnl-2013-304713.full.html>

These include:

References

This article cites 31 articles, 7 of which can be accessed free at:
<http://gut.bmj.com/content/early/2013/05/21/gutjnl-2013-304713.full.html#ref-list-1>

P<P

Published online May 22, 2013 in advance of the print journal.

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Endoscopy](#) (959 articles)

Notes

Advance online articles have been peer reviewed, accepted for publication, edited and typeset, but have not yet appeared in the paper journal. Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>