











## Active sites of nitrogen-doped carbon materials for oxygen reduction reaction clarified using model catalysts

Donghui Guo, Riku Shibuya, Chisato Akiba, Shunsuke Saji, Takahiro Kondo and Junji Nakamura

*Science* **351** (6271), 361-365  
DOI: 10.1126/science.aad0832

### The right kind of dopant

The oxygen reduction reaction is an important step in fuel cells and other electrochemical processes but is still largely dependent on precious metal-containing catalysts. Recently explored alternatives include carbon materials that are doped with different, preferably non-precious metal, atoms. Guo *et al.* studied model graphite catalysts to try to understand the role of nitrogen doping and to elucidate the active catalytic sites. A nitrogen atom bound to two carbons formed an active catalyst site with an activity rivaling that of N-doped graphene catalysts.

*Science*, this issue p. 361

### ARTICLE TOOLS

<http://science.sciencemag.org/content/351/6271/361>

### SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2016/01/20/351.6271.361.DC1>

### REFERENCES

This article cites 21 articles, 2 of which you can access for free  
<http://science.sciencemag.org/content/351/6271/361#BIBL>

### PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)