

RESULTS

	MEAN(yrs)	STD DEV(yrs)	p-value
Early Amniotomy	25.91	5.45	0.9606
Late Amniotomy	25.96	4.77	

Table 1: Table showing the age distribution in the study population.




Figure 1: Chart showing the age distribution in the study population.

AGE GROUP	EARLY (n=70)	(%)	LATE (n=70)	(%)
<21	14	20	13	18.57
21-30	40	57.14	45	64.29
>30	16	22.86	12	17.14

Table 2: Table showing the distribution of women in different age groups in the study population




Figure 2:

	MEAN	STD DEV	p-value
Early Amniotomy	24.10	3.09	0.3531
Late Amniotomy	24.60	3.24	

Table 3: Table showing the BMI distribution in the study population.




Figure 3:

	MEAN(wks)	STD DEV(wks)	p-value
Early Amniotomy	38.61	1.17	0.3319
Late Amniotomy	38.43	1.08	

Table 4: Table showing the distribution of gestational age in the study population.




Figure 4:

INDICATION FOR IOL	EARLY (n=70)	(%)	LATE (n=70)	(%)
GDM	22	31.43	25	35.71
Post dates	22	31.43	15	21.43
Hypertension	3	4.29	2	2.86
Preeclampsia	7	10	6	8.57
Oligohydramnios	16	22.86	22	31.43

Table 5: Table showing the various indication for induction of labour in the study population.




Figure 5:

	MEAN	STD DEV	p-value
Early Amniotomy	6.37	0.49	0.6067
Late Amniotomy	6.41	0.50	

Table 6: Table showing the distribution of bishop's score in the study population.




Figure 6:

	MEAN (hrs:min)	STD DEV (hrs:min)	p-value
Early Amniotomy	3:49	1:30	<0.0001
Late Amniotomy	7:58	2:57	

Table 7: Table showing the time taken to reach active phase in the study population.

Time to reach Active Phase of Labour




Figure 7:

	OXYOCIN INDUCTION TO DELIVERY(hrs:min)	STD DEV (hrs:min)	p-value
Early	2:16	1:28	0.2794
Late	1:59	1:20	

Table 8: Table showing the time interval between oxytocin induction and delivery in the study population.

Time Interval between Oxytocin Induction and Delivery

p value = 0.2794




Figure 8:

	MEAN (hrs:min)	STD DEV (hrs:min)	p-value
Early Amniotomy	6:06	2:25	<0.0001
Late Amniotomy	7:58	2:57	

Table 9: Table showing the time taken from induction to delivery in the study population.




Figure 9:

MODE OF DELIVERY	EARLY (n=70)	(%)	LATE (n=70)	(%)
Vaginal	44	62.86	31	44.29
Caesarean	24	34.29	38	54.29
Instrumental Delivery	2	2.86	1	1.43

Table 10: Table showing the distribution of women in different age groups in the study population.




Figure 10:

CESAREAN INDICATION	EARLY (n=24)	(%)	LATE (n=38)	(%)
Failed induction	6	25	15	39.47
Fetal distress	14	58.33	13	34.21
Arrest of labour	4	16.67	10	26.32

Table 11: Table showing the various indications for caesarean section the study population.




Figure 11:

CTG ABNORMALITY	EARLY (n=70)	%	LATE (n=70)	%
YES	6	8.57	15	21.43
NO	64	91.43	55	78.57

Table 12: Table showing the incidence of Cardiotocographic abnormality in the study population.




Figure 12:

UTERINE HYPERSTIMULATION	EARLY (n=70)	%	LATE (n=70)	%
YES	1	1.43	0	0
NO	69	98.57	70	100

Table 13: Table showing the incidence of post uterine hyperstimulation in the study population.




Figure 13:

Post Partum Hemorrhage	EARLY (n=70)	%	LATE (n=70)	%
YES	5	7.14	9	12.86
NO	65	92.86	61	87.14

Table 14: Table showing the incidence of post partum hemorrhage in the study population.




Figure 14:

	MEAN(mL)	STD DEV(mL)	p-value
Early Amniotomy	464.64	210.44	0.0352
Late Amniotomy	543.21	226.38	

Table 15: Table showing the the distribution of blood loss during delivery in the study population.




Figure 15:

	MEAN(mL)	STD DEV(mL)	p-value
Early Amniotomy	666.67	230.74	0.8114
Late Amniotomy	651.97	237.91	

Table 16: Table showing the the distribution of blood loss during delivery in women undergoing caesarean section in the study population.




Figure 16:

	MEAN(mL)	STD DEV(mL)	p-value
Early Amniotomy	359.24	86.98	0.0228
Late Amniotomy	414.06	121.64	

Table 17: Table showing the the distribution of blood loss during delivery in women undergoing normal vaginal delivery in the study population.




Figure 17:

MATERNAL FEVER	EARLY (n=70)	%	LATE (n=70)	%
YES	1	1.43	2	2.86
NO	69	98.57	68	97.14

Table 18: Table showing the incidence of Maternal fever in the study population.




Figure 18:

NICU ADMISSION	EARLY (n=70)	%	LATE (n=70)	%
YES	13	18.57	22	31.43
NO	57	81.43	48	68.57

Table 19: Table showing the incidence of NICU admission in the study population.

Comparison of incidence of NICU Admissions




Figure 19:

	MEAN(g)	STD DEV(g)	p-value
Early Amniotomy	3017.34	294.58	0.1597
Late Amniotomy	2947.97	285.91	

Table 20: Table showing the the distribution of birth weight of the newborns in the study population.




Figure 20:

APGAR	EARLY (n=70)	(%)	LATE (n=70)	(%)
≤ 7	21	30	24	34.29
> 7	49	70	46	65.71

Table 21: Table showing the APGAR scores in the study population.

Comparison of APGAR Scores between the Sutdy populations




Figure 21:

Comparison of APGAR Scores between the Sutdy populations




Figure 22:

STILL BIRTH	EARLY(n=70)	LATE(n=70)
YES	0	0
NO	70	70

Table 22: Table showing the incidence of stillbirth in the study population.




Figure 23: