

Leinster - Basic Category Theory - Selected problem solutions for Chapter 2

Adam Barber

October 26, 2021

2.1.16

(a) Interesting adjoint functors to G -sets.

The trivial group functor I sends a set to a \mathbf{G} -set with the trivial action $gx = x$.
Interesting functors

Orbit functor sends a G -set with underlying set elements a of A to:

$$A_G = \{g \cdot a, g \in G\}$$

Fixed point functor sends a G -set with underlying set elements a of A to:

$$A^G = \{a \text{ such that } g \cdot a = a \text{ for all } g \in G, a \in A\}$$

Fixed point functor - right adjoint Morphisms in a G -set are functions on the underlying set, where f commutes with g for every $g \in G$.

There is a bijection for each $A \in \mathbf{Set}$ and $B \in [G, \mathbf{Set}]$ as follows

$$\begin{aligned} [G, \mathbf{Set}](I(A), B) &\rightarrow \mathbf{Set}(A, B^G) \\ \psi &\mapsto \bar{\psi} \end{aligned}$$

$\bar{\psi}$ sends each element a of A to $\psi(a)$ if $g \cdot a = a$, otherwise it sends a to $\psi(\emptyset)$.

$$\begin{aligned} \mathbf{Set}(A, B^G) &\rightarrow [G, \mathbf{Set}](I(A), B) \\ \phi &\mapsto \bar{\phi} \end{aligned}$$

ϕ sends each $a \in A$ in the underlying set of the G -set to the G -set $(g, \bar{\phi}(a)), g \in G$.

Orbit functor - left adjoint There is a bijection for each $A \in \mathbf{Set}$ and $B \in [G, \mathbf{Set}]$ as follows

$$\begin{aligned} \mathbf{Set}(A_G, B) &\rightarrow [G, \mathbf{Set}](A, I(B)) \\ \psi &\mapsto \bar{\psi} \end{aligned}$$

So each morphism in \mathbf{Set} sends the orbit set of an element a of A , call it a_G , to $\psi(a_G)$, where ψ is a function of sets. Set G -set morphism to $\bar{\psi} = \psi$, where $\bar{\psi}$ commutes with g for every g in G .

$$\begin{aligned} [G, \mathbf{Set}](A, I(B)) &\rightarrow \mathbf{Set}(A_G, B) \\ \phi &\mapsto \bar{\phi} \end{aligned}$$

Choose $\bar{\phi}$ to be a disjoint union of each orbit of a in A , $\bar{\phi}(a) = \coprod \{\phi(a), g \cdot a, g \in G\}$