

0.10:

Let S be a set. The indiscrete topological space $I(S)$ is the space whose set of points is S and whose only open subsets are \emptyset and S . To find a universal property satisfied by the space $I(S)$ proceed as follows. With this topology any map from a topological space to S is continuous.

Parroting the wording of the question, let us rephrase this in universal parlance. Define a function $i : S \rightarrow I(S)$, by $i(s) = s, s \in S$. Then $I(S)$ has the following property.

$$\begin{array}{ccc} S & \xrightarrow{i} & I(S) \\ & \nwarrow \forall f & \uparrow \bar{f} \\ & & X \end{array}$$

For all topological spaces X and all functions $f : X \rightarrow S$ there exists a unique continuous map $\bar{f} : X \rightarrow I(S)$. What it says is all maps into an indiscrete space are continuous. It also says that given S , the universal property determines $I(S)$ and i , up to isomorphism.

0.11

The universal property that is satisfied by the pair $(\ker(\theta), \iota)$ is depicted in the diagram below.

$$\begin{array}{ccccc} \ker(\theta) & \xrightarrow{\iota} & G & \xrightarrow[\epsilon]{\theta} & H \\ \exists! \bar{f} \uparrow & & \nearrow f & & \\ F & & & & \end{array}$$

The statement of the universal property is as follows. For any $f : F \rightarrow G$ such that $\theta \circ f = \epsilon \circ f$ there is a unique $\bar{f} : F \rightarrow \ker(\theta)$ such that the diagram above commutes. That is $f = \iota \circ \bar{f}$.

0.13:

(a)

Choose $\phi(\sum_{i=1}^n a_i x^i) = \sum_{i=1}^n a_i r^i$. Then ϕ with $\phi(x) = r$ is a homomorphism that satisfies additive and multiplicative properties. To prove uniqueness assume there is another homomorphism ψ , with $\psi(x) = r$. Then $\psi(\sum_{i=1}^n a_i x^i) = \sum_{i=1}^n a_i \psi(x)^i = \sum_{i=1}^n a_i r^i$ by properties of a homomorphism. So $\psi = \phi$.

(b)

$\iota : \mathbb{Z}[x] \rightarrow A$ maps $\sum_{i=1}^n p_i x^i$ to $\sum_{i=1}^n p_i a^i$, using $\iota(x) = a$, the multiplicative property of a homomorphism to get $\iota(x^i) = \iota(x)^i$, and the additive property to get $\iota(p_i) \iota(x)^i = p_i \iota(x)^i$ remembering p_i is in \mathbb{Z} .

Going in the direction $A \rightarrow \mathbb{Z}[x]$ we know as provided in (b) that, taking $R = \mathbb{Z}[x]$, and $\phi = \iota'$, there exists a unique ring homomorphism such that $\iota'(a) = x$. So ι' maps $\sum_{i=1}^n p_i a^i$ to $\sum_{i=1}^n p_i x^i$ and $\iota' \circ \iota = 1_{\mathbb{Z}[x]}$. Also using definitions of ι and ι' easily yields $\iota \circ \iota' = 1_A$.