

AMITA MUNOT
294 Sindh Society Aundh

Reference:Dr.--

SID: 120143692

REPORT

Tel No: 919822033212
PID: 196650

Age:65.20 Years Sex:FEMALE

Collection Date:
01-12-2020 10:08 AM
Registration Date:
01-12-2020 10:08 am
Report Date:
01-12-2020 03:31 PM

Test Description	Observed Value	Biological Reference Interval
------------------	----------------	-------------------------------

TEST NAME

Glycated Hemoglobin (HbA1C), by HPLC	6.00	4.0 to 5.6 %
--------------------------------------	-------------	--------------

On follow up.

Interpretation :

HbA1C level reflects the mean glucose concentration over previous 8-12 weeks and provides better indication of long term glycemic control.

For diagnosis of Diabetes Mellitus (≥ 18 yrs of age) :

5.7 % - 6.4 % : Increased risk for developing diabetes.

≥ 6.5 % : Diabetes

Therapeutic goals for glycemic control :

Adults : < 7%

Toddlers and Preschoolers : < 8.5% (but > 7.5 %)

School age (6-12 yrs) : < 8%

Adolescents and young adults (13 - 19 yrs) : < 7.5 %

Levels of HbA1C may be low as result of shortened RBC life span in case of hemolytic anemia.

Increased HbA1C values may be found in patients with polycythemia or post splenectomy patients.

Patients with Homozygous forms of rare variant Hb(CC,SS,EE,SC) HbA1c can not be quantitated as there is no HbA. In such circumstances glycemic control can be monitored using plasma glucose levels or serum Fructosamine.

The A1c target should be individualized based on numerous factors, such as age, life expectancy, comorbid conditions, duration of diabetes, risk of hypoglycemia or adverse consequences from hypoglycemia, patient motivation and adherence.

Ref : ADA (Standards of Medical Care in Diabetes - 2017)



Page 1 of 4

Golwilkar

Dr.(Mrs.) Vinanti Golwilkar-Patankar
(MD Path) Regn.No.:2008/03/0772
A.G Diagnostics Pvt. Ltd.

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ www.nabl-india.org"

Carrying forward
Dr. Ajit Golwilkar's
legacy of Over
Four Decades

DIAGNOSTICS
BE SURE
BE WELL
ए.जी. डायग्नॉस्टिक्स प्रा. लि. A.G Diagnostics Pvt. Ltd.

Dr. Awanti Golwilkar
MD (Pathology)

Dr. Vinanti Golwilkar
MD (Pathology)

REPORT

AMITA MUNOT
294 Sindh Society Aundh

Tel No: 919822033212
PID: 196650

Age:65.20 Years Sex:FEMALE

Reference:Dr.--

SID: 120143692

Collection Date:
01-12-2020 10:08 AM
Registration Date:
01-12-2020 10:08 am
Report Date:
01-12-2020 03:31 PM

Test Description

Plasma Glucose :

Plasma glucose fasting, by Hexokinase method

Observed Value

83

Biological Reference Interval

< 100 mg/dL
100 to 125 mg/dL : Impaired fasting
glucose tolerance / Prediabetes
>= 126 mg/dL : Suggestive of
diabetes mellitus
(On more than one occasion)
American Diabetes Association
Guidelines 2020



Golwilkar

Dr.(Mrs.) Vinanti Golwilkar-Patankar
(MD Path) Regn.No.:2008/03/0772
A.G Diagnostics Pvt. Ltd.

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ www.agdiagnostics.com"

AMITA MUNOT
294 Sindh Society Aundh

Reference: Dr.--

SID: 120143692

REPORT

Tel No: 919822033212
PID: 196650

Age: 65.20 Years Sex: FEMALE

Collection Date:
01-12-2020 10:08 AM
Registration Date:
01-12-2020 10:08 am
Report Date:
01-12-2020 03:31 PM

Test Description	Observed Value	Biological Reference Interval
TEST NAME		
25 - OH Vitamin D, serum by CMLA	25.90	Severe deficiency : < 10 ng/mL Mild to moderate deficiency : 10 to 19 ng/mL Optimum levels : 20 to 50 ng/mL Increased risk of hypercalciuria: 51 to 80 ng/mL Toxicity possible : > 80 ng/mL Ref. : Mayo Medical Laboratories These reference ranges represent clinical decision values, based on the 2011 Institute of Medicine report

Interpretation :

Vitamin D is vital for strong bones. It also has important, emerging roles in immune function and cancer prevention.

Vitamin D compounds in the body are exogenously derived by dietary means; from plants as 25-hydroxyvitamin D2 (ergocalciferol or calciferol) or from animal products as 25-hydroxyvitamin D3 (cholecalciferol or calcidiol).

Vitamin D may also be endogenously derived by conversion of 7-dihydrocholesterol to 25-hydroxyvitamin D3 in the skin upon ultraviolet exposure.

The total 25-hydroxyvitamin D (25-OH-VitD) level (the sum of 25-OH-vitamin D2 and 25-OH-vitamin D3) is the appropriate indicator of vitamin D body stores.

Patients with renal failure can have very high 25-OH-VitD levels without any signs of toxicity, as renal conversion to the active hormone 1,25-OH-VitD is impaired or absent.

Kindly correlate clinically, with supplementation history & repeat with fresh sample if necessary.



Golwilkar

Dr. (Mrs.) Vinanti Golwilkar-Patankar
(MD Path) Regn.No.:2008/03/0772
A.G. Diagnostics Pvt. Ltd.

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ www.nabl-india.org"

AMITA MUNOT
294 Sindh Society Aundh

Reference: Dr.--

SID: 120143692

REPORT

Tel No: 919822033212
PID: 196650

Age: 65.20 Years Sex: FEMALE

Collection Date:
01-12-2020 10:08 AM
Registration Date:
01-12-2020 10:08 am
Report Date:
01-12-2020 03:31 PM

Test Description	Observed Value	Biological Reference Interval
CRP(hs) - C- Reactive Protein high sensitivity	6.77	See clinical information below Method : Nephelometry / Immunoturbidimetry

Clinical Information :

1. C-reactive protein (CRP) is a biomarker of inflammation. Plasma CRP concentrations increase rapidly and dramatically (100-fold or more) in response to tissue injury or inflammation.

2. High-sensitivity CRP (hs-CRP) is more precise than standard CRP when measuring baseline (i.e. normal) concentrations and enables a measure of chronic inflammation. It is recommended for cardiovascular risk assessment. Atherosclerosis is an inflammatory disease and hs-CRP has been endorsed by multiple guidelines as a biomarker of atherosclerotic cardiovascular disease risk.

Low cardiovascular risk : < 2.0 mg/L
High cardiovascular risk : \geq 2.0 mg/L
Acute inflammation : > 10.0 mg/L

3. A single test for high-sensitivity CRP (hs-CRP) may not reflect an individual patient's basal hs-CRP level. Repeat measurement may be required to firmly establish an individual's basal hs-CRP concentration. The lowest of the measurements should be used as the predictive value.

Reference : Mayo Medical Laboratories

End of Report

Dr. (Mrs.) Vinanti Golwilkar-Patankar
(MD Path) Regn.No.: 2008/03/0772
A.G Diagnostics Pvt. Ltd.

Page 4 of 4

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ www.nabl-india.org"

Carrying forward
Dr. Ajit Golwilkar's
legacy of Over
Four Decades

DIAGNOSTICS
BE SURE
BE WELL
ए.जी. डायग्नॉस्टिक्स प्रा. लि. A.G Diagnostics Pvt. Ltd.

Dr. Awanti Golwilkar
MD (Pathology)

Dr. Vinanti Golwilkar
MD (Pathology)