



Name : Mr. RACHIT GUPTA

Collected

: 13/1/2020 9:03:00AM

Lab No. : 150667093

Age: 35 Years

Received Reported : 13/1/2020 9:19:57AM : 14/1/2020 12:06:15PM

A/c Status : P

Ref By: SELF

Report Status : Final

Results	Units	Bio. Ref. Interval
95.00	mg/dL	70.00 - 100.00
16.50	uU/mL	2.00 - 25.00
143.90	%	
46.80	%	
2.14		<2.50
	95.00 16.50 143.90 46.80	95.00 mg/dL 16.50 uU/mL 143.90 % 46.80 %

Gender:

Male

Note

- 1. As insulin secretion is pulsatile, it is recommended to take mean of three samples at 5 minute intervals to compute HOMA accurately.
- 2. This assay cannot be used to assess beta cell function in those taking exogenous insulin. In such patients HOMA-IR, C-peptide Model is recommended.
- 3. The HOMA IR calculator version 2.2 accepts values only in following validated ranges, Insulin (2.9-57.6uU/mL) and Glucose (54.1-450.5 mg/dL).

Comment

Homeostatic model assessment (HOMA) is a method for assessing beta cell function (%B)and insulin sensitivity (%S) from fasting glucose and insulin concentrations. HOMA can be used to track changes in insulin sensitivity and beta cell function to examine natural history of diabetes. Insulin sensitivity is reduced in normal subjects having first degree relative with type 2 diabetes compared with control subjects. Changes in beta cell sensitivity in subjects on insulin secretogogues may be useful in determining beta cell function over a period.

Usage

- To assess risk of developing diabetes
- To assess response to treatment





S04 - SDA HOME VISIT 0

DELHI



Name : Mr. RACHIT GUPTA

Lab No. : 150667093 Age: 35 Years

Gender: Male

Collected : 13/1/2020 9:03:00AM

Received : 13/1/2020 9:19:57AM Reported : 14/1/2020 12:06:15PM

A/c Status : P Ref By : SELF Report Status : Final

Test Name	Results	Units	Bio. Ref. Interval
HEMOGLOBIN HPLC/ELECTROPHORESIS @ (HPLC)			
Hb F	<1.00	%	<1.50
Peak 2	4.40	%	<9.60
Hb Adult	85.80	%	83.24 - 90.79
Hb A2	2.50	%	1.50 - 3.50
Others (Non Specific)	7.00	%	<10.00
Hemoglobin	15.20	g/dL	13.00 - 17.00
RBC Count	5.48	mill/mm3	4.50 - 5.50
Packed Cell Volume (PCV)	47.10	%	40.00 - 50.00
MCV	85.90	fL	80.00 - 100.00
мсн	27.70	pg	27.00 - 32.00
RDW	13.00	%	11.50 - 14.50

Suggestive Interpretation Normal Hb chromatographic pattern







Name : Mr. RACHIT GUPTA

Collected

: 13/1/2020 9:03:00AM

Lab No. ;

150667093

Gender: Male

Received : 13/1 Reported : 14/1

: 13/1/2020 9:19:57AM : 14/1/2020 12:06:15PM

A/c Status : P Ref By : SELF

Age: 35 Years

Report Status : Final

Test Name Results Units Bio. Ref. Interval

IMPORTANT INSTRUCTIONS

*Test results released pertain to the specimen submitted.*All test results are dependent on the quality of the sample received by the Laboratory.

*Laboratory investigations are only a tool to facilitate in arriving at a diagnosis and should be clinically correlated by the Referring Physician.*Sample repeats are accepted on request of Referring Physician within 7 days post reporting.*Report delivery may be delayed due to unforeseen circumstances. Inconvenience is regretted.*Certain tests may require further testing at additional cost for derivation of exact value. Kindly submit request within 72 hours post reporting.*Test results may show interlaboratory variations.*The Courts/Forum at Delhi shall have exclusive jurisdiction in all disputes/claims concerning the test(s) & or results of test(s).*Test results are not valid for medico legal purposes. *Contact customer care Tel No. +91-11-39885050 for all queries related to test results.

(#) Sample drawn from outside source.





S04 - SDA HOME VISIT

DELHI



Name : Mr. RACHIT GUPTA Collected : 13/1/2020 9:03:00AM

Lab No. : 150667093 Age: 35 Years Gender: Male Received : 13/1/2020 9:19:57AM Reported : 14/1/2020 12:06:15PM

A/c Status : P Ref By : SELF Report Status : Final

Test Name Results Units Bio. Ref. Interval

HIV 1 & 2 ANTIBODIES SCREENING TEST, SERUM @

| Final Result : Negative |

HIV 1 / 2 & P 24 COMBO TEST

(CMIA)

Index Value 0.08

Result Non Reactive

Interpretation

RESULT IN INDEX	REMARKS
< 1.00	Non Reactive
>= 1.00	Provisionally Reactive

Comments

Negative result implies that antibodies to HIV 1 / 2 have not been detected in the sample. This means the patient has either not been exposed to HIV 1 / 2 infection or the sample has been tested during the "window phase" i.e. before the development of detectable levels of antibodies. Hence a Negative result does not exclude the possibility of exposure or infection with HIV 1 / 2.

Recommendations

- 1. Results to be clinically correlated.
- 2. Rarely false negativity/positivity may occur.
- 3. Post test counseling available between 9.00 am to 5:00 pm at LPL laboratories.







Name : Mr. RACHIT GUPTA

Collected

: 13/1/2020 9:03:00AM

Lab No. : 150667093

093 Age: 35 Years

Received Reported : 13/1/2020 9:19:57AM : 14/1/2020 12:06:15PM

A/c Status : P Ref By : SELF Report Status : Final

Test Name Results Units Bio. Ref. Interval

Gender:

IMPORTANT INSTRUCTIONS

Male

*Test results released pertain to the specimen submitted.*All test results are dependent on the quality of the sample received by the Laboratory.

*Laboratory investigations are only a tool to facilitate in arriving at a diagnosis and should be clinically correlated by the Referring Physician .*Sample repeats are accepted on request of Referring Physician within 7 days post reporting.*Report delivery may be delayed due to unforeseen circumstances. Inconvenience is regretted.*Certain tests may require further testing at additional cost for derivation of exact value. Kindly submit request within 72 hours post reporting.*Test results may show interlaboratory variations.*The Courts/Forum at Delhi shall have exclusive jurisdiction in all disputes/claims concerning the test(s) & or results of test(s).*Test results are not valid for medico legal purposes. *Contact customer care Tel No. +91-11-39885050 for all queries related to test results.

(#) Sample drawn from outside source.







Name : Mr. RACHIT GUPTA

Collected

: 13/1/2020 9:03:00AM

Lab No. : 150667093

Age: 35 Years

Male Received Reported

: 13/1/2020 9:19:57AM : 14/1/2020 12:06:15PM

A/c Status : P

Ref By: SELF

Report Status : Final

Test Name	Results	Units	Bio. Ref. Interval
CARDIO C-REACTIVE PROTEIN (hsCRP), SERUM @	4.60	mg/L	<1.00
(Immunoturbidimetry)			

Gender:

Interpretation

CARDIO CRP IN mg/L	CARDIOVASCULAR RISK	
<1	Low	
1-3	Average	
3-10	High	
>10	Persistent elevation may represent Non cardiovascular inflammation	

Note: To assess vascular risk, it is recommended to test hsCRP levels 2 or more weeks apart and calculate the average

Comments

High sensitivity C Reactive Protein (hsCRP) significantly improves cardiovascular risk assessment as it is a strongest predictor of future coronary events. It reveals the risk of future Myocardial infarction and Stroke among healthy men and women, independent of traditional risk factors. It identifies patients at risk of first Myocardial infarction even with low to moderate lipid levels. The risk of recurrent cardiovascular events also correlates well with hsCRP levels. It is a powerful independent risk determinant in the prediction of incident Diabetes.

HEPATITIS B SURFACE ANTIGEN;HBsAg, SERL	M Non Reactive	Non Reactive
@		
(CMIA)		

*Specific Antibody Neutralization Assay is performed on all Reactive results.

Interpretation


ļ	RESULT	REMARKS
į	Reactive	Indicates presence of Hepatitis B Surface Antigen.
İ	Non-Reactive	Indicates absence of Hepatitis B Surface Antigen.

Note

1. This is a screening test and the result should be interpreted in conjunction with clinical findings and



Page 6 of 9







Name : Mr. RACHIT GUPTA Collected : 13/1/2020 9:03:00AM

A/c Status : P Ref By : SELF Report Status : Final

Test Name Results Units Bio. Ref. Interval

other diagnostic tests.

- 2. This assay is used for qualitative detection of Hepatitis B Surface Antigen (HBsAg) in serum samples and cannot differentiate between the stages of Hepatitis B viral infection. Detection of HBsAg may be observed in Hepatitis B viral infection for transient period of time after HBV vaccination.
- 3. Sensitivity and Specificity of the HBsAg test by CMIA technique is 100.0% and >99.5% respectively which is technically superior to other methodologies like ICT (Rapid cards), ELISA etc.
- 4. False positive results may be observed during pregnancy, presence of heterophilic antibodies in serum & patients receiving mouse monoclonal antibodies for diagnosis or therapy.
- 5. False negative reaction may be due to processing of sample collected early in the course of disease or presence of mutant forms of HBsAg.
- 6. For monitoring HBsAg levels, Quantitative HBsAg assay is recommended.
- 7. Test conducted on serum.

Comment

Hepatitis B Virus (HBV) is a member of the Hepadna virus family causing infections of the liver with variable clinical features. Hepatitis B is transmitted by blood and body fluids, sexually and from mother to fetus. In most cases HBV hepatitis is self limiting, but 1-2% adolescents and adults develop Chronic Hepatitis. Frequency of chronic HBV infection is 5-10% in immunocompromised patients and 80% in neonates.

HBsAg is the first serological marker after infection with HBV appearing 1-10 weeks after exposure and 2-8 weeks before the onset of clinical symptoms. HBsAg persists during the acute phase and disappears 12-20 weeks after onset of symptoms late in the convalescence period. Persistence of HBsAg for more than six months indicates development of carrier state or Chronic liver disease.

Uses

- Routine screening of blood and blood products to prevent transmission of Hepatitis B virus (HBV) to recipients
- To diagnose suspected HBV infection and monitor the status of infected individuals
- · For Prenatal Screening of pregnant women







Name : Mr. RACHIT GUPTA Collected : 13/1/2020 9:03:00AM

Received : 13/1/2020 9:19:57AM Lab No. : 150667093 Age: 35 Years Gender: Male Reported : 14/1/2020 12:06:15PM

A/c Status : P Ref By : SELF Report Status : Final

Test Name	Results	Units	Bio. Ref. Interval
HEPATITIS C ANTIBODY (Anti-HCV), SERUM @	0.06	Index	<1.00
(CMIA)			

Interpretation

RESULT (INDEX)	REMARKS
<1.00	Non Reactive / Not Detected
>=1.00	 Reactive/Asymptomatic/ Infective state/ Carrier state

Note

- 1. False positive results are seen Autoimmune diseases, Rheumatoid factor, Hypergammaglobulinemia, antibody Paraproteinemia, passive transfer, Antiidiotypes Anti superoxide dismutase
- 2. Supplemental testing is necessary in Reactive results with index value between 1.0 5.0 to identify and exclude biological false positive results
- 3. All reactive results should be verified by HCV RNA PCR to differentiate between past and present infection (as per CDC recommendation)
- 4. False negative results are seen in early Acute infection, Immunosuppression & Immuno-incompetence

Comments

Hepatitis C (HCV) is an RNA virus of Flavivirus group transmitted via blood transfusions, transplantation, injection drug users, accidental needle punctures in healthcare workers, dialysis patients and rarely from mother to infant.10% of new cases show sexual transmission. As compared to HAV & HBV, chronic infection with HCV occurs in 85% of infected individuals. In high risk populations, the predictive value of Anti HCV for HCV infection is > 99% whereas in low risk populations it is only 25%.

Uses

- Indicator of past or present infection, but does not differentiate between Acute / Chronic / Resolved infection
- Routine screening of low and high prevalence populations including blood donors



Page 8 of 9



Lab No.

A/c Status : P



Name Mr. RACHIT GUPTA

150667093

Age: 35 Years

Ref By: SELF

Male

Collected

: 13/1/2020 9:03:00AM

Received : 13/1/2020 9:19:57AM Reported : 14/1/2020 12:06:15PM

Report Status : Final

Test Name Results Units Bio. Ref. Interval

Gender:

Dr Anil Arora MD, Pathology

HOD Hematology & Immunohematology NRL - Dr Lal PathLabs Ltd

Dr Himangshu Mazumdar MD, Biochemistry Senior Consultant - Clinical Chemistry & Biochemical Genetics NRL - Dr Lal PathLabs Ltd

Dr Kamal Modi MD. Biochemistry

Consultant Biochemist NRL - Dr Lal PathLabs Ltd MD. Biochemistry National Head - Clinical Chemistry &

Biochemical Genetics NRL - Dr Lal PathLabs Ltd

----End of report ----

IMPORTANT INSTRUCTIONS

*Test results released pertain to the specimen submitted.*All test results are dependent on the quality of the sample received by the Laboratory *Laboratory investigations are only a tool to facilitate in arriving at a diagnosis and should be clinically correlated by the Referring Physician .*Sample repeats are accepted on request of Referring Physician within 7 days post reporting.*Report delivery may be delayed due to unforeseen circumstances. Inconvenience is regretted.*Certain tests may require further testing at additional cost for derivation of exact value. Kindly submit request within 72 hours post reporting.*Test results may show interlaboratory variations.*The Courts/Forum at Delhi shall have exclusive jurisdiction in all disputes/claims concerning the test(s) & or results of test(s).*Test results are not valid for medico legal purposes. *Contact customer care Tel No. +91-11-39885050 for all queries related to test results. (#) Sample drawn from outside source.

