PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000







PMF









Name: Mrs. Tannu Hodekar Age/Gender: 27 Ye

27 Year(s) 0 Months(s) 0 Day(s)/Female

Referred By: SELF Client Name: GEO MEDCARE HOSPITAL

**Collection Date:** 12-07-2021 11:40:00 **Report Release Date:** 12-07-2021 15:55:55

Sr.I	No Investigation	Observed Value	Reference Range	Unit	
Complete Haemogram Test					
Erythr	ocytes				
1	Total RBC	5.46	4.1-5.1	10^6/μL	
2	Hemoglobin	11.9	12.3-15.3	g/dL	
3	Hematocrit (PCV)	39.1	33-57	%	
4	Mean Corpuscular Volume (MCV)	71.6	80-96	$\mathbf{fL}$	
5	Mean Corpuscular Hemoglobin (MCH)	21.8	27.5-33.2	pg	
6	Mean Corpuscular Hemoglobin Concentration (MCHC)	30.4	29.4- 34.5	g/dL	
7	Red Cell Distribution Width (RDW-CV)	15.2	12-15	%	
8	Red Cell Distribution Width-SD(RDW-SD)	38.5	32-60.4	fl	
9	Nucleated Red Blood Cells	0.02	0 - 1.36	cells/μL	
10	Nucleated Red Blood Cells Percentage	0.1	0-4	%	
Platele	ts				
11	Platelet Count	257.0	150-450	10^3/μL	
12	Mean Platelet Volume (MPV)	8.4	6 - 12	fL	
13	Platelet Distribution Width (PDW)	16.9	15.5-18.3	%	
14	Plateletcrit (PCT)	0.215	0.12-0.37	%	
Leucoc	ytes				
15	<b>Total Leucocytes Count</b>	13.0	4.4-11	10^3/μL	
16	Neutrophils	71.5	40-77	%	
17	Lymphocyte Percentage	19.1	16-44	%	
18	Monocytes Percentage	3.9	2.0-10.0	%	
19	Eosinophils Percentage	4.9	0-7	%	
20	Basophils Percentage	0.6	0 - 1	%	
21	<b>Neutrophils-Absolute Count</b>	9.30	1.8-7.8	10^3/μL	
22	Lymphocytes-Absolute Count	2.48	1-4.8	$10^3/\mu$ L	
23	Monocytes-Absolute Count	0.51	0.1-1.0	$10^3/\mu$ L	
24	<b>Eosinophils-Absolute Count</b>	0.64	0-0.45	10^3/μL	
25	Basophils-Absolute Count	0.08	0-0.2	10^3/μL	



CRM No: 2873284

Sample Recd. Time: 12-07-2021 11:49

Report Time: 12-07-2021 15:55 Patient Name: Mrs. Tannu Hodekar

Patient ID: 2873284



Authorized Signatory Dr. Dina Abhani DCP, DNB (Pathology)





Scan For Report
Page 1 of 8

PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000

**Collection Date:** 



12-07-2021 11:40:00





PMF









Name: Mrs. Tannu Hodekar

Age/Gender:

27 Year(s) 0 Months(s) 0 Day(s)/Female

GEO MEDCARE HOSPITAL

Duy(8)/1 em

Referred By: SELF Client Name:

**Report Release Date:** 12-07-2021 15:55:55

Sr.No Investigation Observed Value Reference Range Unit

# **Peripheral Blood Smear**

26	RBC Morphology	Normocytic Normochromic
27	WBC Morphology	Leucocytosis
28	Platelets	Adequate On Smear



CRM No :2873284

Sample Recd. Time: 12-07-2021 11:49 Report Time: 12-07-2021 15:55 Patient Name: Mrs. Tannu Hodekar

Patient ID: 2873284

Davis

Authorized Signatory Dr. Dina Abhani DCP, DNB (Pathology)





PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000







PMF









Name: Mrs. Tannu Hodekar Age/Gender:

27 Year(s) 0 Months(s) 0 Day(s)/Female

Referred By: **SELF**  **Client Name:** 

GEO MEDCARE HOSPITAL

**Collection Date:** 12-07-2021 11:40:00 **Report Release Date:** 12-07-2021 15:55:55

Sr.No	Investigation	Observed Value	Reference Range	Unit
Liver F	unction Test			
1	Bilirubin Total Serum, Method: Jendrassik Grof	0.67	0.2-1.2	mg/ dL
2	Bilirubin Direct Serum, Method: Diazotization	0.14	0.01 - 0.4	mg/ dL
3	Bilirubin Indirect Serum, Method: Calculated	0.53	0.01-1.0	mg/dL
4	Aspartate Transaminase (AST/SGOT) Serum, Method: UV Kinetic with P5P	15.6	<35	U/L
5	Alanine Transaminase (ALT/SGPT) Serum, Method: UV Kinetic with P5P	7.2	<35	U/L
6	Alkaline Phosphatase Serum, Method: AMP – pNPP Kinetic	71.0	30 - 130	U/L
7	Total Protein Serum, Method: Biuret end point	6.22	6.4 - 8.2	g/dL
8	Albumin Serum, Method: Bromocresol Purple (BCP)	4.09	3.4 - 5	g/dL
9	Globulin Serum, Method: Calculated	2.13	1.9-3.9	g/dL
10	A/G ratio Serum, Method: Calculated	1.92	1.0 - 2.0	
11	Gamma GT Serum, Method: G glutamyl carboxy nitroanilide	21.0	5 - 55	U/L



CRM No:2873284

Sample Recd. Time: 12-07-2021 11:49 Report Time: 12-07-2021 15:55

Patient Name: Mrs. Tannu Hodekar

Patient ID: 2873284



Authorized Signatory Dr. Dina Abhani DCP, DNB (Pathology)





PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000







PMF...









Name: Mrs. Tannu Hodekar

Age/Gender:

27 Year(s) 0 Months(s) 0 Day(s)/Female

**Referred By:** SELF

Client Name:

GEO MEDCARE HOSPITAL

**Collection Date:** 12-07-2021 11:40:00

**Report Release Date:** 12-07-2021 15:55:55

Sr.No	Investigation	<b>Observed Value</b>	Reference Range	Unit
Lipid P	rofile			
1	Total Cholesterol Serum, Method: Photometry	184.0	Desirable: <200 Borderline high = 200-239 High: > 240	mg/dl
2	Triglycerides Serum, Method: Enzymatic, end point coupled assay	77.5	Desirable: <150 Borderline High: 150 - 199 High: > 200 - 499	mg/dl
3	HDL-Cholesterol Serum, Method: Photometry	53.3	30 - 60	mg/dL
4	LDL- Cholesterol Serum, Method: Photometry	115.20	Optimal: <100; Near Optimal: 100-129; Borderline High: 130-159; High: 160-189; Very high: >190	mg/dl
5	Cholesterol/HDL ratio Serum, Method: Calculated	3.45	Optimal: <3.5 Near Optimal: 3.5 - 5.0 High >5.0	
6	VLDL Cholesterol Serum, Method: Calculated	15.50	6 - 40	mg/dL
7	Non HDL Cholesterol Serum, Method: Calculated	130.70	Desirable: <130 Borderline high: 130-159 High: 160-189 Very High:>190	mg/dl
8	LDL /HDL ratio Serum, Method: Calculated	2.16	Optimal: <2.5 Near Optimal: 2.5-3.5 High >3.5	

## **Interpretation**

- 1.Triglycerides: When triglycerides are very high greater than 1000 mg/dL, there is a risk of developing pancreatitis in children and adults. Triglycerides change dramatically in response to meals, increasing as much as 5 to 10 times higher than fasting levels just a few hours after eating. Even fasting levels vary considerably day to day. Therefore, modest changes in fasting triglycerides measured on different days are not considered to be abnormal.
- 2. HDL-Cholesterol: HDL- C is considered to be beneficial, the so-called "good" cholesterol, because it removes excess cholesterol from tissues and carries it to the liver for disposal. If HDL-C is less than 40 mg/dL for men and less than 50 mg/dL for women, there is an increased risk of heart disease that is independent of other risk factors, including the LDL-C level. The NCEP guidelines suggest that an HDL cholesterol value greater than 60 mg/dL is protective and should be treated as a negative risk factor.
- 3. LDL-Cholesterol: Desired goals for LDL-C levels change based on individual risk factors. For young adults, less than 120 mg/dL is acceptable. Values between 120-159 mg/dL are considered Borderline high. Values greater than 160 mg/dL are considered high. Low levels of LDL cholesterol may be seen in people with an inherited lipoprotein deficiency and in people with hyperthyroidism, infection, inflammation, or cirrhosis.



CRM No: 2873284

Sample Recd. Time: 12-07-2021 11:49 Report Time: 12-07-2021 15:55

Patient Name: Mrs. Tannu Hodekar

Patient ID: 2873284



Authorized Signatory Dr. Dina Abhani DCP, DNB (Pathology)





PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000







PMF...









Name: Mrs. Tannu Hodekar

**SELF** 

Age/Gender:

27 Year(s) 0 Months(s) 0

Day(s)/Female

Referred By:

Client Name:

GEO MEDCARE HOSPITAL

**Collection Date:** 12-07-2021 11:40:00

**Report Release Date:** 12-07-2021 15:55:55

Sr.No	Investigation	<b>Observed Value</b>	Reference Range	Unit
Thyroid	l Profile - Total T3,Total T4,TSH (TFT	)		
1	Total T3 Serum, Method: CLIA	90.56	60 - 200	ng/dL
2	Total T4 Serum, Method: CLIA	10.44	4.5 - 14.5	μg/dL
3	TSH (Thyroid Stimulating Hormone) Serum, Method: CLIA	1.548	0.35 - 5.5	μIU/ml

## **Interpretation**

- 1. Triodothyronine (T3) is produced by the thyroid gland and along with thyroxine (T4) help control the rate at which the body uses energy. Elevated T3 denote hyperthyroidism while low levels indicate hypothyroidism.
- 2.The most common causes of thyroid dysfunction are related to autoimmune disorders. Graves disease causes hyperthyroidism, but it can also be caused by thyroiditis, thyroid cancer, and excessive production of TSH. Total T3 is used to assess thyroid function.
- 3. Elevated T4 levels may indicate hyperthyroidism. They may also indicate other thyroid problems, such as thyroiditis or toxic multinodular goiter. Abnormally low levels of T4 may indicate: dietary issues, such as fasting, malnutrition, or an iodine deficiency, medications that affect protein levels, hypothyroidism, illness.
- 4. Thyroid-stimulating hormone (TSH) stimulates the production and release of T4 (primarily) and T3. They help control the rate at which the body uses energy and are regulated by a feedback system. Most of the T4 circulates in the blood bound to protein, while a small percentage is free (not bound).
- 5. Lab has estimated Total T4 reference intervals that are specific for India, using the indirect sampling technique following CLSI EP28-A3c document: Defining Establishing, and Verifying Reference Intervals in the Clinical Laboratory: Approved Guideline-Third Edition.
- 5. Thyroid hormone status during pregnancy:



CRM No :2873284

Sample Recd. Time: 12-07-2021 11:49

Report Time: 12-07-2021 15:55 Patient Name: Mrs. Tannu Hodekar

Patient ID: 2873284

Dans

Authorized Signatory
Dr. Dina Abhani
DCP, DNB (Pathology)

Authorized Sign



PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000







PMF









Name: Mrs. Tannu Hodekar

Age/Gender:

27 Year(s) 0 Months(s) 0

Day(s)/Female

Referred By: SELF Client Name:

GEO MEDCARE HOSPITAL

**Collection Date:** 12-07-2021 11:40:00 **Report Release Date:** 12-07-2021 15:55:55

Sr.No	Investigation	Observed Value	Reference Range	Unit
GD We	llness Kidney Profile			
1	BUN (Blood Urea Nitrogen) Serum, Method: Calculated	8.93	3.3 - 18.7	mg/dL
2	Creatinine Serum, Method: Alkaline picrate kinetic	0.6	0.4 - 1.2	mg/dL
3	BUN/Creatinine ratio Serum, Method: Calculated	14.88	5.0 - 23.5	
4	Uric Acid Serum, Method: Uricase, UV	4.2	2.0 - 6.5	mg/ dL
5	Calcium Serum, Method: O cresolphthalein complexone	9.4	8.5 - 10.5	mg/dL
6	eGFR (estimated Glomerular Filtration Rate) Serum, Method: Calculated	127.23	Normal: > 90 Mild decrease in GFR: 60 -89 Moderate decrease in GFR: 30-59 Severe decrease in GFR: 15- 29 Kidney failure: < 15	mL/min/1.73 m²

# Interpretation

A renal function panel could be ordered when a patient has risk factors for kidney dysfunction such as high blood pressure (hypertension), diabetes, cardiovascular disease, obesity, elevated cholesterol, or a family history of kidney disease. A renal function panel may also be ordered when someone has signs and symptoms of kidney disease, though early kidney disease often does not cause any noticeable symptoms. It may be initially detected through routine blood or urine testing. Renal function panel results are not diagnostic but rather indicate that there may be a problem with the kidneys and that further testing is required to make a diagnosis and determine the cause. Results of the panel are usually considered together, rather than separately. Individual test result can be abnormal due to causes other than kidney disease, but taken together with risks and signs and symptoms, they may give an indication of whether kidney disease is present.



CRM No :2873284

Sample Recd. Time: 12-07-2021 11:49 Report Time: 12-07-2021 15:55

Patient Name: Mrs. Tannu Hodekar Patient ID: 2873284 Dains

Authorized Signatory
Dr. Dina Abhani
DCP, DNB (Pathology)





PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000





PMF









Name: Mrs. Tannu Hodekar

Age/Gender:

27 Year(s) 0 Months(s) 0 Day(s)/Female

Referred By:

SELF

Client Name:

GEO MEDCARE HOSPITAL

**Collection Date:** 

12-07-2021 11:40:00

**Report Release Date:** 12-07-2021 15:55:55

Sr.No	Investigation	Observed Value	Reference Range	Unit
Iron Stu	ıdies (Iron,TIBC, Transferrin saturatio	n)		
1	Iron Serum, Method: Ferene	102.34	50 - 170	μg/dL
2	TIBC Serum, Method: Ferene	300.55	250-450	μg/dL
3	Transferrin saturation Serum, Method: Calculated	34.05	20 - 50	%

## Interpretation

- 1. Serum iron measures the level of iron in the liquid portion of the blood. Low iron levels may seen in anemia (microcytic and hypochromic) . High levels of serum iron in hereditary hemochromatosis, multiple blood transfusions, and a few other conditions.
- 2. TIBC (Total iron-binding capacity) measures all the proteins in blood available to bind with iron, including transferrin.TIBC test is a good indirect measurement of transferrin. The body produces transferrin in relationship to the need for iron. When iron stores are low, transferrin levels increase and vice versa. Since transferrin is the primary iron-binding protein, the TIBC test is a good indirect measurement of transferrin availability.



CRM No:2873284

Sample Recd. Time: 12-07-2021 11:49

Report Time: 12-07-2021 15:55 Patient Name: Mrs. Tannu Hodekar

Patient ID: 2873284



Authorized Signatory
Dr. Dina Abhani
DCP, DNB (Pathology)





PreventiNe Life Care (P) Ltd. Plot-6, Sector - 24, Navi Mumbai Maharashtra, India. Pin - 400705 Tel: +91-22-40450000

Referred By:







PMF...









Name: Mrs. Tannu Hodekar

Age/Gender:

27 Year(s) 0 Months(s) 0 Day(s)/Female

Client Name:

GEO MEDCARE HOSPITAL

**Collection Date:** 12-07-2021 11:40:00

**SELF** 

**Report Release Date:** 12-07-2021 15:55:55

Sr.No	Investigation	Observed Value	Reference Range	Unit
HbA1c	(Whole Blood)			
1	HBA1c-Glycated Haemoglobin EDTA Whole Blood, Method: HPLC	5.4	Non-diabetic: 4-6 Excellent Control: 6-7 Fair to good control: 7-8 Unsatisfactory control: 8-10 Poor Control: >10	%
2	Estimated Average Glucose (eAG) EDTA Whole Blood, Method: Calculated	108.28	90-120 mg/dL : Good control 121-150 mg/dL : Fair control 151-180 mg/dL : Unsatisfactory control >180 mg/dL : Poor control	mg/dL

## **Interpretation**

- 1.The term HbA1c refers to Glycated Haemoglobin. Measuring HbA1c gives an overall picture of what the average blood sugar levels have been over a period of weeks/month. Higher the HbA1c, the greater the risk of developing diabetes-related complications.
- 2.HbA1c has been endorsed by clinical groups and ADA (American Diabetes Assocation) guidelines 2012, for the diagnosis of diabetes using a cut-off point of 6.5%. ADA defined biological reference range for HbA1c is between 4-6%. Patients with HBA1c value between 6.0-6.5% are considered at risk for developing diabetes in the future. Trends in HbA1c area a better indicator of glucose control than standalone test.
- 3.To estimate the eAG from the HbA1c value, the following equation is used: eAG(mg/dl) = 28.7\*A1c-46.7.
- 4.Diabetic must aspire to keep values under 7% to avoid the various complications resulting from diabetes.

## **End Of Report**



CRM No: 2873284

Sample Recd. Time: 12-07-2021 11:49

Report Time: 12-07-2021 15:55 Patient Name: Mrs. Tannu Hodekar

Patient ID: 2873284



Authorized Signatory
Dr. Dina Abhani
DCP, DNB (Pathology)

Destronde

