115 3 Sanghar Bungalow Lane No 14 Prabhat Road

Pune

REPORT

Tel No: 919823316416

PID: 121895

Reference:Dr.--SID: 120165451

> Collection Date: 31-12-2020 11:17 AM Registration Date: 31-12-2020 11:17 am Report Date: 31-12-2020 04:19 PM

Age:42.70 Years Sex: MALE

Complete Blood Count (EDTA Whole Blood)	Result	Biological Reference Interval
Hemoglobin (Hb), EDTA whole blood	14.80	14.0 - 17.50 g/dL
Method: Photometry	14100	1110 17100 g/ d_
Total Leucocytes (WBC) count	5,300	4000-10000/µL
Method : Coulter Principle / Microscopy	0,000	1000 10000/µ2
Platelet count	258,000	150000 - 450000 /µL
Method : Coulter Principle / Microscopy	200,000	100000 100000 /μ2
Red blood cell (RBC) count	4.99	4.52 - 5.90 x 10^6 /μL
Method: Coulter Principle	4.00	1.02 0.00 χ 10 0 /μ2
PCV (Packed Cell Volume)	43.20	41.5 - 50.4 %
Method: Calculated	10.20	
MCV (Mean Corpuscular Volume)	86.50	80.0 - 96.0 fL
Method: Derived from RBC histogram	00.00	33.3 33.3
MCH (Mean Corpuscular Hb)	29.60	27.5 - 33.2 pgms
Method: Calculated	20.00	c
MCHC (Mean Corpuscular Hb Conc.)	34.20	33.4 - 35.5 g/dL
Method: Calculated	020	30.1. 30.0 g/ u_
RDW (RBC distribution width)	13.30	11.6 - 14.6 %
Method: Derived from RBC Histogram		
WBC Differential Count		
Method: VCSn / Microscopy / Calculated		
Neutrophils	57	40 - 80 %
Absolute Neutrophils	3,021	2000 - 7000 /μL
	-,-	•
Eosinophils	3	1 - 6 %
Absolute Eosinophils	159	20 - 500 /μL
Basophils	0	0 - 2 %
Absolute Basophils	0	0 - 100 /µL
	-	
Lymphocytes	34	20 - 40 %
Absolute Lymphocytes	1,802	1000 - 3000 /µL
	-,	
Monocytes	6	2 - 10 %
Absolute Monocytes	318	200 - 1000 /μL
-	@	
-		



Page 1 of 8

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.

rehendale





115 3 Sanghar Bungalow Lane No 14 Prabhat Road

Pune

Tel No: 919823316416

PID: 121895

Age:42.70 Years Sex:MALE

Reference:Dr.-- SID: 120165451

Collection Date: 31-12-2020 11:17 AM Registration Date: 31-12-2020 11:17 am Report Date:

31-12-2020 04:19 PM

Complete Blood Count Findings

R.B.C. : Normocytic, Normochromic

W.B.C. : No abnormality detected

Platelets : Adequate

Remark : ON FOLLOW UP.

-

-

REPORT

•

.



Page 2 of 8

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.

nehendale

115 3 Sanghar Bungalow Lane No 14 Prabhat Road

Pune

Tel No: 919823316416

PID: 121895

Age:42.70 Years Sex:MALE

Reference:Dr.-- SID: 120165451

Collection Date: 31-12-2020 11:17 AM Registration Date: 31-12-2020 11:17 am Report Date:

31-12-2020 04:19 PM

Test Description Observed Value Biological Reference Interval

TEST NAME

REPORT

Glycated Hemoglobin (HbA1C), by HPLC 5.30 4.0 to 5.6 %

Interpretation:

HbA1C level reflects the mean glucose concentration over previous 8-12 weeks and provides better indication of long term glycemic control.

For diagnosis of Diabetes Mellitus (>/= 18 yrs of age) :

5.7 % - 6.4 %: Increased risk for developing diabetes.

>/= 6.5 % : Diabetes

Therapeutic goals for glycemic control:

Adults: < 7%

Toddlers and Preschoolers: < 8.5% (but > 7.5%)

School age (6-12 yrs): < 8%

Adolescents and young adults (13 - 19 yrs): < 7.5 %

Levels of HbA1C may be low as result of shortened RBC life span in case of hemolytic anemia. Increased HbA1C values may be found in patients with polycythemia or post splenectomy patients. Patients with Homozygous forms of rare variant Hb(CC,SS,EE,SC) HbA1c can not be quantitated as there is no HbA. In such circumstances glycemic control can be monitored using plasma glucose levels or serum Fructosamine.

The A1c target should be individualized based on numerous factors, such as age, life expectancy, comorbid conditions, duration of diabetes, risk of hypoglycemia or adverse consequences from hypoglycemia, patient motivation and adherence.

Ref: ADA (Standards of Medical Care in Diabetes - 2017)



Page 3 of 8

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.

rehendale







JITAL RAMESH SHAH 115 3 Sanghar Bungalow

Lane No 14 Prabhat Road

Pune

Tel No: 919823316416

PID: 121895

Age:42.70 Years Sex: MALE

Reference: Dr.--

SID: 120165451

Collection Date: 31-12-2020 11:17 AM Registration Date: 31-12-2020 11:17 am Report Date:

31-12-2020 04:19 PM

Test Description Observed Value Biological Reference Interval

Haematology:

Erythrocyte Sedimentation Rate, EDTA Whole Blood 09

•

Male under 50 Yrs: Upto 15mm/hr. Male 50 - 85 Yrs: Upto 20mm/hr. Male > 85 yrs: Upto 30mm/hr. Results corrected to 18 deg. celsius

Technique : Automated Westergren Method .

- 1. ESR is markedly elevated in monodonal gammopathy such as multiple myeloma, in severe polyclonal hyperglobulinemia due to inflammatory disease, and in hyperfibrinogenemia. 2. Moderate elevations are common in active inflammatory disease such as rheumatoid arthritis, chronic infections, collagen disease and neoplastic disease
- 3. ESR has little diagnostic value in these disorders but can be useful in monitoring disease activity.
- 4. Useful in the diagnosis and in monitoring polymyalgia rheumatica and temporal arteritis.
- 5. Moderate increase is seen in pregnancy (beginning at the 10th to 12th week) and returns to normal about 1 month postpartum.
- 6. Red cells with an abnormal or irregular shape, such as sickle cells or spherocytes, hinder rouleaux formation and lower the ESR.



Page 4 of 8

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.

rehendale







JITAL RAMESH SHAH 115 3 Sanghar Bungalow

Lane No 14 Prabhat Road

Pune

Tel No: 919823316416

PID: 121895

Age:42.70 Years Sex:MALE

Reference: Dr.--

SID: 120165451

Collection Date: 31-12-2020 11:17 AM Registration Date: 31-12-2020 11:17 am Report Date: 31-12-2020 04:19 PM

Test Description	Observed Value	Biological Reference Interval
Plasma Glucose :		
Plasma glucose, random by Hexokinase method	97	< 200 mg/dL American Diabetes Association
		Guidelines 2020

Hormones

1. Levels are increased in insulinomas, factitious hypoglycemia, insulin autoimmune syndrome, acromegaly (after ingestion of glucose), Cushings syndrome, corticosteroid administration and levodopa usage. 2. Levels are depressed to absent in diabetes mellitus, pituitary tumors and chronic pancreatic diseases i.e. cystic fibrosis.
3. Insulin/C-peptide ratio is used for differentiating between factitious hypoglycemia and insulinomas where a ratio < 1.0 indicates insulinoma; but results may vary in renal failure. 4. Antibodies to insulin form is longstanding diabetes mellitus treated with insulin hence in these



Page 5 of 8

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.

rehendale







JITAL RAMESH SHAH 115 3 Sanghar Bungalow Lane No 14 Prabhat Road

Pune

Tel No: 919823316416

PID: 121895

Age:42.70 Years Sex:MALE

Reference:Dr.--

Collection Date: 31-12-2020 11:17 AM Registration Date: 31-12-2020 11:17 am Report Date:

SID: 120165451

31-12-2020 04:19 PM

Test Description Hormones:

Observed Value

Biological Reference Interval



Page 6 of 8

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.

nehendale

Dr. Vinanti Golwilkar MD (Pathology)

ilkar

MD (Pathology)

115 3 Sanghar Bungalow Lane No 14 Prabhat Road

Pune

REPORT Tel No: 919823316416

PID: 121895

Reference: Dr.--

SID: 120165451

Collection Date: 31-12-2020 11:17 AM Registration Date: 31-12-2020 11:17 am Report Date: 31-12-2020 04:19 PM

Age:42.70 Years Sex:MALE

CRP(hs) - C- Reactive Protein high sensitivity

Test Description Observed Value

2.05 See clinical information below

Method: Nephelometry / Immunoturbidimetry

Biological Reference Interval

Clinical Information:

- 1. C-reactive protein (CRP) is a biomarker of inflammation. Plasma CRP concentrations increase rapidly and dramatically (100-fold or more) in response to tissue injury or inflammation.
- 2. High-sensitivity CRP (hs-CRP) is more precise than standard CRP when measuring baseline (i.e. normal) concentrations and enables a measure of chronic inflammation. It is recommended for cardiovascular risk assessment. Atherosclerosis is an inflammatory disease and hs-CRP has been endorsed by multiple guidelines as a biomarker of atherosclerotic cardiovascular disease risk.

Low cardiovascular risk : < 2.0 mg/L High cardiovascular risk : >/= 2.0 mg/L Acute inflammation : > 10.0 mg/L

3. A single test for high-sensitivity CRP (hs-CRP) may not reflect an individual patient's basal hs-CRP level. Repeat measurement may be required to firmly establish an individual's basal hs-CRP concentration. The lowest of the measurements should be used as the predictive value.

Reference: Mayo Medical Laboratories

Inchendale

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.

Page 7 of 8







115 3 Sanghar Bungalow Lane No 14 Prabhat Road

Age:42.70 Years Sex: MALE

Pune

PID: 121895

Tel No: 919823316416

Observed Value Negative (0.04)

Reference: Dr.--

31-12-2020 04:19 PM **Biological Reference Interval**

Negative : < 1.4 Index (S/C) Positive : $>/= 1.4 \ln (S/C)$

SID: 120165451

31-12-2020 11:17 AM

31-12-2020 11:17 am

Registration Date:

Collection Date:

Report Date:

Test Description

SARS-CoV-2 IgG Antibodies, Serum by CMIA

Remarks:

REPORT

- * IgG test is not useful for diagnosis of acute infection.
- * IgG antibodies usually appear after 2 weeks (14 days) of infection. Presence of IgG antibodies may / may not indicate immunity.
- * Detection of IgG antibodies may be useful for :
- a. Understanding whether an individual is exposed to infection with SARS-CoV-2 including asymptomatic individuals.
- b. Understanding the seroprevalence in communities and especially high risk or vulnerable populations.

Reference: ICMR Advisory dated 23/06/2020

End of Report

Page 8 of 8

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv

rehendale

Dr.(Mrs.) Awanti Golwilkar Mehendale MD(Path) Regn.No.: 2000/02/1052 A.G Diagnostics Pvt. Ltd.



