Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

# **REPORT**

Age:37.80 Years Sex:MALE

Complete Blood Count	Result	Biological Reference Interval
(EDTA Whole Blood)		
Hemoglobin (Hb), EDTA whole blood	<u>13.90</u>	14.0 - 17.50 g/dL
Method: Photometry		
Total Leucocytes (WBC) count	6,000	4000-10000/μL
Method : Coulter Principle / Microscopy		
Platelet count	389,000	150000 - 450000 /μL
Method : Coulter Principle / Microscopy		
Red blood cell (RBC) count	4.93	4.52 - 5.90 x 10^6 /µL
Method: Coulter Principle		
PCV (Packed Cell Volume)	42.20	41.5 - 50.4 %
Method: Calculated		
MCV (Mean Corpuscular Volume)	85.70	80.0 - 96.0 fL
Method: Derived from RBC histogram		
MCH (Mean Corpuscular Hb)	28.30	27.5 - 33.2 pgms
Method: Calculated		
MCHC (Mean Corpuscular Hb Conc.)	<u>33.00</u>	33.4 - 35.5 g/dL
Method: Calculated		
RDW (RBC distribution width)	13.30	11.6 - 14.6 %
Method: Derived from RBC Histogram		
WBC Differential Count		
Method: VCSn / Microscopy / Calculated		
Neutrophils	50	40 - 80 %
Absolute Neutrophils	3,000	2000 - 7000 /μL
Essinantilo	06	4 6 9/
Eosinophils	06	1 - 6 %
Absolute Eosinophils	360	20 - 500 /μL
Basophils	00	0 - 2 %
Absolute Basophils	0	0 - 100 /µL
Abbolato Bacopillo	· ·	ο 100/μΕ
Lymphocytes	36	20 - 40 %
Absolute Lymphocytes	2,160	1000 - 3000 /μL
Monocytes	08	2 - 10 %
Absolute Monocytes	480	200 - 1000 /μL
-	<>	

Reference:Dr.--



rehendale







Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28

**REPORT** 

Tel No: 918879699138 PID: 11531066

Age:37.80 Years Sex:MALE

Reference:Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

## **Complete Blood Count Findings**

R.B.C. Normocytic, Normochromic

W.B.C. No abnormality detected

**Platelets Adequate** 

Remark



rehendale

Page 2 of 15 Pr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ w. \_\_\_\_\_A.G Diagnostics Pvt. Ltd.





Flat No-M-301, Marvel Diva,

Magarpatta Road, Near Seasons Mall,

**REPORT** 

Hadapsar, Pune-28 Tel No: 918879699138

PID: 11531066

Age:37.80 Years Sex:MALE

Reference:Dr.--SID: 120196846

> Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date:

10-02-2021 08:04 PM

Test Desciption	Observed Value	Biological Reference Interval
<u>Lipid Profile Maxi :</u>		
Serum Appearance	Clear	
Cholesterol (Total), serum by Enzymatic method	117	Desirable : < 200 mg/dL Borderline high : 200 - 239 mg/dL High : >/= 240 mg/dL
Triglycerides, serum by Enzymatic method	97	Normal: < 150 mg/dL Borderline high: 150-199 mg/dL High: 200-499 mg/dL Very high: >/= 500 mg/dL
HDL Cholesterol, serum by Enzymatic method	<u>35</u>	Men : > 40 mg/dL Women : > 50 mg/dL
VLDL Cholestrol, serum by calculation	19	< 30 mg/dL
LDL Cholesterol, serum by calculation	63	Optimal: <100 mg/dL Near optimal/above optimal: 100-129 mg/dL Borderline high: 130-159 mg/dL High: 160-189 mg/dL Very high: >/= 190 mg/dL
Cholesterol(Total)/HDL Cholesterol Ratio	3.34	Males : Acceptable ratio = 5.00<br Females : Acceptable ratio = 4.50</td
LDL Cholesterol/HDL Cholesterol Ratio	1.79	Males : Acceptable ratio = 3.60<br Females : Acceptable ratio = 3.20</td
Apolipoprotein A1, serum by Nephelometry	127	Male : 110 to 205 mg/dL
Apolipoprotein B, serum by Nephelometry	58	55 to 140 mg/dL

## Reference: ATP III, NCEP Guidelines and National Lipid Association (NLA) 2014 Recommendations

As per most international and national guidelines including Lipid Association of India 2016:

- 1. Lipoprotein and lipid levels should be considered in conjunction with other atherosclerotic cardiovascular disease (ASCVD) risk determinants to assess treatment goals and strategies.
- 2. Non-fasting lipid levels can be used in screening and in general risk estimation.



rehendale

Page 3 of 15 Page





**REPORT** 

Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

**AASHIESH GUPTA** 

Reference:Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

Age:37.80 Years Sex:MALE

Test Description Liver Function Test :	Observed	Biological Reference Interval
Bilirubin-Total, serum by Diazo method	0.70	0.10 - 1.20 mg/dL Neonates : Upto 15.0 mg/dL
Bilirubin-Conjugated, serum by Diazo method	0.32	Upto 0.5 mg/dL
Bilirubin-Unconjugated, serum by calculation	0.38	0.1 to 1.0 mg/dL
SGOT (AST), serum by Enzymatic method	31	>or= 14 years : 8 - 48 U/Lt
SGPT (ALT), serum by Enzymatic Method	41	7 to 55 U/Lt
Alkaline Phosphatase,serum by pNPP-kinetic	59	Adult Male: (Unit: U/Lt.) 15 - < 17 years: 82 - 331 17 - < 19 years: 55 - 149 > or = 19 years: 40 - 129
Protein (total), serum by Biuret method	6.50	6.4 to 8.2 g/dL
Albumin, serum by Bromocresol purple method	4.05	3.4 to 5.0 g/dL
Globulin, serum by calculation	2.45	2.3 - 3.5 g/dL



--XX--

rehendale

Page 4 of 15 Pr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ w. \_\_\_\_\_A.G Diagnostics Pvt. Ltd.





Flat No-M-301, Marvel Diva,

Magarpatta Road, Near Seasons Mall,

REPORT

Hadapsar, Pune-28 Tel No: 918879699138

PID: 11531066

Age:37.80 Years Sex:MALE

Reference: Dr.--

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date:

SID: 120196846

10-02-2021 08:04 PM

**Test Description Observed Value Biological Reference Interval** 

**TEST NAME** 

Glycated Hemoglobin (HbA1C), by HPLC 4.0 to 5.6 % 6.80

## Interpretation:

HbA1C level reflects the mean glucose concentration over previous 8-12 weeks and provides better indication of long term glycemic control.

## For diagnosis of Diabetes Mellitus (>/= 18 yrs of age) :

5.7 % - 6.4 %: Increased risk for developing diabetes.

>/= 6.5 % : Diabetes

## Therapeutic goals for glycemic control:

Adults: < 7%

Toddlers and Preschoolers: < 8.5% (but > 7.5%)

School age (6-12 yrs): < 8%

Adolescents and young adults (13 - 19 yrs): < 7.5 %

Levels of HbA1C may be low as result of shortened RBC life span in case of hemolytic anemia. Increased HbA1C values may be found in patients with polycythemia or post splenectomy patients. Patients with Homozygous forms of rare variant Hb(CC,SS,EE,SC) HbA1c can not be quantitated as there is no HbA. In such circumstances glycemic control can be monitored using plasma glucose levels or serum Fructosamine.

The A1c target should be individualized based on numerous factors, such as age, life expectancy, comorbid conditions, duration of diabetes, risk of hypoglycemia or adverse consequences from hypoglycemia, patient motivation and adherence.

Ref: ADA (Standards of Medical Care in Diabetes - 2017)



Page 5 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd.

ehendale

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv. .....



Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28

REPORT Tel No

Tel No: 918879699138 PID: 11531066

Age:37.80 Years Sex:MALE

Reference: Dr.--

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date:

SID: 120196846

10-02-2021 08:04 PM

Test Description Observed Value Biological Reference Interval

Gamma Glutamyl Transferase (GGT)

Gamma GT(GGT), Serum by Carboxy substrate-kinetic 44.00 Male: (Unit: U/Lt.)

13 - 17 years : < 43 >or= 18 years : 8 - 61

## Interpretation

- \* GGT is used to diagnose and monitor hepatobiliary diseases.
- \* Increased GGT and Alkaline Phosphatase indicate hepatobiliary diseases.
- \* Normal GGT activity and increased Alkaline Phosphatase is consistent with skeletal disease.
- \* May be used a screening test for occult alcoholism.
- \* Elevated GGT is seen in:
  - 1) Intra or post hepatic biliary obstruction (5 to 30 times normal)
  - 2) Infectious hepatitis (2 to 5 times normal)
  - 3) Alcoholism
  - 4) Sclerosing cholangitis
  - 5) Primary or secondary neoplasm
  - 6) Medications such as phenytoin and phenobarbitone

Reference: Mayo Medical Laboratories, 2018 Interpretive Handbook.

--<>--



Page 6 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 M. A.G Diagnostics Pvt. Ltd.

ehendale

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ w. ......





Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

REPORT

Age:37.80 Years Sex:MALE

Reference: Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

Test Description	Observed Value	Biological Reference Interval
DI Al		

Plasma Glucose : Plasma glucose fasting, by Hexokinase method 113

< 100 mg/dL

100 to 125 mg/dL: Impaired fasting glucose tolerance / Prediabetes >/= 126 mg/dL : Suggestive of diabetes mellitus

(On more than one occasion) American Diabetes Association

Guidelines 2020

Plasma glucose post prandial, by Hexokinase method 199 < 140 mg/dL

> 140 to 199 mg/dL: Impaired glucose tolerance / Prediabetes >/= 200 mg/dL : Suggestive of

diabetes mellitus

(On more than one occasion) American Diabetes Association


Guidelines 2020

## **Clinical Chemistry**

17 to 49 mg/dL Urea, serum by GLDH-urease 20 BUN-Blood Urea Nitrogen, serum by calculation 9.35 8 to 23 mg/dL Creatinine, serum by Jaffe w/o deproteinization 0.96 0.6 to 1.2 mg/dL



rehendale







Flat No-M-301, Marvel Diva,

Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

Age:37.80 Years Sex: MALE

Reference:Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

Test Description Observed Value Biological Reference Interval

**Clinical Chemistry:** 

REPORT

Uric Acid, serum by Uricase method 4.70 Male: 3.50 to 7.20 mg/dL

- \* Uric acid is useful for 1. Diagnosis and follow up of renal failure. 2. Monitoring patients receiving cytotoxic drugs and a variety of other disorders, including gout, leukemia, psoriasis, starvation and other wasting conditions . \* Increased uric acid is seen in following conditions:
- 1. Increased purine synthesis 2. Inherited metabolic disorders 3. Excess dietary purine intake
- 4. Increased nucleic acid turnover 5. Malignancy, cytotoxic drugs 6. Decreased urinary excretion (due to CRF) 7. Increased renal reabsorption .
- \* Uric acid is decreased in : 1. Hepatocellular disease with reduced purine synthesis
- 2. Defective renal reabsorption 3. Overtreatment of uricemia (allopurinol or cancer therpies like 6-mercaptopurine, etc).



Page 8 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd.

rehendale

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv. ......

Dr. Awanti Golwilkar
MD (Pathology)

Dr. Vinanti Golwilkar
MD (Pathology)

Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

REPORT

Age:37.80 Years Sex: MALE

Reference: Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

**Test Description Observed Value Biological Reference Interval Clinical Chemistry:** 

Calcium, serum by OCPC method 9.40 Adult: 8.4 to 10.2 mg/dL

Method: Colorimetric (o-cresolpthalein substrate).

1. Calcium is useful for diagnosis and monitoring of a wide range of disorders including diseases of bone, kidney, parathyroid gland, or gastrointestinal tract.

2. Calcium ions play an important role in blood clotting, bone mineralization, musculature contractility and CNS functioning. .

3. Hypocalcemia is due to the absence or impaired function of the parathyroid glands or impaired vitamin-D synthesis. Chronic renal failure is also frequently associated with hypocalcemia due to decreased vitamin-D synthesis as well as hyperphosphatemia and skeletal resistance to the action of parathyroid hormone (PTH). 4. Hypercalcemia is mainly due to primary hyperparathyroidism (pHPT), and bone metastasis of carcinoma of the breast, thyroid gland, or lung. Severe hypercalcemia may result in cardiac arrhythmia.



Page 9 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd.

rehendale

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv. .....





Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

**REPORT** 

Age:37.80 Years Sex:MALE

Reference: Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

Test Description Clinical Chemistry:	Observed Value	Biological Reference Interval
<u>Hormones</u>		
Free T3, serum by CMIA	2.04	1.71 to 3.71 pg/mL
Free T4, serum by CMIA	0.96	0.71 to 1.85 ng/dL
TSH(Ultrasensitive), serum by CMIA	0.82	0.40 - 4.00 μIU/mL

MC-3143

Page 10 of 15

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv

DIAGNOSTICS
BE SURE
BE WELL
ए.जी डायग्जॉस्टिक्स प्रा. लि.\_\_\_\_\_\_\_A.G Diagnostics Pvt. Ltd.

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd.

pehendale

Dr. Vinanti Golwilkar

MD (Pathology)

**Carrying forward** 

Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138

**REPORT** PID: 11531066

Age:37.80 Years Sex: MALE

Reference:Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

**Test Description Observed Value Biological Reference Interval** 

**TEST NAME** 

Vitamin B12, serum by CMIA 195.0 187 - 883 pg/mL

## Interpretation:

- 1. Vitamin B12 (cobalamin) is necessary for hematopoiesis and normal neuronal function.
- 2. Vitamin B12 is decreased in

Decreased Serum B12	
Pregnancy	
Contraceptive hormones	
Malabsorption	
Ethanol ingestion	
Smoking	
Strict vegan diet	
Pernicious anemia	

- 3. Serum methylmalonic acid and homocysteine levels are also elevated in vitamin B12 deficiency states. Active B12 (Holotranscobalamin) is low in Vitamin B12 deficiency.
- 4. Please correlate in case of patients taking vitamin B12 supplementation.



Page 11 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd. "Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv. .....

ehendale



Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

Age:37.80 Years Sex:MALE

Reference: Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

Test Description Observed Value Reference range & Units

**TEST NAME** 

REPORT

Homocysteine, plasma by CMIA 13.13 Male: 5.08 to 15.39 µmol/Lt

Homocysteine concentration is an indicator of acquired folate or cobalamin deficiency, and is a contributing factor in the pathogenesis of neural tube defects. Currently, the use of homocysteine for assessment of cardiovascular risk is uncertain and controversial. Based on several meta-analyses, at present, homocysteine may be regarded as a weak risk factor for coronary heart disease, and there is a lack of direct causal relationship between hyperhomocysteinemia and cardiovascular disease. It is most likely an indicator of poor lifestyle and diet. Homocysteine concentrations >13 mcmol/L are considered abnormal in patients evaluated for suspected nutritional deficiencies (B12, folate) and inborn errors of metabolism. Homocysteine concentrations < or =10 mcmol/L are desirable when utilized for cardiovascular risk.



Page 12 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd.

ehendale

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv......





Magarpatta Road, Near Seasons Mall,

Flat No-M-301, Marvel Diva,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

Age:37.80 Years Sex:MALE

**Test Description Observed Value** 

**Biological Reference Interval** 

**TEST NAME** 

REPORT

25 - OH Vitamin D, serum by CMIA

**18.40** Severe deficiency: < 10 ng/mL

Reference: Dr .--

Mild to moderate deficiency: 10 to 19 ng/mL

Optimum levels: 20 to 50 ng/mL

Increased risk of hypercalciuria: 51 to 80

ng/mL

Toxicity possible : > 80 ng/mL Ref.: Mayo Medical Laboratories These reference ranges represent clinical decision values, based on the 2011 Institute of Medicine report

## Interpretation:

Vitamin D is vital for strong bones. It also has important, emerging roles in immune function and cancer prevention.

Vitamin D compounds in the body are exogenously derived by dietary means; from plants as 25-hydroxyvitamin D2 (ergocalciferol or calciferol) or from animal products as 25-hydroxyvitamin D3 (cholecalciferol or calcidiol).

Vitamin D may also be endogenously derived by conversion of 7-dihydrocholesterol to 25-hydroxyvitamin D3 in the skin upon ultraviolet exposure.

The total 25-hydroxyvitamin D (25-OH-VitD) level (the sum of 25-OH-vitamin D2 and 25-OH-vitamin D3) is the appropriate indicator of vitamin D body stores.

Patients with renal failure can have very high 25-OH-VitD levels without any signs of toxicity, as renal conversion to the active hormone 1,25-OH-VitD is impaired or absent.

Kindly corelate clinically, with supplementation history & repeat with fresh sample if necessary.



Page 13 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd.

ehendale

"Laboratory is accredited as per ISO 15189:2012, Certificate Number MC-3143. Scope available on request / @ wv. .....





Flat No-M-301, Marvel Diva,

Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28

Tel No: 918879699138 PID: 11531066

**Urine Routine Examination** 

REPORT

Reference: Dr.--SID: 120196846

**Biological Reference Interval** 

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

Age:37.80 Years Sex: MALE

(Sample: Urine, Automated / Semiautomated) **Physical Quantity Examined** 5.0 ml Method: Visual Clear **Appearance** Method: Visual / Automated Colour Pale yellow Method: Visual / Automated **Chemical (Dipstick)** pН 6.0 4.6 - 8.0Method: Indicator Principle **Protein Present Trace** Absent Method: Sulphosalycylic Acid/ pH Indicator Glucose **Absent** Absent Method: GOD-POD/Benedict's Acetone **Absent** Absent Method: Sodium Nitroprusside reaction **Absent Bile Pigments** Absent Method: Diazo Reaction / Fouchet's test Urobilinogen Not significant Not Significant Method: Modified Ehrlich / Watson Schwartz Microscopy / Flow cytometry R.B.Cs 1-2 0 - 2 per hpf Pus cells 2-3 0 - 5 per hpf **Epithelial cells** 1-2 0 - 5 per hpf

Result

Casts **Not Detected** 


**Crystals Not Detected** 

@@#

Kindly correlate clinically and follow up.



rehendale







Flat No-M-301, Marvel Diva, Magarpatta Road, Near Seasons Mall,

Hadapsar, Pune-28 Tel No: 918879699138 PID: 11531066

Age:37.80 Years Sex:MALE

Reference:Dr.--

SID: 120196846

Collection Date: 10-02-2021 11:27 AM Registration Date: 10-02-2021 11:27 am Report Date: 10-02-2021 08:04 PM

**Test Description** 

SARS-CoV-2 IgG Antibodies, Serum by CMIA

Observed Value Negative (0.04)

**Biological Reference Interval** Negative: < 1.4 Index (S/C) Positive: >/= 1.4 Index (S/C)

## Remarks:

REPORT

- \* IgG test is not useful for diagnosis of acute infection.
- \* IgG antibodies usually appear after 2 weeks (14 days) of infection. Presence of IgG antibodies may / may not indicate immunity.
- \* Detection of IgG antibodies may be useful for :
- a. Understanding whether an individual is exposed to infection with SARS-CoV-2 including asymptomatic individuals.
- b. Understanding the seroprevalence in communities and especially high risk or vulnerable populations.

Reference: ICMR Advisory dated 23/06/2020

**End of Report** 

Page 15 of 15

Dr.(Mrs.) Awanti Golwilkar Mehendale MBBS,MD(Path) Regn.No:2000/02/1052 A.G Diagnostics Pvt. Ltd.

