#### **NV-Centers in** Diamond as a **Qubit Platform** Alex Heilman

NV-Centers in Diamond as a Qubit Platform An Overview

Alex Heilman

April 18, 2023

### Overview

• What's a qubit?

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond

Hamiltonian Simplifications Energy Levels

IV as Qubit

### Overview

• What's a qubit? What can be a qubit?

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond Hamiltonian Simplifications

NV as Qubit

Gates

4 □ ト 4 圖 ト 4 圖 ト 4 圖 ・ 9 Q (♡)

lefects in liamond lamiltonian limplifications

NV as Qubit

Gates

• What's a qubit? What can be a qubit?

• What's an NV Center?

NV as Qubit

Gates

• What's a qubit? What can be a qubit?

• What's an NV Center?

• How is an NV Center a qubit?

NV as Qubit

Gates

• What's a qubit? What can be a qubit?

• What's an NV Center?

• How is an NV Center a qubit?

### What's a Qubit?

I think we've seen this plenty of times

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in

Hamiltonian Simplification Energy Level

IV as Qubit

Gates

VV as Qubit

Initialization Gates

Gates Measurement

I think we've seen this plenty of times, in case you haven't:

Two-level Quantum System:

$$|0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \qquad |1\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle$$

Normalized: 
$$\alpha^2 + \beta^2 = 1$$

# I think we've seen this plenty of times, in case you haven't:

Two-level Quantum System:

$$|0\rangle = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \qquad |1\rangle = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle$$

Normalized:  $\alpha^2 + \beta^2 = 1$ 

What criteria for physical systems do we need to satisfy?

DiVincenzo's Criteria: The first five are for quantum computers.

- Scalable and discernible
- Fiduciary initial state
- Long decoherence time
- Universal gate set
- Measureable

These next two are for quantum communication

- Memory → Computation
- Faithful transmission

### Criteria for Qubit Platforms cont.

Let's just focus on a few here:

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

#### Qubit Criteria

Defects i Diamond

> Hamiltonian Simplifications Energy Levels

IV as Qubit

### Criteria for Qubit Platforms cont.

Let's just focus on a few here:

Initialization

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

#### Qubit Criteria

Diamond Hamiltonian

Simplifications Energy Levels

V as Qubit

### Criteria for Qubit Platforms cont.

Let's just focus on a few here:

Initialization

Control

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

#### Qubit Criteria

Diamond
Hamiltonian

Energy Levels

V as Qubit

Let's just focus on a few here:

#### Qubit Criteria

Diamond
Hamiltonian
Simplifications

V as Qubit

Gates

Control

Measurement

Initialization

V as Qubit

Gates

Let's just focus on a few here:

• Initialization

Control

Measurement

• Coherence Time

## Why Defects?

Defects can give us localized electronic and spin states trapped in a solid state system.

#### **NV-Centers in** Diamond as a **Qubit Platform**

Alex Heilman

#### Defects in Diamond

V as Qubit

Gates

Measurement

Defects can give us localized electronic and spin states trapped in a solid state system.

Also, gives us more prospective systems, since for every solid state system we'll have (atleast) a few possible defects


Defects can give us localized electronic and spin states trapped in a solid state system.

Also, gives us more prospective systems, since for every solid state system we'll have (atleast) a few possible defects

Really, the concepts discussed below should apply to most similar systems, it's just that this system is well-studied and has convenient energy levels/properties

### Why Diamond?

Diamond is an ideal candidate due to low density of phonon modes (relatively high Debye temperature)



NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond


Simplification

IV as Qubit

itialization

Measuremen

Diamond is an ideal candidate due to low density of phonon modes (relatively high Debye temperature)



This leaves spins of defects and electrons less influenced by phonon modes, increasing their coherence times

### What's an NV-Center

NV-Center refers to a type of defect in diamond lattices with several charge states (-, +, neutral).

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond

Simplifications Energy Levels

NV as Qubit

Gates

Measurement

Simplification Energy Levels


IV as Qubit

Measuremen

NV-Center refers to a type of defect in diamond lattices with several charge states (-, +, neutral). The most commonly discussed charge state is the (-) NV-Center. If unspecified, this is most likely the defect under consideration.

# What's an (-) NV-Center

An (-) NV-Center is a neighboring nitrogen-vacancy defect in diamond with an extra electron



NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond

implifications nergy Levels


IV as Qubit

Initialization Gates

Measurement

# What's an (-) NV-Center

An (-) NV-Center is a neighboring nitrogen-vacancy defect in diamond with an extra electron



This results in a 'center' between the two with localized energy/spin states.

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond


iamiitonian implifications nergy Levels

V as Qubit

Gates Measurement

## What's an (-) NV-Center

An (-) NV-Center is a neighboring nitrogen-vacancy defect in diamond with an extra electron



This results in a 'center' between the two with localized energy/spin states.

These energy states may be used as a Qubit platform.

**NV-Centers** in Diamond as a **Qubit Platform** 

Alex Heilman

Defects in Diamond

We're really interested in the spin state of the electron.

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Derects

Hamiltonian

Simplification Energy Level

IV as Qubit

Gates

Measuremen

We're really interested in the spin state of the electron. Primarily dependent on spin-spin and spin-magnetic field interactions (we'll ignore electric fields here); with the relevant effects being:

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Diamond

Hamiltonian

Energy Levels

V as Qubit

Gates

Measurement

We're really interested in the spin state of the electron. Primarily dependent on spin-spin and spin-magnetic field interactions (we'll ignore electric fields here); with the relevant effects being:

(Electron State) Zero-Field Splitting (ZFS): SDS

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond

Hamiltonian

Energy Levels

V as Qubit

Gates

Measuremer

We're really interested in the spin state of the electron. Primarily dependent on spin-spin and spin-magnetic field interactions (we'll ignore electric fields here); with the relevant effects being:

(Electron State) Zero-Field Splitting (ZFS): SDS

(Nuclear Spin) Quadrapole Interaction: IQI

**NV-Centers** in Diamond as a **Qubit Platform** 

Alex Heilman

Hamiltonian

We're really interested in the spin state of the electron. Primarily dependent on spin-spin and spin-magnetic field interactions (we'll ignore electric fields here); with the relevant effects being:

(Electron State) Zero-Field Splitting (ZFS): SDS

(Nuclear Spin) Quadrapole Interaction: IQI

(Spin-Spin) Hyper-fine Interactions: SAI

**NV-Centers** in Diamond as a **Qubit Platform** 

Alex Heilman

Hamiltonian

Primarily dependent on spin-spin and spin-magnetic field interactions (we'll ignore electric fields here); with the relevant effects being:

(Electron State) Zero-Field Splitting (ZFS): SDS

(Nuclear Spin) Quadrapole Interaction: IQI

(Spin-Spin) Hyper-fine Interactions: SAI

(Spin-B Field) Zeeman:  $g_sSB$ ,  $g_iI_iB$ 

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

efects in Diamond

Hamiltonian

Energy Levels

V as Qubit

Measuremer

We're really interested in the spin state of the electron. Primarily dependent on spin-spin and spin-magnetic field interactions (we'll ignore electric fields here); with the relevant effects being:

(Electron State) Zero-Field Splitting (ZFS): SDS

(Nuclear Spin) Quadrapole Interaction: IQI

(Spin-Spin) Hyper-fine Interactions: SAI

(Spin-B Field) Zeeman:  $g_sSB$ ,  $g_iI_iB \leftarrow$  This we can control!

We're really interested in the spin state of the electron. Primarily dependent on spin-spin and spin-magnetic field interactions (we'll ignore electric fields here); with the relevant effects being:

(Electron State) Zero-Field Splitting (ZFS): SDS

(Nuclear Spin) Quadrapole Interaction: IQI

(Spin-Spin) Hyper-fine Interactions: SAI

(Spin-B Field) Zeeman:  $g_sSB$ ,  $g_iI_iB \leftarrow$  This we can control!

Let's see the Hamiltonian...

Hamiltonian Simplification

Energy Levels

Initialization Gates

Gates Measuremer

The electron will be taken to be localized to the defect. We'll take this localized two electron state to be the unperturbed solution and apply some interaction effects, with the corresponding interaction Hamiltonian below (again, assuming no external electric field/Stark effects):

$$H = \underbrace{\hat{S}\mathbf{D}\hat{S}}_{\mathsf{ZFS}} + \underbrace{\hat{I}_{N}\mathbf{Q}\hat{I}_{N}}_{\mathsf{Quadrapole}} + \underbrace{\hat{S}(\mathbf{A}_{N}\hat{I}_{N}^{'} + \sum_{i} \mathbf{A}_{C_{i}}\hat{I}_{C_{i}})}_{\mathsf{Hyperfine}} + \underbrace{(\hat{S}\mathbf{g}_{s} + \hat{I}_{N}\mathbf{g}_{N} + \sum_{i}\hat{I}_{C_{i}}\mathbf{g}_{C_{i}})B}_{\mathsf{Zeeman}} [3]$$

### **Simplifications**

Assuming we apply a magnetic field only along the symmetry axis, which we define to be the z direction

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond

Simplifications

Energy Levels

VV as Qubit

Moscuromont

Diamond
Hamiltonian
Simplifications

NV as Qubi

Gates Measurement

Assuming we apply a magnetic field only along the symmetry axis, which we define to be the z direction, and neglecting the surrounding carbon's effects (both Zeeman and hyperfine, though these allow us to use the nv-center as register for these surrounding sites [1]):

$$H_{NV} \approx DS_z^2 + \omega_e S_z + QI_z^2 + \omega_n I_z + AS_z I_z$$
 [2]

where  $\omega_i = \gamma_i B$ , the Larmor frequency.

 $D=2\pi \times (2.87\, GHz)$ , the dipole coupling constant  $Q=2\pi \times (-4.95\, GHz)$ , the nuclear quadrapole coupling constant

 $A = 2\pi \times (-2.16\,\text{GHz})$ , the hyperfine coupling constant


Hamiltonian Simplifications

Energy Levels

NV as Qubit

Gates

#### Triplet



# Singlet



# $m_s=\pm 1$ $m_s=-1$ Zeeman Effect $m_s=0$ 3 A

# **Qubit States**

The localized spin states of the defect may be used as a two-level system.

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Diamond
Hamiltonian
Simplifications

NV as Qubit

Gates


Energy Levels

NV as Qubit

Gates

The localized spin states of the defect may be used as a two-level system.

We make the identifications:



The localized spin states of the defect may be used as a two-level system.

We make the identifications:

$${}^{3} ext{A}_{1} egin{array}{c} m_{s} = 1 \ m_{s} = -1 \ m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \ m_{s} = 0 \ \end{array} egin{array}{c} m_{s} = 0 \ m_{s} = 0 \$$

The transitions between these states are in the microwave regime.


Defects in Diamond Hamiltonian

Energy Levels

Initialization

Measuremer

Initialization to the zero state can be achieved by an asymmetric relaxation from the first excited triplet state  $(^3E)$  to the triplet ground state  $(^3A)$ .



Applying a resonant pulse (532 nm) excites all triplet ground states to their corresponding excited states ( $\Delta m_s = 0$ ).

 $m_s = 0$  ground state.

 $m_s = 0$ 

efects in liamond

NV as Qubit

Initialization

 $m_s=\pm 1$   $m_s=0$  3E 1E  $1A_2$ 


Excited  $m_s=0$  states have been shown to decay back into  $m_s=0$  ground states, but  $m_s=\pm 1$  excited states favor a non-radiative (vibrational) decay mode via the singlet states, back into the

This may be exploited to intialize the spin state into our  $|0\rangle$  state.

JV as Qubit

Gates

Perpendicular (polarization relative symmetry axis) microwave pulses can induce Rabi flopping of state between  $|0\rangle$  and  $|1\rangle$ 




[6]

Energy Levels

Gates Measurement

Measuremen

Much like the initialization technique, optically detected magnetic resonance (ODMR) takes advantage of the asymmetric relaxation modes of the excited state.



Hence, if the defect flouresces when hit with a resonant pulse, it was measured to be in the  $|0\rangle$  state, but if it's 'dark' after a resonant pulse, it can be considered to have been in the  $|1\rangle$  state.

### Measurement Cont.

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

fects in amond miltonian applifications ergy Levels

NV as Qubit

Gates

Measurement

Note also that these allows our read-out to essentially be part of the next computation's initialization.

Measurement

Note also that these allows our read-out to essentially be part of the next computation's initialization.

One problem, however, is the possibility of internal reflection of the emitted photon.

Measurement

Note also that these allows our read-out to essentially be part of the next computation's initialization.

One problem, however, is the possibility of internal reflection of the emitted photon. This can be solved with an immersion lens built into the diamond [4].

## Conclusion & Outlook

Promising candidate due to practical methods of initialization, measurement, and qubit manipulation.

NV-Centers in Diamond as a Qubit Platform

Alex Heilman

Qubit Criteria

Defects in Diamond

Hamiltonian Simplifications Energy Levels

IV as Qubit

Measuremer

Defects in Diamond

Hamiltonian Simplification Energy Levels

13.7

V as Qubit

Measuremen

Promising candidate due to practical methods of initialization, measurement, and qubit manipulation.

Long coherence times, room temperature stability

Promising candidate due to practical methods of initialization, measurement, and qubit manipulation.

Long coherence times, room temperature stability

Haven't discussed scalability, two (or more) qubit operations, quantum networking, etc.

Promising candidate due to practical methods of initialization, measurement, and qubit manipulation.

Long coherence times, room temperature stability

Haven't discussed scalability, two (or more) qubit operations, quantum networking, etc.

Recent works have used neighboring spins (of carbon and nitrogen) as additional qubits [1] or used spin-photon coupling to get distant centers to interact [5]

Promising candidate due to practical methods of initialization, measurement, and qubit manipulation.

Long coherence times, room temperature stability

Haven't discussed scalability, two (or more) qubit operations, quantum networking, etc.

Recent works have used neighboring spins (of carbon and nitrogen) as additional qubits [1] or used spin-photon coupling to get distant centers to interact [5]

Some have also considered defect levels as qutrits [2]

C. E. Bradley, J. Randall, M. H. Abobeih, R. C. Berrevoets, M. J. Degen, M. A. Bakker, M. Markham, D. J. Twitchen, and T. H. Taminiau.

A ten-qubit solid-state spin register with quantum memory up to one minute.

Phys. Rev. X, 9:031045, Sep 2019.

Yue Fu, Wenquan Liu, Xiangyu Ye, Ya Wang, Chengjie Zhang, Chang-Kui Duan, Xing Rong, and Jiangfeng Du. Experimental investigation of quantum correlations in a two-qutrit spin system.

Phys. Rev. Lett., 129:100501, Aug 2022.

Ádám Gali.

Ab initio theory of the nitrogen-vacancy center in diamond.

Nanophotonics, 8(11):1907–1943, 2019.

Alex Heilman

JP Hadden, JP Harrison, Antony C Stanley-Clarke, L Marseglia, Y-LD Ho, BR Patton, Jeremy L O'Brien, and JG Rarity.

Strongly enhanced photon collection from diamond defect centers under microfabricated integrated solid immersion lenses

Applied Physics Letters, 97(24):241901, 2010.

Kae Nemoto, Michael Trupke, Simon J Devitt, Burkhard Scharfenberger, Kathrin Buczak, Jörg Schmiedmayer, and William J Munro.

Photonic quantum networks formed from nv- centers. Scientific reports, 6(1):1–12, 2016.

Alex Heilman

Lucio Robledo, Hannes Bernien, Ilse van Weperen, and Ronald Hanson.

Control and coherence of the optical transition of single nitrogen vacancy centers in diamond.

Phys. Rev. Lett., 105:177403, Oct 2010.