

Localization, Factorization and Dualities for Elliptic Kernels

Alessio Fontanarossa^{1,2}, Fabrizio Nieri^{1,2}, and Antonio Pittelli^{1,2}

¹*Dipartimento di Matematica, Università di Torino, Via Carlo Alberto 10, 10123
Torino, Italy*

²*INFN, Sezione di Torino, Via Pietro Giuria 1, 10125 Torino, Italy*

Abstract

We study the exact partition function of 4d $\mathcal{N} = 1$ supersymmetric gauge theories on a torus times a cylinder $\text{Cyl} = I \times S^1$, where I is a finite interval carrying two boundary components. Each endpoint supports an independent Dirichlet or Robin-like boundary polarization, so that the partition function is a boundary-to-boundary elliptic kernel. We construct the rigid supersymmetric geometry, determine the BPS locus, and compute the chiral-multiplet 1-loop determinants for the four possible boundary polarizations via equivariant localization. The resulting elementary building blocks are theta functions dressed by cubic phases. We then prove rank-changing Seiberg-type dualities as identities of Jeffrey–Kirwan residues of these elliptic kernels. We also discuss factorization into holomorphic-block cap wavefunctions represented by elliptic Gamma functions, dimensional reductions to three and two dimensions, complete-intersection gauged linear sigma models, and elliptic kernels for 4d $\mathcal{N} = 4$ super Yang–Mills and the Klebanov–Witten theory, useful for holographic applications.

Contents

1	Introduction and Summary	2
2	Supersymmetric Geometry	8
2.1	Background and domain	8
2.2	Bilinears as tangent vectors and dual one-forms	10
3	Localization and BPS loci	11
4	One-loop determinants and Boundary Polarizations	14
4.1	Kernel equations, supercharge algebra, and adjointness	14
4.2	Index interpretation	15
4.3	Boundary polarizations and one-loop determinants	16

5	Applications and Examples	21
5.1	Dualities	21
5.1.1	$U(N_c) \leftrightarrow U(N_f - N_c)$	21
5.1.2	$SU(N_c) \leftrightarrow SU(N_f - N_c)$ projection	29
5.1.3	Theta-function confining identity	30
5.2	Complete intersections on the cylinder	30
5.3	Factorization and cap wavefunctions	35
5.3.1	Four-dimensional factorization on $\text{Cyl} \times T^2$	36
5.3.2	The elliptic kernel interpretation	38
5.3.3	Two-dimensional factorization on Cyl	41
5.4	Holographic theories on $\text{Cyl} \times T^2$	43
5.4.1	$\mathcal{N} = 4$ SYM with gauge group $SU(N)$	44
5.4.2	Klebanov–Witten theory	45
A	Conventions and Auxiliary Formulae	46
A.1	Spinors and sigma matrices	46
A.2	Shintani–Barnes regularization	48
A.3	Jeffrey–Kirwan residue conventions	49
B	Anomaly Equations	50
C	Lower Dimensional Limits	52
C.1	Three-dimensional limit	52
C.2	Two-dimensional limit	53
C.3	The Z-product and the cylinder determinant	54

1 Introduction and Summary

In supersymmetric and topological settings, exact amplitudes of quantum field theories on manifolds with boundaries naturally behave as kernels. In two dimensions, this viewpoint is standard: interval amplitudes compute overlaps of boundary states, cap or hemisphere amplitudes assign wavefunctions to boundary conditions, and factorization follows from inserting a complete set of supersymmetric ground states. Exact results for 2d $\mathcal{N} = (2, 2)$ gauge theories on a disk geometry (D^2) provide a concrete realization of this rich structure [1, 2]. A related lesson is that exact answers on compact spaces [3–5] often factorize into more elementary blocks attached to simpler pieces of the geometry, and gluing such blocks reconstructs indices and partition functions on closed backgrounds. In three dimensions, this far-reaching property has been widely explored for theories defined on manifolds admitting a genus-one Heegaard splitting, starting from [6, 7]. In four dimensions, the natural playground is provided by geometries containing a two-torus (T^2), most notably the localization of 4d $\mathcal{N} = 1$ theories on $D^2 \times T^2$ and the associated four-dimensional holomorphic blocks [8, 9].

Interval (I) geometries add a further layer: observables become kernels between two boundary sectors. This is already visible in three-dimensional indices on $I \times T^2$, where bulk multiplets couple to boundary $2d$ degrees of freedom and the supersymmetric observables are related to elliptic genera, open-string Witten indices and interval factorizations [10–12]. Related gluing, cigar, line-segment and slab constructions, including interfaces and boundary localization, have been developed in [13–18]. The purpose of this paper is to construct and analyze such kernels for 4d $\mathcal{N} = 1$ gauge theories on $\text{Cyl} \times T^2$, where the cylinder $\text{Cyl} = I \times S^1$ has two boundary components. Throughout the paper we use the equivalent descriptions $\text{Cyl} \times T^2 = (I \times S^1) \times T^2 \simeq I \times T^3$, depending on which aspect of the geometry we wish to emphasize. The title refers to elliptic kernels for a concrete reason: fixing two supersymmetric boundary conditions at the northern and southern ends, the interval direction makes the path integral a boundary-to-boundary amplitude, i.e. a transition matrix element between boundary sectors, while the torus makes this amplitude elliptic in the sense of the $\text{Cyl} \times pt/S^1/T^2$ hierarchy.¹ Eventually, supersymmetric localization turns the matrix element into a finite-dimensional contour integral of a meromorphic function of gauge and flavor fugacities, whose elementary factors are theta functions or, equivalently, products of elliptic-Gamma cap wavefunctions. Once the elliptic kernels are obtained, we study how they behave under changes of boundary conditions/polarizations, anomaly cancellation conditions and compatibility with non-perturbative dualities. We also study how to recover the lower-dimensional results by taking suitable limits and the concatenation of several kernels, possibly coupled to cap wavefunctions.

This paper sits at the intersection of three lines of work which naturally connect supersymmetric quantum field theory to geometry, representation theory and special functions. The first is localization on manifolds with boundary, where one must keep track not only of the bulk BPS locus but also of admissible boundary conditions, boundary terms and possible boundary degrees of freedom [1, 8, 10, 13, 14, 16, 25]. The second is the theory of holomorphic blocks and their elliptic lifts, where wavefunctions are assigned to elementary pieces of a background and gluing is expressed by integration over gauge fugacities [6, 7, 9]. The third is the study of dual boundary conditions and duality interfaces, where infrared equivalences of bulk theories act non-trivially on the space of boundary conditions and must be tested by anomaly matching and supersymmetric indices [26]. The present construction combines these ingredients in a setting with two independent boundaries in four dimensions and lowest supersymmetry.

¹We use the word “kernel” in the boundary-to-boundary sense appropriate to an interval geometry. This should be distinguished from another common usage in the literature on supersymmetric partition functions and elliptic hypergeometric integrals, where compact-space indices depending on two sets of flavor fugacities can be interpreted, for instance, as integral or distributional kernels; see e.g. the elliptic hypergeometric identities of [19, 20], related recent work on elliptic kernels and duality walls [21, 22], and the holomorphic-block/bad-theory literature [6, 7, 9, 23, 24]. In this paper we analyze only the elementary D/R boundary polarizations and their flip determinants. We leave for future work a kernel with matrix elements labelled by a complete basis of boundary conditions or boundary defects.

Main results. The paper starts from the study of the rigid supersymmetric background $\text{Cyl} \times T^2$, making the Killing-spinor bilinears and the supercharge vector explicit. The first main result is the equivariant index evaluation of the 1-loop determinants for the four basic boundary polarizations of chiral multiplets, namely for the four possible choices of Dirichlet (D) or Robin(-like) (R) boundary conditions at the two ends. These are the same conditions considered in [8] and further studied in [27]. The mixed polarizations are acyclic away from the resonance hyperplanes of the boundary problem, meaning that the relevant first-order complex has no admissible kernel or cokernel modes and hence contributes the unit determinant

$$z_{\text{DR}}(u) = z_{\text{RD}}(u) = 1. \quad (1.1)$$

The unmixed polarizations give the two non-trivial elementary kernels, built out of theta or elliptic-Gamma functions dressed by *cubic* polynomial phases in the relevant fugacities

$$z_{\text{DD}}(u) := e^{-\frac{i\pi}{3}P_3(-u)} \Theta(-u; \omega), \quad z_{\text{RR}}(u) := \frac{e^{\frac{i\pi}{3}P_3(u)}}{\Theta(u; \omega)}, \quad (1.2)$$

the ω parameter representing the rotational fugacity of the cylinder. For q -Pochhammer symbols, theta and elliptic-Gamma functions we follow the conventions of [28–30], as summarized in the appendix. In particular

$$P_3(u) := B_{3,3}(u; 1, \tau, \omega) + B_{3,3}(\omega - u; 1, \tau, \omega) = -\frac{\tau + 1}{2\omega\tau} (6u^2 - 6\omega u + \omega^2 + \tau). \quad (1.3)$$

is the relevant combination of *cubic* Bernoulli polynomials, τ being the torus modular parameter. This is a distinguished feature with respect to the three-dimensional case, where the polynomial phases are quadratic. Another key difference is the diverse factorization into cap wavefunctions, namely

$$\Theta(u; \omega) = \frac{1}{\Gamma_e(u; \tau, \omega) \Gamma_e(\omega - u; \tau, \omega)}. \quad (1.4)$$

For interacting theories and general gauge group G , the exact partition function can be written in a matrix-integral form, with 1-loop factors given by boundary-polarized cylinder blocks. Let u denote a gauge fugacity in the Cartan, Δ_G the set of roots, and ρ the weights of a chiral representation. For a chiral multiplet Φ_I of R-charge r_I , flavor fugacity ν_I , gauge representation $\mathcal{R}_I$ and boundary polarization $\mathbf{q}_I \in \{\text{DD}, \text{DR}, \text{RD}, \text{RR}\}$, we define the full chiral 1-loop determinant by

$$\begin{aligned} Z_{\Phi_I}^{\text{DD}}(u) &= \prod_{\rho \in \mathcal{R}_I} z_{\text{DD}}(\rho(u) + \nu_I + (r_I - 2)\gamma_R), \\ Z_{\Phi_I}^{\text{RR}}(u) &= \prod_{\rho \in \mathcal{R}_I} z_{\text{RR}}(\rho(u) + \nu_I + r_I\gamma_R), \\ Z_{\Phi_I}^{\text{DR}}(u) &= Z_{\Phi_I}^{\text{RD}}(u) = 1. \end{aligned} \quad (1.5)$$

The non-zero-weight contribution of a vector multiplet with Neumann boundary conditions is obtained from the same first-order complex as a chiral multiplet with Dirichlet conditions in the

adjoint representation with effective R-charge $r = 2$, after the usual ghost and Cartan zero-mode cancellations. This gives a 1-loop factor $z_{\text{DD}}(\alpha(u))$ for each root $\alpha \in \Delta_G$

$$Z_{\text{vec}}^G(u) = \prod_{\alpha \in \Delta_G} z_{\text{DD}}(\alpha(u)). \quad (1.6)$$

The total localized expression is thus

$$\mathcal{Z}_{\text{Cyl} \times T^2}^{G, \{\mathbf{q}_I\}} = \frac{1}{|W_G|} \oint_{\mathcal{C}_\eta} \prod_{a=1}^{\text{rk } G} \frac{du_a}{2\pi i} Z_{\text{vec}}^G(u) \prod_I Z_{\Phi_I}^{\mathbf{q}_I}(u). \quad (1.7)$$

Boundary degrees of freedom, when present, multiply the integrand by additional factors. In the main text we consider the flip determinants between RR and DD conditions only. The contour prescription is a key ingredient, and we use the Jeffrey–Kirwan (JK) residue [31–35].

The second main result is the boundary-flip calculus on a two-boundary geometry. On a disk, a boundary multiplet can flip one boundary condition into another by multiplying the bulk determinant by a lower-dimensional determinant. On a cylinder, the effect of flipping one end depends on what is imposed at the other end. For this reason we keep track of all elementary flips at the northern and southern boundaries. The resulting flip factors are full theta-function determinants of the cylinder kernel. Elliptic-Gamma functions appear when the cylinder determinant is factorized into cap wavefunctions. This separation between boundary determinants and cap wavefunctions is one of the conceptual points of the paper.

The third main result is a rank-changing duality of Seiberg type [36]. The electric theory has gauge group $U(N_c)$, while the magnetic theory has gauge group $U(N_f - N_c)$. The JK residues on the two sides can be matched term by term. Electric poles are labelled by subsets $\mathcal{I}$ of N_c flavor labels, magnetic poles by complementary subsets $\mathcal{J} = \mathcal{I}^c$ of cardinality $N_f - N_c$. At the endpoint $N_f = N_c$, the magnetic gauge group becomes trivial and the same identity reduces to a theta-function confining evaluation formula, which is new as far as we know. From the viewpoint of special functions, the resulting identities are close in spirit to elliptic hypergeometric residue formulae and elliptic beta-type evaluations [19, 20]. From the viewpoint of field theory, they are the cylinder analogues of integral identities produced by supersymmetric dualities.

The fourth main result is the analysis of dimensional limits. In three dimensions, the cylinder blocks reduce to theta-function determinants factorizable into pairs of q -Pochhammer symbols, each representing a cap wavefunction. In two dimensions, they degenerate to trigonometric determinants factorizable into pairs of Euler-Gamma functions. The complete-intersection examples reproduce the expected equivalence of geometric and Landau–Ginzburg residue chambers, in the spirit of the GLSM phase structure of [37].

Last but not least, matter contents relevant to holographic SCFTs are analyzed in order to pave the way for some of the future research directions. For instance, writing $u_{ab} := u_a - u_b$, $SU(N)$ $\mathcal{N} = 4$ admits the following cylinder partition function, obtained by taking two adjoint

chirals in RR polarization and one in DD polarization,

$$Z_{\mathcal{N}=4}^{SU(N)}(\Delta_1, \Delta_2, \Delta_3) = \frac{[z_{\text{RR}}(\Delta_1) z_{\text{RR}}(\Delta_2) z_{\text{DD}}(\Delta_3)]^{N-1}}{N!} \oint_{\mathcal{C}_{SU(N)}} \prod_{a=1}^{N-1} \frac{du_a}{2\pi i} \\ \times \prod_{\substack{a,b=1 \\ a \neq b}}^N z_{\text{DD}}(u_{ab}) z_{\text{RR}}(u_{ab} + \Delta_1) z_{\text{RR}}(u_{ab} + \Delta_2) z_{\text{DD}}(u_{ab} + \Delta_3). \quad (1.8)$$

with $u_N = -\sum_{a=1}^{N-1} u_a$. Here the Δ_I are the fugacities of the three adjoint chiral multiplets. Equivalently, keeping the Bernoulli contact phase explicit, the same partition function can be written directly in terms of theta functions as

$$Z_{\mathcal{N}=4}^{SU(N)}(\Delta_1, \Delta_2, \Delta_3) = \frac{1}{N!} \oint_{\mathcal{C}_{SU(N)}} \prod_{a=1}^{N-1} \frac{du_a}{2\pi i} \exp\left[\frac{i\pi}{3} \mathcal{P}_{\mathcal{N}=4}^{SU(N)}(\Delta)\right] \\ \times \left[\frac{\Theta(-\Delta_3; \omega)}{\Theta(\Delta_1; \omega) \Theta(\Delta_2; \omega)} \right]^{N-1} \prod_{\substack{a,b=1 \\ a \neq b}}^N \frac{\Theta(-u_{ab}; \omega) \Theta(-(u_{ab} + \Delta_3); \omega)}{\Theta(u_{ab} + \Delta_1; \omega) \Theta(u_{ab} + \Delta_2; \omega)}. \quad (1.9)$$

Using (1.3), the Bernoulli contact phase in the theta-function presentation simplifies to the gauge-independent polynomial

$$\mathcal{P}_{\mathcal{N}=4}^{SU(N)}(\Delta) = - \frac{(N-1)(\tau+1)(\omega^2 + \tau)}{2\omega\tau} \\ - \frac{3(N^2-1)(\tau+1)}{\omega\tau} \left(\Delta_1^2 + \Delta_2^2 - \Delta_3^2 - \omega(\Delta_1 + \Delta_2 + \Delta_3) \right). \quad (1.10)$$

All dependence on the gauge fugacities cancels in (1.10); before the cancellation the possible linear terms are proportional to $\sum_{a \neq b} u_{ab} = 0$, while the quadratic u_{ab} -terms cancel root by root. Thus (1.8) is the compact block form, while (1.9) is the fully equivalent theta-function form. The contour is again specified by the JK prescription. This finite-rank formula is one of the inputs for a possible large- N analysis of cylinder kernels and their comparison with supersymmetric gravitational backgrounds containing cylindrical factors, in the spirit of the twisted-index, spindle-index and black-hole microstate literature [38–42].

Outlook. Several directions remain open. A first one is the systematic inclusion of boundary interactions, anomaly inflow [43] and defects or interfaces [26]. The two ends of the cylinder suggest a categorical interpretation: boundary polarizations should define objects, cylinder kernels should define morphisms or pairings, and boundary matter should act by composition, in analogy with the categorical description of branes and boundary conditions in supersymmetric boundary field theory, topological strings and Landau–Ginzburg models [44–47].

A second direction is a classification of cylinder dualities. The Seiberg-like determinant-pair family studied here is deliberately conservative, because it is the class for which JK chambers and complementary residue matching can all be controlled explicitly. The anomaly-screening equations allow more general matter assignments, including adjoint and flavor-adjoint variants.

A third direction may connect the present construction to derived categories of boundary conditions [45–47], analytic kernels for elliptic special functions [28, 29], and the tt^* interpretation

of block fusion [48, 49]. The word “fusion” is meant here in the broad sewing/gluing sense familiar from conformal field theory and modular tensor category constructions [50, 51]. At the level established in this paper the concrete algebraic statements are the reflected inverse relation for the two unmixed blocks and the flip identities (4.41)–(4.42); associativity of more general boundary-matter compositions is left for future work.

A fourth direction is the large-rank limit. Since the observable is a boundary-to-boundary kernel rather than a closed-space index, its saddle-point interpretation should differ from standard compact-space formulas. The $SU(N)$ $\mathcal{N} = 4$ contour integral (1.8) is the most direct starting point. A holographic interpretation, if any, should be closer to a transition amplitude in an asymptotically locally AdS_5 spacetime with two three-dimensional boundary components than to a closed-space partition function. In a local decompactification limit $T^3 \simeq \mathbb{R}^3$, the relevant bulk geometry may admit an AdS_4 -sliced description, namely a warped $AdS_4 \times I$ form. This is the $D = 4$ instance of the general $AdS_D \times I$ interval geometries studied recently in axio-dilaton gravity, while related defect-CFT constructions recover observables from warped $AdS_p \times S^q \times I$ backgrounds [52–54]. It would be interesting to understand whether the two boundary components give rise to two gravitational blocks [55] and whether their gluing reproduces black-string, black-brane or accelerating-black-hole observables [56] in backgrounds containing an elliptic factor, as happens in related compact computations [38–42].

Finally, one can study degenerations of the cylinder. If one boundary collapses, becomes asymptotic or is replaced by an orbifold point, the cylinder kernel should interpolate between the present two-boundary construction, elliptic disk blocks [8], twisted indices on hyperbolic manifolds [25, 57] and localization on orbifold geometries [41, 58, 59]. Understanding this interpolation may clarify which degrees of freedom are genuinely localized at a boundary or singular point and which are artifacts of a chosen polarization.

Outline. Section 2 constructs the supersymmetric background by solving the conformal new minimal equations. The Killing-spinor bilinears are evaluated explicitly and used to identify the equivariant parameters entering localization.

Section 3 describes the BPS locus. The vector multiplet equations imply that a particular combination of boundary holonomies is constant along the interval; this constant forms the complex gauge fugacity together with the toric holonomies.

Section 4 derives the 1-loop determinants using the cohomological approach. The elementary flips of boundary conditions at the two boundaries are listed and the distinction between full theta-function boundary determinants and elliptic-Gamma cap wavefunctions explained.

Section 5 contains the main applications. Subsection 5.1 proves the Seiberg-like duality between $U(N_c)$ and $U(N_f - N_c)$, discusses the SU projection and records the theta-function confining identity at the $U(0)$ endpoint. Subsection 5.2 studies complete-intersection GLSMs on the two-dimensional cylinder. Subsection 5.3 discusses factorization into cap wavefunctions, while Subsection 5.4 studies the consistency of cylinder integrals for matter contents familiar from holographic SCFTs.

[Appendix A](#) collects notation, boundary-condition labels, 1-loop determinant conventions, spinor and curvature conventions, Shintani–Barnes regularization, Jeffrey–Kirwan residue conventions. [Appendix B](#) records the anomaly-screening equations used to test candidate boundary polarizations. [Appendix C](#) discusses dimensional limits to three and two dimensions, distinguishing literal Shintani–Barnes evaluations from contact-term-normalized blocks.

Acknowledgements. The work of FN was partially supported by the CRT Foundation under the grant 108399/2024.0434.

2 Supersymmetric Geometry

2.1 Background and domain

Fix parameters

$$0 < \theta_N < \theta_S < \pi, \quad \theta \in [\theta_N, \theta_S], \quad \varphi \sim \varphi + 2\pi, \quad x \sim x + 2\pi, \quad y \sim y + 2\pi,$$

and define

$$\tau := \tau_1 + i\tau_2 \quad (\tau_2 > 0). \quad (2.1)$$

Let $h(\theta)$ be a smooth real-valued function on $[\theta_N, \theta_S]$. Throughout, $h(\theta)$ is assumed to be strictly positive on the closed interval:

$$h(\theta) > 0 \quad \text{for all } \theta \in [\theta_N, \theta_S]. \quad (2.2)$$

This condition ensures that the φ -fiber has strictly positive proper length at both boundary circles; if h vanishes at an endpoint, the φ -circle collapses and the geometry ceases to be a cylinder.

Let $f(\theta)$ be a smooth, real-valued and strictly positive function on $[\theta_N, \theta_S]$. Usually,

$$\lim_{\theta \rightarrow 0, \pi} f(\theta) \rightarrow b, \quad (2.3)$$

but we will not need it as the closed interval $[\theta_N, \theta_S]$ is strictly contained into $(0, \pi)$. Introduce the toric one-form

$$\Upsilon := d\varphi + \Omega_3 dx + \Omega_4 dy. \quad (2.4)$$

The metric is taken to be

$$ds^2 = L^2 f(\theta)^2 d\theta^2 + L^2 b^2 h(\theta)^2 \Upsilon^2 + L^2 \beta^2 \left((dx + \tau_1 dy)^2 + \tau_2^2 dy^2 \right). \quad (2.5)$$

Introduce the complex combination

$$\omega := \Omega_3 \tau - \Omega_4. \quad (2.6)$$

At fixed (x, y) one has $\Upsilon = d\varphi$, and the (θ, φ) -submanifold is a warped cylinder with boundary circles at $\theta = \theta_N$ and $\theta = \theta_S$, of proper lengths $2\pi L b h(\theta_N)$ and $2\pi L b h(\theta_S)$, respectively. The Riemannian density in coordinates (θ, φ, x, y) is

$$\sqrt{g} = L^4 b \beta^2 \tau_2 f(\theta) h(\theta). \quad (2.7)$$

The positivity assumption (2.2) guarantees non-degeneracy of (2.5) on the closed cylinder.

Consider the orthonormal coframe

$$e^1 = Lf(\theta) d\theta, \quad e^2 = Lbh(\theta) \Upsilon, \quad e^3 = L\beta(dx + \tau_1 dy), \quad e^4 = L\beta\tau_2 dy, \quad (2.8)$$

so that $ds^2 = \sum_{a=1}^4 (e^a)^2$.

Let $\omega_{ab} = -\omega_{ba}$ denote the Levi-Civita spin connection one-forms defined by the torsion-free Cartan equation

$$de^a + \omega^a{}_b \wedge e^b = 0. \quad (2.9)$$

In the coframe (2.8),

$$\omega_{12} = -\omega_{21} = -\frac{bh'(\theta)}{f(\theta)} \Upsilon, \quad (2.10)$$

and $\omega_{ab} = 0$ for all other pairs $(a, b) \neq (1, 2)$. Indeed, one has $de^1 = de^3 = de^4 = 0$ and, since $d\Upsilon = 0$,

$$de^2 = Lbh'(\theta) d\theta \wedge \Upsilon.$$

The $a = 2$ Cartan equation forces $\omega^2{}_3 = \omega^2{}_4 = 0$ and requires $\omega^2{}_1 = (bh'(\theta)/f(\theta)) \Upsilon$, which implies (2.10). The remaining Cartan equations are solved by setting all other components to zero.

Let E_a be dual to e^a . From (2.8),

$$\begin{aligned} E_1 &= \frac{1}{Lf(\theta)} \partial_\theta, \\ E_2 &= \frac{1}{Lbh(\theta)} \partial_\varphi, \\ E_3 &= \frac{1}{L\beta} (\partial_x - \Omega_3 \partial_\varphi), \\ E_4 &= \frac{1}{L\beta\tau_2} (\partial_y - \tau_1 \partial_x - (\Omega_4 - \Omega_3 \tau_1) \partial_\varphi). \end{aligned} \quad (2.11)$$

We use the Euclidean spinor and sigma-matrix conventions collected in Appendix A. In particular, curved sigma matrices are obtained from the frame and its inverse, and the covariant derivatives on Weyl spinors are defined by (A.13).

The conformal new minimal equations are [4, 5, 60]

$$(\nabla_\mu - iA_\mu^C) \zeta + \frac{1}{4} \sigma_\mu \tilde{\sigma}^\nu (\nabla_\nu - iA_\nu^C) \zeta = 0, \quad (2.12)$$

$$(\nabla_\mu + iA_\mu^C) \tilde{\zeta} + \frac{1}{4} \tilde{\sigma}_\mu \sigma^\nu (\nabla_\nu + iA_\nu^C) \tilde{\zeta} = 0. \quad (2.13)$$

Take

$$A^C = \frac{1}{2} (\omega_{12} + \alpha_\varphi d\varphi + \alpha_x dx + \alpha_y dy), \quad (2.14)$$

and define

$$\Xi := \alpha_\varphi \varphi + \alpha_x x + \alpha_y y. \quad (2.15)$$

The choice (2.14) retains its form for general $h(\theta)$: the θ -dependence of the Levi-Civita connection is entirely captured by ω_{12} in (2.10), and its inclusion in A^C ensures that the conformal equations admit constant-frame spinors with purely toric phases.

A convenient solution is

$$\zeta_\alpha = \sqrt{\kappa_0} e^{\frac{i}{2}\Xi} \begin{pmatrix} 1 \\ 0 \end{pmatrix}_\alpha, \quad \tilde{\zeta}^{\dot{\alpha}} = \sqrt{\kappa_0} e^{-\frac{i}{2}\Xi} \begin{pmatrix} 0 \\ 1 \end{pmatrix}^{\dot{\alpha}}, \quad (2.16)$$

with constant $\kappa_0 \in \mathbb{C}$. The pair $\zeta, \tilde{\zeta}$ has opposite chirality and opposite $U(1)_R$ charge; the supercharge used below is the real combination whose square closes on the vector field $K = \zeta \sigma \tilde{\zeta}$, together with gauge and R-symmetry transformations. Periodicity along T^2 and the boundary of the cylinder implies

$$\alpha_\varphi, \alpha_x, \alpha_y \in \mathbb{Z}. \quad (2.17)$$

The new minimal Killing spinor equations are [4, 5, 60]

$$(\nabla_\mu - iA_\mu)\zeta + iV_\mu\zeta + iV^\nu\sigma_{\mu\nu}\zeta = 0, \quad (2.18)$$

$$(\nabla_\mu + iA_\mu)\tilde{\zeta} - iV_\mu\tilde{\zeta} - iV^\nu\tilde{\sigma}_{\mu\nu}\tilde{\zeta} = 0, \quad (2.19)$$

and are solved by the spinors (2.16) provided the background fields

$$V = L\beta\kappa(dx + \tau dy), \quad A = A^C + \frac{3}{2}V, \quad (2.20)$$

are chosen, where $\kappa \in \mathbb{C}$ is constant.

2.2 Bilinears as tangent vectors and dual one-forms

Introduce the bilinears as tangent vectors

$$K := K^\mu \partial_\mu, \quad Y := Y^\mu \partial_\mu, \quad \tilde{Y} := \tilde{Y}^\mu \partial_\mu, \quad \tilde{K} := \tilde{K}^\mu \partial_\mu. \quad (2.21)$$

We use the following normalization for the vector bilinears:

$$K^\mu := \zeta \sigma^\mu \tilde{\zeta}, \quad Y^\mu := \frac{1}{2|\tilde{\zeta}|^2} \zeta \sigma^\mu \tilde{\zeta}^\dagger, \quad \tilde{Y}^\mu := -\frac{1}{2|\zeta|^2} \zeta^\dagger \sigma^\mu \tilde{\zeta}, \quad \tilde{K}^\mu := \frac{1}{4|\zeta|^2 |\tilde{\zeta}|^2} \tilde{\zeta}^\dagger \tilde{\sigma}^\mu \zeta^\dagger. \quad (2.22)$$

The metric dual one-forms are denoted by the musical isomorphism

$$K^\flat := g_{\mu\nu} K^\nu dx^\mu, \quad Y^\flat := g_{\mu\nu} Y^\nu dx^\mu, \quad \tilde{Y}^\flat := g_{\mu\nu} \tilde{Y}^\nu dx^\mu, \quad \tilde{K}^\flat := g_{\mu\nu} \tilde{K}^\nu dx^\mu. \quad (2.23)$$

Using $\sigma_{\alpha\dot{\alpha}}^\mu \tilde{\sigma}_\mu^{\dot{\beta}\beta} = -2\delta_\alpha^\beta \delta_{\dot{\alpha}}^{\dot{\beta}}$, one obtains

$$K^\mu (K_\mu)^* = 4|\zeta|^2 |\tilde{\zeta}|^2 K^\mu \tilde{K}_\mu, \quad K^\mu \tilde{K}_\mu = \frac{1}{2}, \quad (2.24)$$

and, consistently with (2.22),

$$(K^\mu)^* = 4|\zeta|^2 |\tilde{\zeta}|^2 \tilde{K}^\mu, \quad (Y^\mu)^* = -\frac{|\zeta|^2}{|\tilde{\zeta}|^2} \tilde{Y}^\mu, \quad (\tilde{Y}^\mu)^* = -\frac{|\tilde{\zeta}|^2}{|\zeta|^2} Y^\mu. \quad (2.25)$$

Using (2.11), one finds

$$K = \frac{i\kappa_0}{L\beta\tau_2} (\omega \partial_\varphi - \tau \partial_x + \partial_y), \quad (2.26)$$

$$Y = \frac{1}{2Lf(\theta)} \partial_\theta - \frac{i}{2Lbh(\theta)} \partial_\varphi, \quad (2.27)$$

$$\tilde{Y} = -\frac{1}{2Lf(\theta)} \partial_\theta - \frac{i}{2Lbh(\theta)} \partial_\varphi, \quad (2.28)$$

and

$$\widetilde{K} = \frac{1}{4|\zeta|^2|\widetilde{\zeta}|^2} K^* = \frac{1}{4|\kappa_0|^2} K^*. \quad (2.29)$$

The metric dual one-forms are

$$K^\flat = L\beta \kappa_0 (dx + \tau dy), \quad (2.30)$$

$$\begin{aligned} Y^\flat &= \frac{Lf(\theta)}{2} d\theta - \frac{iLbh(\theta)}{2} \Upsilon, \\ \widetilde{Y}^\flat &= -\frac{Lf(\theta)}{2} d\theta - \frac{iLbh(\theta)}{2} \Upsilon, \end{aligned} \quad (2.31)$$

and

$$\widetilde{K}^\flat = \frac{L\beta}{4\kappa_0} (dx + \bar{\tau} dy), \quad \bar{\tau} = \tau_1 - i\tau_2. \quad (2.32)$$

Since the components of $K^\flat$ are constant in the coordinate basis and the vector K is orthogonal to the fibre direction, one finds

$$\nabla_\mu K_\nu = 0. \quad (2.33)$$

In particular, K is Killing. The contraction with the toric one-form Υ is

$$\iota_K \Upsilon = 0. \quad (2.34)$$

3 Localization and BPS loci

The localization setup below uses the standard cohomological form of rigid supersymmetry on curved backgrounds, adapted to the presence of supersymmetric boundary conditions as in the disk and hyperbolic computations of Refs. [4, 5, 8, 25, 61, 62].

Define $\varrho = \varrho(\theta)$ by

$$\frac{d\varrho}{d\theta} = \frac{f(\theta)}{bh(\theta)}, \quad \varrho(\theta) := \int_{\theta_*}^{\theta} \frac{f(\theta')}{bh(\theta')} d\theta'. \quad (3.1)$$

Under (2.2), $\varrho(\theta)$ is smooth and strictly monotone if $f(\theta)$ stays positive on $[\theta_N, \theta_S]$, which holds for the present $f(\theta)$ in (2.3). Consequently, ϱ provides a global coordinate along the cylinder direction and is adapted to the $(Y, \widetilde{Y})$ -complex.

Then

$$E_1 = \frac{1}{Lbh(\theta)} \partial_\varrho, \quad E_2 = \frac{1}{Lbh(\theta)} \partial_\varphi,$$

so that (up to a nonzero scalar factor)

$$Y \propto \partial_\varrho + i\partial_\varphi, \quad \widetilde{Y} \propto \partial_\varrho - i\partial_\varphi. \quad (3.2)$$

Introduce

$$z := e^{\varrho+i\varphi}, \quad \bar{z} := e^{\varrho-i\varphi}. \quad (3.3)$$

For a field X of R -charge q_R and gauge charge q_G define

$$D_\mu X := \nabla_\mu X - iq_R A_\mu X - iq_G \mathcal{A}_\mu X. \quad (3.4)$$

Define

$$L_Y := Y^\mu D_\mu, \quad L_{\tilde{Y}} := \tilde{Y}^\mu D_\mu. \quad (3.5)$$

Along the φ -circle at fixed (x, y) one has $\Upsilon = d\varphi$, hence

$$A_\varphi = \frac{1}{2} \left(\alpha_\varphi - \frac{b h'(\theta)}{f(\theta)} \right), \quad \mathcal{A}_\varphi = \mathcal{A}(\partial_\varphi). \quad (3.6)$$

Using (3.1), one checks

$$\frac{b h'(\theta)}{f(\theta)} = \frac{d}{d\theta} (\log h(\theta)). \quad (3.7)$$

A vector multiplet $(\mathcal{A}_\mu, \lambda, \tilde{\lambda}, D)$ has supersymmetry variations

$$\begin{aligned} \delta \mathcal{A}_\mu &= i \tilde{\zeta} \tilde{\sigma}_\mu \lambda + i \zeta \sigma_\mu \tilde{\lambda}, \\ \delta \lambda &= \mathcal{F}_{\mu\nu} \sigma^{\mu\nu} \zeta + i D \zeta, \\ \delta \tilde{\lambda} &= \mathcal{F}_{\mu\nu} \tilde{\sigma}^{\mu\nu} \tilde{\zeta} - i D \tilde{\zeta}, \\ \delta D &= \tilde{\zeta} \tilde{\sigma}^\mu \left(D_\mu \lambda + \frac{3i}{2} V_\mu \lambda \right) - \zeta \sigma^\mu \left(D_\mu \tilde{\lambda} - \frac{3i}{2} V_\mu \tilde{\lambda} \right), \end{aligned} \quad (3.8)$$

where $\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu$ in the Abelian case, and D_μ is the appropriate covariant derivative on the gaugini.

A chiral multiplet (ϕ, ψ, F) of R -charge r has SUSY variations

$$\begin{aligned} \delta \phi &= \sqrt{2} \zeta \psi, \\ \delta \psi &= \sqrt{2} F \zeta + i \sqrt{2} \sigma^\mu \tilde{\zeta} D_\mu \phi, \\ \delta F &= i \sqrt{2} \tilde{\zeta} \tilde{\sigma}^\mu \left(D_\mu \psi - \frac{i}{2} V_\mu \psi \right) - 2i (\tilde{\zeta} \tilde{\lambda}) \phi, \end{aligned} \quad (3.9)$$

and for the anti-chiral multiplet $(\tilde{\phi}, \tilde{\psi}, \tilde{F})$,

$$\begin{aligned} \delta \tilde{\phi} &= \sqrt{2} \tilde{\zeta} \tilde{\psi}, \\ \delta \tilde{\psi} &= \sqrt{2} \tilde{F} \tilde{\zeta} + i \sqrt{2} \tilde{\sigma}^\mu \zeta D_\mu \tilde{\phi}, \\ \delta \tilde{F} &= i \sqrt{2} \zeta \sigma^\mu \left(D_\mu \tilde{\psi} + \frac{i}{2} V_\mu \tilde{\psi} \right) + 2i \tilde{\phi} (\zeta \lambda). \end{aligned} \quad (3.10)$$

Assume an Abelian ansatz depending only on θ ,

$$\mathcal{A} = (\mathcal{A}_\varphi(\theta) + \beta_\varphi) d\varphi + (\mathcal{A}_x(\theta) + \beta_x) dx + (\mathcal{A}_y(\theta) + \beta_y) dy. \quad (3.11)$$

Here $\beta_\varphi, \beta_x, \beta_y$ denote constant holonomies along the three circles (not to be confused with the metric parameter β in (2.5)).

A standard component of the vector-multiplet BPS equations in the cohomological presentation is

$$\iota_K \mathcal{F} = 0, \quad (3.12)$$

with $\mathcal{F} = d\mathcal{A}$ in the Abelian case. Under the assumption that the components in (3.11) depend only on θ , one has

$$\mathcal{F} = \mathcal{A}'_\varphi(\theta) d\theta \wedge d\varphi + \mathcal{A}'_x(\theta) d\theta \wedge dx + \mathcal{A}'_y(\theta) d\theta \wedge dy,$$

and using $\iota_K(d\theta) = 0$ and $\iota_K(d\varphi) = K^\varphi$, $\iota_K(dx) = K^x$, $\iota_K(dy) = K^y$, one finds

$$\iota_K \mathcal{F} = -\left(K^\varphi \mathcal{A}'_\varphi(\theta) + K^x \mathcal{A}'_x(\theta) + K^y \mathcal{A}'_y(\theta)\right) d\theta.$$

With K as in (2.26), the condition (3.12) is equivalent to the first-order ODE

$$\omega \mathcal{A}'_\varphi(\theta) - \tau \mathcal{A}'_x(\theta) + \mathcal{A}'_y(\theta) = 0, \quad (3.13)$$

which integrates to

$$\omega \mathcal{A}_\varphi(\theta) - \tau \mathcal{A}_x(\theta) + \mathcal{A}_y(\theta) = \mathcal{A}_0, \quad (3.14)$$

for some integration constant $\mathcal{A}_0 \in \mathbb{C}$. Equivalently, using (2.6),

$$\mathcal{A}_y(\theta) = \tau \mathcal{A}_x(\theta) + (\Omega_4 - \tau \Omega_3) \mathcal{A}_\varphi(\theta) + \mathcal{A}_0. \quad (3.15)$$

Introduce the component along the fibre direction,

$$\mathcal{A}_\Upsilon(\theta) := \mathcal{A}_\varphi(\theta) + \Omega_3 \mathcal{A}_x(\theta) + \Omega_4 \mathcal{A}_y(\theta). \quad (3.16)$$

Then $d\mathcal{A}$ contains $\mathcal{A}'_\Upsilon(\theta) d\theta \wedge \Upsilon$, and using (2.8),

$$\mathcal{F}_{12} = \frac{1}{L^2 b f(\theta) h(\theta)} \frac{d\mathcal{A}_\Upsilon}{d\theta}. \quad (3.17)$$

On the cohomological BPS locus, the auxiliary field is fixed accordingly:

$$D|_{\text{BPS}} = \frac{1}{L^2 b f(\theta) h(\theta)} \frac{d\mathcal{A}_\Upsilon}{d\theta}. \quad (3.18)$$

The chiral-multiplet BPS equations follow by setting to zero the fermionic variations in (3.9) and (3.10). On a contour where bosons and auxiliaries are taken to obey standard reality conditions, a consistent supersymmetric saddle is obtained by

$$\psi = \tilde{\psi} = 0, \quad F = \tilde{F} = 0, \quad (3.19)$$

together with the first-order constraints obtained from the kinetic terms in $\delta\psi$ and $\delta\tilde{\psi}$. In particular, the projected equations

$$\sigma^\mu \tilde{\zeta} D_\mu \phi = 0, \quad \tilde{\sigma}^\mu \zeta D_\mu \tilde{\phi} = 0 \quad (3.20)$$

can be expressed in the present background in terms of the first-order operators $L_{\tilde{\Upsilon}}$ and L_Y introduced in (3.5). Concretely, up to an overall nonzero scalar factor,

$$\sigma^\mu \tilde{\zeta} D_\mu \phi = 0 \iff L_{\tilde{\Upsilon}} \phi = 0, \quad \tilde{\sigma}^\mu \zeta D_\mu \tilde{\phi} = 0 \iff L_Y \tilde{\phi} = 0. \quad (3.21)$$

In the localization computation, the one-loop determinant is obtained by expanding around the BPS locus (and imposing supersymmetric boundary conditions on the two boundary circles).

Let $\mathcal{B}_2 = \{(\theta, \varphi) : \theta \in [\theta_N, \theta_S], \varphi \sim \varphi + 2\pi\}$ at fixed (x, y) , with boundary circles S_S^1 at $\theta = \theta_S$ and S_N^1 at $\theta = \theta_N$. For any Abelian connection $A_{(0)}$ on $\mathcal{B}_2$,

$$\frac{1}{2\pi} \int_{\mathcal{B}_2} dA_{(0)} = \frac{1}{2\pi} \oint_{S_S^1} A_{(0)} - \frac{1}{2\pi} \oint_{S_N^1} A_{(0)}. \quad (3.22)$$

For the vector multiplet connection (3.11),

$$\frac{1}{2\pi} \int_{\mathcal{B}_2} d\mathcal{A} = \mathcal{A}_\Upsilon(\theta_S) - \mathcal{A}_\Upsilon(\theta_N). \quad (3.23)$$

4 One-loop determinants and Boundary Polarizations

4.1 Kernel equations, supercharge algebra, and adjointness

For the spectral analysis, one expands fluctuations in Fourier modes along the three circles and treats the constant holonomies $\beta_\varphi, \beta_x, \beta_y$ as background parameters. In this setting the dependence on the BPS profile enters through the combination fixed by (3.14), while the mode labels remain in $\mathbb{Z}^3$.

Consider Fourier modes on the three-torus,

$$X(\theta, \varphi, x, y) = e^{i(m_\varphi \varphi + m_x x + m_y y)} F(\theta), \quad (m_\varphi, m_x, m_y) \in \mathbb{Z}^3. \quad (4.1)$$

Using (3.2) and (3.6), the kernel equations reduce to first-order equations on (ϱ, φ) :

$$L_Y X = 0 \iff (\partial_\varrho + iD_\varphi)X = 0, \quad (4.2)$$

$$L_{\tilde{Y}} X = 0 \iff (\partial_\varrho - iD_\varphi)X = 0, \quad (4.3)$$

where $D_\varphi = \partial_\varphi - iq_R A_\varphi - iq_G \mathcal{A}_\varphi$ and $\mathcal{A}_\varphi = \mathcal{A}(\partial_\varphi)$.

For $L_Y X = 0$, solutions can be written as

$$X_{m_\varphi, m_x, m_y}^{(Y)} = c_{m_\varphi, m_x, m_y}^{(Y)} \exp(i(m_\varphi \varphi + m_x x + m_y y)) \exp\left((m_\varphi - \frac{q_R}{2} \alpha_\varphi - q_G \mathcal{A}_\varphi) \varrho(\theta)\right) (h(\theta))^{\frac{q_R}{2}}. \quad (4.4)$$

For $L_{\tilde{Y}} X = 0$,

$$X_{m_\varphi, m_x, m_y}^{(\tilde{Y})} = c_{m_\varphi, m_x, m_y}^{(\tilde{Y})} \exp(i(m_\varphi \varphi + m_x x + m_y y)) \exp\left(-(m_\varphi + \frac{q_R}{2} \alpha_\varphi + q_G \mathcal{A}_\varphi) \varrho(\theta)\right) (h(\theta))^{-\frac{q_R}{2}}. \quad (4.5)$$

On a field X of charges (q_R, q_G) one has

$$\delta^2 X = 2i(\mathcal{L}_K X - iq_R \Phi_R X - iq_G \Phi_G X), \quad (4.6)$$

with $\Phi_R := \iota_K A$ and $\Phi_G := \iota_K \mathcal{A}$. Using (2.26) and (2.34), the spin-connection term in A^C drops out of the contraction and one finds

$$\Phi_R = \iota_K A = \frac{i\kappa_0}{L\beta\tau_2} \gamma_R, \quad \gamma_R := \frac{1}{2}(\omega\alpha_\varphi - \tau\alpha_x + \alpha_y). \quad (4.7)$$

For the Abelian BPS connection (3.11), the contraction with K gives

$$\Phi_G = \iota_K \mathcal{A} = \frac{i\kappa_0}{L\beta\tau_2} \left(\omega(\mathcal{A}_\varphi(\theta) + \beta_\varphi) - \tau(\mathcal{A}_x(\theta) + \beta_x) + (\mathcal{A}_y(\theta) + \beta_y) \right).$$

On the BPS locus, the θ -dependent part is constant by (3.14), hence

$$\Phi_G = \iota_K \mathcal{A} = \frac{i\kappa_0}{L\beta\tau_2} \gamma_G, \quad \gamma_G := \mathcal{A}_0 + \beta_y - \tau\beta_x + \omega\beta_\varphi. \quad (4.8)$$

The expression (4.8) makes explicit the three integer shifts of the underlying flat connection:

$$\beta_y \mapsto \beta_y + 1 : \gamma_G \mapsto \gamma_G + 1, \quad \beta_\varphi \mapsto \beta_\varphi + 1 : \gamma_G \mapsto \gamma_G + \omega, \quad \beta_x \mapsto \beta_x + 1 : \gamma_G \mapsto \gamma_G - \tau. \quad (4.9)$$

The Shintani–Barnes representative used for the cylinder determinant below is chosen so that the final theta functions have nome $e^{2\pi i\omega}$. Hence the elementary cylinder blocks are meromorphic functions of the gauge fugacity on the ω -elliptic lattice $\mathbb{Z} + \omega\mathbb{Z}$. All single-valuedness statements in the residue identities of this paper therefore refer to ω -ellipticity in the gauge fugacities, not to invariance under an independent shift by τ .

On Fourier modes (4.1),

$$\mathcal{L}_K X = \frac{i\kappa_0}{L\beta\tau_2} (\omega m_\varphi - \tau m_x + m_y) X,$$

hence

$$\delta^2 X_{m_\varphi, m_x, m_y} = -\frac{2i\kappa_0}{L\beta\tau_2} (\omega m_\varphi - \tau m_x + m_y - q_G \gamma_G - q_R \gamma_R) X_{m_\varphi, m_x, m_y}. \quad (4.10)$$

The common prefactor $-2i\kappa_0/(L\beta\tau_2)$ cancels in determinant ratios.

Let $\mathcal{F}$ be the L^2 completion of the relevant field space on the cylindrical region, with inner product

$$\langle X_1, X_2 \rangle := \int d^4x \sqrt{g} (X_1)^\dagger X_2, \quad (4.11)$$

and impose one of the supersymmetric boundary polarizations described below at each end of the interval. These polarizations are chosen so that the boundary bilinear generated by integrating L_Y by parts vanishes. Equivalently, the localization deformation δV has no uncanceled boundary variation on the allowed field space. We therefore use the standard localization argument only within a fixed boundary polarization: deformations of the positive localizing functional that preserve the same boundary complex do not change the determinant ratio, while changing the polarization is a different observable and is implemented by the flip factors of Subsection 4. With these assumptions one has

$$L_Y^\dagger = -L_{\tilde{Y}}, \quad (4.12)$$

and therefore

$$\text{coker}(L_Y) \cong \ker(L_{\tilde{Y}}), \quad \text{coker}(L_{\tilde{Y}}) \cong \ker(L_Y). \quad (4.13)$$

4.2 Index interpretation

The same algebraic data also has an index interpretation. We take $y = t_E$ to be the Euclidean time circle and use the Wick rotation $t = it_E$. With the Lorentzian Hamiltonian convention $\mathcal{H} = i\partial_t$, this gives $\mathcal{H} = \partial_y$. The path integral on $\text{Cyl} \times T^2$ is then a graded trace over the Hilbert space of the theory on $\text{Cyl} \times S_x^1$. The two boundary polarizations are kept fixed throughout this quantization. We use the convention, common in cohomological constructions of protected sectors, that symmetry generators act on local operators by commutators and hence by the corresponding differential operators; for instance the sphere construction of [63] writes the angular generator along the protected circle as $P_\varphi = i\partial_\varphi$. With the Fourier convention (4.1), we therefore introduce

$$J := i\partial_\varphi, \quad \mathcal{P} := i\partial_x, \quad -i\partial_y X_{m_\varphi, m_x, m_y} = m_y X_{m_\varphi, m_x, m_y}. \quad (4.14)$$

Thus a Fourier mode has eigenvalues $J = -m_\varphi$ and $\mathcal{P} = -m_x$. The vanishing of δ^2 on a state contributing to the protected cohomology is not the separate vanishing of each charge, but the vanishing of the equivariant combination appearing in (4.10). Equivalently, for a state of charges (q_R, q_G) ,

$$\delta^2 X_{\text{BPS}} = 0 \quad \Longleftrightarrow \quad \omega m_\varphi - \tau m_x + m_y - q_G \gamma_G - q_R \gamma_R = 0. \quad (4.15)$$

In terms of the operators (4.14), this becomes

$$-i\partial_y = \omega J - \tau \mathcal{P} + \gamma_G Q_G + \gamma_R Q_R \quad \text{on } \mathcal{H}_{\text{BPS}}, \quad (4.16)$$

where Q_G and Q_R denote the corresponding gauge, flavor, or R-symmetry charge operators. Therefore the Euclidean translation around the y -circle can be rewritten on Q -cohomology as

$$\exp(-2\pi\partial_y) = \exp\left[-2\pi i(\omega J - \tau \mathcal{P} + \gamma_G Q_G + \gamma_R Q_R)\right] \quad \text{on } \mathcal{H}_{\text{BPS}}. \quad (4.17)$$

Introducing

$$q := \exp(2\pi i \omega), \quad p := \exp(2\pi i \tau), \quad (4.18)$$

the cylinder partition function may thus be viewed as the protected index

$$\begin{aligned} Z_{\text{Cyl} \times T^2} &= \text{Tr}_{\mathcal{H}(\text{Cyl} \times S_x^1)} \left[(-1)^F \exp(-2\pi \mathcal{H}) \right] = \text{Tr}_{\mathcal{H}(\text{Cyl} \times S_x^1)} \left[(-1)^F \exp(-2\pi \partial_y) \right] \\ &= \text{Tr}_{\mathcal{H}_{\text{BPS}}} \left[(-1)^F q^{-J} p^{\mathcal{P}} \exp(-2\pi i(\gamma_G Q_G + \gamma_R Q_R)) \right]. \end{aligned} \quad (4.19)$$

With the Wick-rotation convention used above, $\mathcal{H} = \partial_y$ on the Euclidean functional integral. The protected trace (4.19) is independent of this convention once the Euclidean algebra (4.10) is kept fixed.

4.3 Boundary polarizations and one-loop determinants

The boundary of $\mathcal{B}_2$ consists of two circles at $\theta = \theta_N$ and $\theta = \theta_S$. In this paper a boundary polarization is simply the choice of supersymmetric boundary condition for the chiral multiplet at a given boundary circle. We allow Dirichlet (D) and Robin (R) boundary conditions. Since the cylinder has two ends, a bulk chiral multiplet is assigned an ordered pair of boundary conditions,

$$DD, \quad DR, \quad RD, \quad RR,$$

where the first label refers to the northern boundary and the second to the southern boundary.

In the cohomological presentation, Robin boundary conditions are characterized by compatibility with the δ -variation of the fermionic cohomological variable B . Schematically,

$$\delta B = L_{\tilde{Y}} \phi + \cdots, \quad (4.20)$$

and therefore Robin enforces

$$(L_{\tilde{Y}} \phi)|_{\partial} = 0. \quad (4.21)$$

In particular, bulk modes in $\ker L_{\tilde{Y}}$ satisfy (4.21) automatically. Written in the radial coordinate ϱ , this is a spectral Robin condition of the form $(\partial_\varrho - iD_\varphi)\phi|_{\partial} = 0$, with the connection and

charge-dependent terms contained in D_φ ; it is not an additional arbitrary parameter but the supersymmetric Robin operator paired with the Dirichlet condition $\phi|_\partial = 0$.

For mixed boundary choices, the cylinder supports no nontrivial cohomology for the first-order complex under generic background holonomies. This follows from the incompatibility of holomorphic/antiholomorphic transport data across the two boundary circles, as made explicit below.

Consider a field X obeying the bulk equation $L_Y X = 0$, hence of the form (4.4). Acting on such a mode with $L_{\tilde{Y}}$ yields

$$L_{\tilde{Y}} X_{m_\varphi, m_x, m_y}^{(Y)} = 2 \left(m_\varphi - \frac{q_R}{2} \alpha_\varphi - q_G \mathcal{A}_\varphi \right) \mathcal{W}(\theta) e^{i(m_\varphi \varphi + m_x x + m_y y)}, \quad (4.22)$$

where $\mathcal{W}(\theta)$ is a nonzero weight obtained from the $\varrho(\theta)$ and $h(\theta)$ factors in (4.4). In particular, at any boundary circle $\theta = \theta_*$ the Robin condition $L_{\tilde{Y}} X|_{\theta_*} = 0$ implies that each Fourier coefficient vanishes unless

$$m_\varphi - \frac{q_R}{2} \alpha_\varphi - q_G \mathcal{A}_\varphi = 0. \quad (4.23)$$

For generic holonomies and background parameters, (4.23) has no solution with $m_\varphi \in \mathbb{Z}$ and thus enforces $X \equiv 0$. The exceptional loci defined by (4.23) are resonance hyperplanes of the boundary problem. In the gauge-theory residue formulae below we assume the flavor masses and spectator fugacities are generic enough that the JK poles selected by the unmixed RR factors do not lie on these resonance hyperplanes. Under this genericity assumption the mixed-sector determinant is the meromorphic continuation of the identically trivial determinant on the complement of the resonance arrangement, hence it contributes the factor one at the selected poles. If a resonance hyperplane is tuned to pass through a JK pole, extra boundary zero modes may appear and the corresponding wall-crossed boundary theory must be specified separately.

An analogous statement holds with $Y \leftrightarrow \tilde{Y}$ interchanged. Consequently, for mixed boundary polarizations (DR or RD) one obtains, generically,

$$\ker L_Y = \ker L_{\tilde{Y}} = 0 \quad \implies \quad \text{coker}(L_Y) = \text{coker}(L_{\tilde{Y}}) = 0. \quad (4.24)$$

This provides the mechanism underlying the triviality of the bulk one-loop determinant for mixed polarizations. In this precise sense the mixed boundary complexes are acyclic: the cohomology of the first-order complex is zero, because both the kernel and the cokernel vanish on the allowed space of modes. Consequently the determinant ratio contains no non-trivial bulk factor and is equal to one, unless the parameters are tuned to a resonance hyperplane where extra boundary zero modes must be included explicitly.

After dropping the common prefactor in (4.10), the universal δ^2 eigenvalue factor on a Fourier mode labelled by $(m_\varphi, m_x, m_y) \in \mathbb{Z}^3$ and carrying charges (q_R, q_G) is

$$\lambda(m_\varphi, m_x, m_y; q_R, q_G) := \omega m_\varphi - \tau m_x + m_y - q_G \gamma_G - q_R \gamma_R. \quad (4.25)$$

For the Dirichlet polarization, the contributing cohomological scalar has effective charges

$$(q_R, q_G) = (r - 2, q_G), \quad (4.26)$$

hence the corresponding raw product is

$$Z_{1\text{-loop}}^{\text{CM}}|_D = \prod_{(m_\varphi, m_x, m_y) \in \mathbb{Z}^3} \left(\omega m_\varphi - \tau m_x + m_y - q_G \gamma_G - (r-2)\gamma_R \right). \quad (4.27)$$

For the Robin polarization, compatibility with (4.20) enforces (4.21). The relevant modes are those of the chiral scalar ϕ in $\ker L_{\tilde{Y}}$, carrying charges

$$(q_R, q_G) = (r, q_G). \quad (4.28)$$

With the placement of this sector in the denominator of the determinant ratio, the raw product reads

$$Z_{1\text{-loop}}^{\text{CM}}|_R = \prod_{(m_\varphi, m_x, m_y) \in \mathbb{Z}^3} \left(\omega m_\varphi - \tau m_x + m_y + q_G \gamma_G + r \gamma_R \right)^{-1}. \quad (4.29)$$

In order to regularize these products, we employ a scheme similar to [64, 65], namely we first consider the torus modes and then the cylinder ones by breaking the remaining product into two halves. Geometrically this splitting is the one-loop shadow of cutting the cylinder into two oppositely oriented caps: after regularization, each half is naturally interpreted as a $D^2 \times T^2$ cap wavefunction, while the full $\text{Cyl} \times T^2$ determinant is obtained by multiplying the two cap contributions and retaining the Bernoulli contact terms. Thus the factorization used later is already visible at the level of the regulated triple product:

$$\begin{aligned} Z_{1\text{-loop}}^{\text{CM}}|_D &= \prod_{m_\varphi \geq 0} \prod_{(m_x, m_y) \in \mathbb{Z}^2} (-1) \left(\tau m_x - m_y + \omega m_\varphi - q_G \gamma_G - (r-2)\gamma_R \right) \times \\ &\times \left(\tau m_x - m_y + \omega m_\varphi + \omega + q_G \gamma_G + (r-2)\gamma_R \right), \end{aligned} \quad (4.30)$$

and similarly

$$\begin{aligned} Z_{1\text{-loop}}^{\text{CM}}|_R &= \prod_{m_\varphi \geq 0} \prod_{(m_x, m_y) \in \mathbb{Z}^2} (-1) \left(\tau m_x - m_y + \omega m_\varphi + q_G \gamma_G + r \gamma_R \right)^{-1} \times \\ &\times \left(\tau m_x - m_y + \omega m_\varphi + \omega - q_G \gamma_G - r \gamma_R \right)^{-1}. \end{aligned} \quad (4.31)$$

Then, using the standard Shintani-Barnes regularization as summarized in Subsection A.2, such procedure gives us the final results (up to a normalization constant)

$$\begin{aligned} Z_{1\text{-loop}}^{\text{CM}}|_D &= \frac{e^{-i\frac{\pi}{3}P_3(-q_G\gamma_G-(r-2)\gamma_R)}}{\Gamma_e(-q_G\gamma_G-(r-2)\gamma_R; \tau, \omega) \Gamma_e(\omega + q_G\gamma_G + (r-2)\gamma_R; \tau, \omega)} \\ &= e^{-i\frac{\pi}{3}P_3(-q_G\gamma_G-(r-2)\gamma_R)} \Theta(-q_G\gamma_G - (r-2)\gamma_R; \omega), \end{aligned} \quad (4.32)$$

and

$$\begin{aligned} Z_{1\text{-loop}}^{\text{CM}}|_R &= e^{i\frac{\pi}{3}P_3(q_G\gamma_G+r\gamma_R)} \Gamma_e(q_G\gamma_G + r\gamma_R; \tau, \omega) \Gamma_e(\omega - q_G\gamma_G - r\gamma_R; \tau, \omega) \\ &= \frac{e^{i\frac{\pi}{3}P_3(q_G\gamma_G+r\gamma_R)}}{\Theta(q_G\gamma_G + r\gamma_R; \omega)}. \end{aligned} \quad (4.33)$$

With the conventions used above, the unmixed boundary polarizations define the two elementary four-dimensional cylinder blocks used in the applications below. For a generic argument u , we set

$$z_{\text{DD}}(u) := e^{-\frac{i\pi}{3}P_3(-u)}\Theta(-u;\omega), \quad z_{\text{RR}}(u) := \frac{e^{\frac{i\pi}{3}P_3(u)}}{\Theta(u;\omega)}. \quad (4.34)$$

With this normalization the DD and RR blocks are inverse after reflection of the argument:

$$z_{\text{DD}}(u) z_{\text{RR}}(-u) = 1. \quad (4.35)$$

Thus inserting a DD/RR pair with opposite arguments gives an exact inverse pair. By contrast, the same-argument product is a holomorphic non-vanishing factor and introduces no polar hyperplane. We also introduce

$$\Theta'_0 := \left. \frac{\partial}{\partial u} \Theta(u;\omega) \right|_{u=0}, \quad \varphi_0 := \exp\left[\frac{i\pi}{3}P_3(0)\right]. \quad (4.36)$$

The constant φ_0 is the value of the cubic phase of the RR block at $u = 0$; no additional polynomial is being introduced. Thus a simple positive-charge pole of $z_{\text{RR}}(u)$ contributes φ_0/Θ'_0 in the oriented JK convention used below.

The corresponding formula for a general gauge theory is obtained by multiplying these elementary factors over weights. Let G be a compact gauge group with Weyl group W_G , Cartan fugacities $u = (u_1, \dots, u_{\text{rk } G})$, root system Δ_G , and chiral multiplets Φ_I of R-charge r_I , flavor fugacity ν_I , representation $\mathcal{R}_I$, and boundary polarization $\mathbf{q}_I \in \{\text{DD}, \text{DR}, \text{RD}, \text{RR}\}$. With the normalized contour measure used throughout this paper, the localized partition function is

$$\mathcal{Z}_{\text{Cyl} \times T^2}^{G, \{\mathbf{q}_I\}} = \frac{1}{|W_G|} \oint_{\mathcal{C}_\eta} \prod_{a=1}^{\text{rk } G} \frac{du_a}{2\pi i} Z_{\text{vec}}^G(u) \prod_I Z_{\Phi_I}^{\mathbf{q}_I}(u), \quad (4.37)$$

where

$$Z_{\text{vec}}^G(u) = \prod_{\alpha \in \Delta_G} z_{\text{DD}}(\alpha(u)) \quad (4.38)$$

and

$$\begin{aligned} Z_{\Phi_I}^{\text{DD}}(u) &= \prod_{\rho \in \mathcal{R}_I} z_{\text{DD}}(\rho(u) + \nu_I + (r_I - 2)\gamma_R), \\ Z_{\Phi_I}^{\text{RR}}(u) &= \prod_{\rho \in \mathcal{R}_I} z_{\text{RR}}(\rho(u) + \nu_I + r_I\gamma_R), \\ Z_{\Phi_I}^{\text{DR}}(u) &= Z_{\Phi_I}^{\text{RD}}(u) = 1 \end{aligned} \quad (4.39)$$

for the bulk complex with no additional boundary multiplets. The product over ρ is the product over the weights of the representation $\mathcal{R}_I$; flavor fugacities are included by replacing ν_I with the appropriate linear combination of flavor holonomies. The vector factor (4.38) is the non-zero-weight part of the adjoint one-loop determinant. It can be obtained from the chiral determinant by taking the adjoint representation and the effective R-charge $r = 2$; the non-zero adjoint weights are the roots $\alpha \in \Delta_G$, while the Cartan zero weights are cancelled by the ghost/zero-mode sector and leave the normalized Cartan measure in (4.37). With our orientation convention

this gives the DD root factor $z_{\text{DD}}(\alpha(u))$. Using the reflected inverse relation one may equivalently trade it for inverse RR factors with reflected arguments, but the DD presentation is the one in which Vandermonde zeros remove coincident gauge eigenvalues most transparently. Boundary multiplets are not included in (4.39); they multiply the bulk kernel by the flip ratios described below. The notation $\mathcal{C}_\eta$ always denotes a JK cycle on the chosen ω -elliptic fugacity torus. Products of blocks which fail the ω -ellipticity test should be regarded as local one-loop building blocks, not as standalone meromorphic kernels on this torus.

For mixed choices (DR or RD), the acyclicity statement (4.24) implies the absence of admissible kernel/cokernel modes in the bulk complex and hence a trivial elementary bulk block:

$$z_{\text{DR}}(u) = z_{\text{RD}}(u) = 1, \quad (4.40)$$

up to possible finite-dimensional boundary degrees of freedom if included explicitly. Equivalently, for a full chiral multiplet one has $Z_\Phi^{\text{DR}} = Z_\Phi^{\text{RD}} = 1$ in the minimal bulk complex.

The acyclicity statement (4.40) refers to the bulk complex without additional boundary degrees of freedom. As in the disk computation of boundary-condition flips [8, 25, 26], boundary multiplets can change the effective boundary polarization. On the cylinder there are two independent boundary components, hence one can flip the northern and southern boundary conditions separately.

We denote the boundary circles at $\theta = \theta_N$ and $\theta = \theta_S$ by ∂_N and ∂_S , respectively. The first label in a pair such as DR refers to ∂_N , while the second refers to ∂_S . We do not introduce separate symbols for the four bulk determinants: the non-trivial unmixed ones are the blocks z_{RR} and z_{DD} , while the mixed elementary blocks are equal to one by (4.40). A boundary multiplet which flips one end of the cylinder is represented by the ratio of the final determinant to the initial one. These ratios are full boundary determinants, and therefore are written in terms of theta functions rather than elliptic Gamma cap wavefunctions.

Flipping the northern boundary gives four possibilities:

$$\begin{aligned} Z_{\text{RR} \rightarrow \text{DR}}^{\partial_N}(u) &:= \frac{1}{z_{\text{RR}}(u)} = \exp\left[-\frac{i\pi}{3}P_3(u)\right] \Theta(u; \omega), \\ Z_{\text{DR} \rightarrow \text{RR}}^{\partial_N}(u) &:= z_{\text{RR}}(u) = \frac{\exp\left[+\frac{i\pi}{3}P_3(u)\right]}{\Theta(u; \omega)}, \\ Z_{\text{RD} \rightarrow \text{DD}}^{\partial_N}(u) &:= z_{\text{DD}}(u) = \exp\left[-\frac{i\pi}{3}P_3(-u)\right] \Theta(-u; \omega), \\ Z_{\text{DD} \rightarrow \text{RD}}^{\partial_N}(u) &:= \frac{1}{z_{\text{DD}}(u)} = \frac{\exp\left[+\frac{i\pi}{3}P_3(-u)\right]}{\Theta(-u; \omega)}. \end{aligned} \quad (4.41)$$

Flipping the southern boundary gives the analogous four possibilities:

$$\begin{aligned}
Z_{RR \rightarrow RD}^{\partial_S}(u) &:= \frac{1}{z_{RR}(u)} = \exp\left[-\frac{i\pi}{3}P_3(u)\right] \Theta(u; \omega), \\
Z_{RD \rightarrow RR}^{\partial_S}(u) &:= z_{RR}(u) = \frac{\exp\left[+\frac{i\pi}{3}P_3(u)\right]}{\Theta(u; \omega)}, \\
Z_{DR \rightarrow DD}^{\partial_S}(u) &:= z_{DD}(u) = \exp\left[-\frac{i\pi}{3}P_3(-u)\right] \Theta(-u; \omega), \\
Z_{DD \rightarrow DR}^{\partial_S}(u) &:= \frac{1}{z_{DD}(u)} = \frac{\exp\left[+\frac{i\pi}{3}P_3(-u)\right]}{\Theta(-u; \omega)}. \tag{4.42}
\end{aligned}$$

These equations are the cylinder analogue of the boundary-condition flip: multiplying the full cylinder kernel by the appropriate boundary determinant changes one end of the cylinder from Robin to Dirichlet, or conversely. The dependence on the other end is encoded in the initial and final labels, for instance $RR \rightarrow DR$ and $RD \rightarrow DD$ are different northern flips. The flip operation is symmetric under exchanging Dirichlet and Robin: reversing an arrow replaces the determinant by its inverse, and exchanging the two boundary labels maps the mixed blocks z_{DR} and z_{RD} into each other. This symmetry is consistent with the bulk acyclicity of both mixed sectors, but the two flips must still be distinguished because they live on different boundary components. If instead one factorizes the cylinder into two caps, the corresponding one-sided ratios are elliptic-Gamma cap wavefunctions; those are useful for gluing, but they should not be confused with the full theta-function boundary determinants displayed above.

5 Applications and Examples

The cylinder blocks constructed above lead to four-dimensional $\text{Cyl} \times T^2$ residue identities of electric–magnetic type and to two-dimensional cylinder degenerations reproducing the chamber decomposition of GLSMs on $\text{Cyl} \simeq I \times S^1$.

5.1 Dualities

The duality tests in this subsection are modeled on Seiberg-type rank-changing dualities [36], while the contour prescription is the Jeffrey–Kirwan prescription used in equivariant localization, elliptic genera and twisted indices [32–35]. The selected matter content and boundary conditions follow from the anomaly cancellation conditions surveyed in the appendix. In this subsection, ω -ellipticity refers to meromorphicity of the completed gauge integrand on the auxiliary gauge-fugacity torus $u \sim u + 1 \sim u + \omega$. This is the torus relevant for the JK contour prescription, and should not be confused with the physical spacetime torus of modulus τ .

5.1.1 $U(N_c) \leftrightarrow U(N_f - N_c)$

Set

$$n := N_f - N_c \geq 0, \tag{5.1}$$

and

$$M := \sum_{i=1}^{N_f} m_i, \quad \widetilde{M} := \sum_{i=1}^{N_f} \widetilde{m}_i. \quad (5.2)$$

The electric theory has gauge group $U(N_c)$, gauge fugacities u_a , $a = 1, \dots, N_c$, and integrand

$$\begin{aligned} Z_{\text{el}}(u) &= \frac{1}{N_c!} z_{\text{DD}} \left[\sum_{a=1}^{N_c} u_a + M + \lambda \right] z_{\text{DD}} \left[- \sum_{a=1}^{N_c} u_a + \widetilde{M} + \lambda \right] \\ &\times \prod_{\substack{a,b=1 \\ a \neq b}}^{N_c} z_{\text{DD}}[u_a - u_b] \\ &\times \prod_{a=1}^{N_c} \prod_{i=1}^{N_f} z_{\text{RR}}[-u_a - m_i] z_{\text{RR}}[u_a - \widetilde{m}_i] \\ &\times \prod_{a=1}^{N_c} \prod_{\alpha=1}^n z_{\text{DD}}[u_a + b_\alpha] z_{\text{DD}}[-u_a + \widetilde{b}_\alpha]. \end{aligned} \quad (5.3)$$

The corresponding electric partition function is the JK-contour integral of this meromorphic form

$$\mathcal{Z}_{\text{el}}^{U(N_c)} := \oint_{\mathcal{C}_\eta^{\text{el}}} \prod_{a=1}^{N_c} \frac{du_a}{2\pi i} Z_{\text{el}}(u) = \text{JK}_{U(N_c), \eta} Z_{\text{el}}. \quad (5.4)$$

The factor $1/N_c!$ is already included in (5.3). The matter content consists of N_f fundamental RR chirals, N_f anti-fundamental RR chirals, n fundamental DD chirals, n anti-fundamental DD chirals, and two determinant-type DD chirals

$$S_+ : \quad z_{\text{DD}} \left[\sum_a u_a + M + \lambda \right], \quad S_- : \quad z_{\text{DD}} \left[- \sum_a u_a + \widetilde{M} + \lambda \right]. \quad (5.5)$$

Notice the ω -ellipticity of Z_{el} on the gauge fugacity space follows from the quadratic condition $n = N_f - N_c$ and from imposing the linear balancing condition

$$\sum_{\alpha=1}^n b_\alpha = \sum_{\alpha=1}^n \widetilde{b}_\alpha. \quad (5.6)$$

Hence, when studying the possible contributing poles, we omit those given by integer shifts of the ω -lattice. The determinant shifts are not independent masses: they are fixed by the total flavor fugacities in (5.2). In checking the multiplier under a gauge shift $u_a \mapsto u_a + \omega$, the dependence on the fundamental and anti-fundamental masses enters only through the total combinations M and $\widetilde{M}$. These are precisely the combinations appearing in the two determinant shifts. Hence the mass-dependent part of the gauge ω -elliptic multiplier cancels without imposing any balancing condition on the individual m_i or $\widetilde{m}_i$. The individual masses remain, of course, genuine flavor fugacities of the integrand and of the JK residues.

We use the oriented JK prescription with the orientation fixed so that each simple pole of z_{RR} contributes φ_0/Θ'_0 , cf. (4.36). The charge covector of a polar hyperplane is the coefficient of the gauge fugacities in the argument of the corresponding RR block. Thus a factor $z_{\text{RR}}[q \cdot u + m]$ contributes a polar hyperplane with charge covector q . Throughout this subsection we work in the chamber

$$\eta = (1, \dots, 1), \quad (5.7)$$

which selects precisely the RR poles whose arguments have coefficient +1 with respect to the relevant gauge fugacity. On the electric side these are the factors $z_{\text{RR}}[u_a - \tilde{m}_i]$. The electric JK residue is therefore supported at

$$u_a = \tilde{m}_i. \quad (5.8)$$

The vector-multiplet Vandermonde factor removes coincident flavor labels; the contributing poles are indexed by subsets

$$\mathcal{I} \subset \{1, \dots, N_f\}, \quad |\mathcal{I}| = N_c. \quad (5.9)$$

Thus the electric JK residue is a finite sum of $\binom{N_f}{N_c}$ non-vanishing sectors. Equivalently, before quotienting by the Weyl group one may assign ordered flavor labels to the N_c gauge variables; the Vandermonde factor kills coincident labels and the factor $1/N_c!$ removes the ordering. With

$$S_{\mathcal{I}} := \sum_{i \in \mathcal{I}} \tilde{m}_i, \quad (5.10)$$

Before writing the residue formula it is useful to spell out one finite example. Take $N_f = 5$ and $N_c = 2$, so that $n = N_f - N_c = 3$. One electric residue sector is

$$\mathcal{I} = \{2, 5\}, \quad u_1 = \tilde{m}_2, \quad u_2 = \tilde{m}_5. \quad (5.11)$$

Equivalently, after relabelling the two electric gauge variables, this is the pole $u_1 = \tilde{m}_2, u_2 = \tilde{m}_5$. The complementary magnetic sector is

$$\mathcal{J} = \mathcal{I}^c = \{1, 3, 4\}, \quad v_1 = -\tilde{m}_1, \quad v_2 = -\tilde{m}_3, \quad v_3 = -\tilde{m}_4, \quad (5.12)$$

up to Weyl permutations of the magnetic gauge variables. The electric determinant factor depends on $S_{\mathcal{I}} = \tilde{m}_2 + \tilde{m}_5$, while the magnetic determinant factor depends on $\sum_{j \in \mathcal{J}} \tilde{m}_j$. Since $\mathcal{I} \sqcup \mathcal{J} = \{1, \dots, 5\}$, the two expressions are converted into each other by the total anti-fundamental mass $\tilde{M}$. This is the concrete mechanism behind the complementary-residue matching proved below. In general, the residue is

$$\text{JK}_{U(N_c), \eta} Z_{\text{el}} = \left(\frac{\varphi_0}{\Theta'_0} \right)^{N_c} \sum_{\substack{\mathcal{I} \subset \{1, \dots, N_f\} \\ |\mathcal{I}| = N_c}} \mathcal{E}_{\mathcal{I}}, \quad (5.13)$$

where

$$\begin{aligned} \mathcal{E}_{\mathcal{I}} &= z_{\text{DD}}[S_{\mathcal{I}} + M + \lambda] z_{\text{DD}}[-S_{\mathcal{I}} + \tilde{M} + \lambda] \\ &\times \prod_{i \in \mathcal{I}} \prod_{\ell=1}^{N_f} z_{\text{RR}}[-\tilde{m}_i - m_{\ell}] \\ &\times \prod_{i \in \mathcal{I}} \prod_{j \notin \mathcal{I}} z_{\text{RR}}[\tilde{m}_i - \tilde{m}_j] \\ &\times \prod_{i \in \mathcal{I}} \prod_{\alpha=1}^n z_{\text{DD}}[\tilde{m}_i + b_{\alpha}] z_{\text{DD}}[-\tilde{m}_i + \tilde{b}_{\alpha}]. \end{aligned} \quad (5.14)$$

The magnetic theory has gauge group

$$U(n) = U(N_f - N_c), \quad (5.15)$$

with fugacities v_r , $r = 1, \dots, n$. In the chamber (5.7) we use the presentation

$$\begin{aligned}
Z_{\text{mag}}(v) = & \frac{1}{n!} z_{\text{DD}} \left[-\sum_{r=1}^n v_r + \lambda \right] z_{\text{DD}} \left[\sum_{r=1}^n v_r + M + \widetilde{M} + \lambda \right] \\
& \times \prod_{\substack{r,s=1 \\ r \neq s}}^n z_{\text{DD}}[v_r - v_s] \\
& \times \prod_{r=1}^n \prod_{i=1}^{N_f} z_{\text{RR}}[v_r + \widetilde{m}_i] z_{\text{DD}}[-v_r + m_i] \\
& \times \prod_{r=1}^n \prod_{\alpha=1}^n z_{\text{RR}}[v_r - b_\alpha] z_{\text{DD}}[v_r - b_\alpha] z_{\text{RR}}[-v_r + b_\alpha] z_{\text{RR}}[-v_r - \widetilde{b}_\alpha].
\end{aligned} \tag{5.16}$$

The magnetic integrand is written in a completed selected-chamber presentation. It is obtained from the uncompleted ω -elliptic magnetic integrand by inserting the exact inverse pairs $z_{\text{DD}}[v_r - b_\alpha] z_{\text{RR}}[-v_r + b_\alpha] = 1$. Therefore its ω -elliptic multiplier is unchanged, and it is ω -elliptic under the same conditions (5.2) and (5.6) as the electric integrand. The same-argument pair

$$z_{\text{RR}}[v_r - b_\alpha] z_{\text{DD}}[v_r - b_\alpha] \tag{5.17}$$

is holomorphic and non-vanishing, and hence introduces no polar hyperplane. The reflected pair satisfies

$$z_{\text{DD}}[v_r - b_\alpha] z_{\text{RR}}[-v_r + b_\alpha] = 1. \tag{5.18}$$

These identities are used only to specify the polar presentation of the magnetic integrand. The chamber $\eta = (1, \dots, 1)$ selects, for each variable v_r , the RR factors whose arguments have coefficient +1 with respect to v_r . In the magnetic integrand the selected factors are

$$z_{\text{RR}}[v_r + \widetilde{m}_i], \tag{5.19}$$

and they generate the selected polar hyperplanes. No further hyperplanes are selected. The determinant factors, the vector factor, and the $z_{\text{DD}}[-v_r + m_i]$ factors are of DD type and do not produce poles. The same-argument product in (5.17) is regular, while the remaining RR factors in the b_α -sector have arguments $-v_r + b_\alpha$ and $-v_r - \widetilde{b}_\alpha$, hence coefficient -1 with respect to v_r . They belong to the opposite JK chamber. Thus the selected magnetic cell is generated only by the hyperplanes

$$v_r + \widetilde{m}_i = 0. \tag{5.20}$$

The singlet sector is

$$Z_{\text{sing}} = Z_M Z_B, \tag{5.21}$$

where the meson sector is

$$Z_M = \prod_{i,j=1}^{N_f} z_{\text{RR}}[-m_i - \widetilde{m}_j], \tag{5.22}$$

and the boundary singlet sector is

$$Z_B = \prod_{i=1}^{N_f} \prod_{\alpha=1}^n z_{\text{DD}}[\widetilde{m}_i + b_\alpha] z_{\text{DD}}[-\widetilde{m}_i + \widetilde{b}_\alpha]. \tag{5.23}$$

The magnetic gauge integral and the full magnetic partition function are

$$\mathcal{Z}_{\text{mag}}^{U(n)} := \oint_{\mathcal{C}_\eta^{\text{mag}}} \prod_{r=1}^n \frac{dv_r}{2\pi i} Z_{\text{mag}}(v) = \text{JK}_{U(n), \eta} Z_{\text{mag}}, \quad \mathcal{Z}_{\text{mag, full}}^{U(n)} := Z_{\text{sing}} \mathcal{Z}_{\text{mag}}^{U(n)}. \quad (5.24)$$

The factors in $Z_{\text{mag}}(v)$ are cylinder one-loop determinants in the indicated polarizations. The exact reflected DD/RR pairs inserted in (5.16) are not additional boundary multiplets; they are a change of polar presentation equal to one. By contrast, the gauge-neutral sector Z_B can be interpreted as a boundary singlet sector implementing the required flip/cancellation of the spectator polarizations. The same-argument product (5.17) is a holomorphic contact factor and should also not be counted as a pole-carrying boundary degree of freedom. Genuine boundary multiplets enter through the flip ratios (4.41)–(4.42).

Lemma 5.1 (Selected JK poles and multiplicities). *Under the genericity assumptions of Theorem 5.4, the chamber $\eta = (1, \dots, 1)$ selects the following isolated poles, with unit multiplicity after the Weyl factors $1/N_c!$ and $1/n!$ are included. On the electric side the selected poles are exactly*

$$u_a = \tilde{m}_i, \quad i \in \mathcal{I}, \quad |\mathcal{I}| = N_c, \quad (5.25)$$

with distinct flavor labels. On the magnetic side the selected poles are exactly

$$v_r = -\tilde{m}_j, \quad j \in \mathcal{J}, \quad |\mathcal{J}| = n, \quad (5.26)$$

again with distinct labels. Thus the magnetic JK residue is a finite sum of $\binom{N_f}{n} = \binom{N_f}{N_f - N_c} = \binom{N_f}{N_c}$ sectors. The complement map $\mathcal{I} \mapsto \mathcal{J} = \mathcal{I}^c$ is therefore a bijection between the electric and magnetic residue sectors. The magnetic b_α -hyperplanes and the same-argument contact pairs do not contribute in this chamber. At the magnetic pole labelled by $\mathcal{J}$, the vector factor cancels the selected RR factors with both flavor labels in $\mathcal{J}$; the only off-diagonal RR factors left after this cancellation are those with one label in $\mathcal{J}$ and the other in $\mathcal{J}^c$.

Proof. A polar hyperplane is selected by the JK covector only if it is produced by an RR block whose argument has positive coefficient with respect to the corresponding gauge fugacity. For the electric integrand (5.3), the only such factors are $z_{\text{RR}}[u_a - \tilde{m}_i]$. The local charge matrix at a pole with one such hyperplane for each a is a permutation matrix and hence has determinant ± 1 , with the sign absorbed by the oriented JK convention. The Weyl factor $1/N_c!$ removes the ordering of the variables. If two variables are assigned the same flavor label, the vector factor contains $z_{\text{DD}}[0]$, so the corresponding would-be residue vanishes. Thus only subsets $\mathcal{I}$ of cardinality N_c remain, each once. For distinct labels, the RR factors whose flavor label is also selected are paired with the ordered vector factors by the reflected inverse relation, leaving precisely the factors with labels outside $\mathcal{I}$, as in (5.14).

For the magnetic integrand (5.16), the same charge test selects $z_{\text{RR}}[v_r + \tilde{m}_i]$ and gives (5.26). The determinant factors, the vector factor and all DD factors are holomorphic at the selected hyperplanes and therefore do not generate JK charges. The RR factors $z_{\text{RR}}[-v_r + b_\alpha]$ and $z_{\text{RR}}[-v_r - \tilde{b}_\alpha]$ have charge -1 with respect to v_r , hence lie in the opposite chamber;

they can contribute only after crossing a wall, which is excluded by the genericity assumption. The product $z_{\text{RR}}[v_r - b_\alpha]z_{\text{DD}}[v_r - b_\alpha]$ is holomorphic and non-vanishing, so it defines no polar hyperplane. As on the electric side, the local charge matrix is a permutation matrix, the Weyl factor removes the ordering, and coincident labels are killed by the Vandermonde zero. For distinct labels, the vector factor cancels the RR factors with both labels in $\mathcal{J}$, leaving only the mixed factors with labels in $\mathcal{J}$ and $\mathcal{J}^c$. This proves both the list of selected poles and the unit multiplicity statement. $\square$

Lemma 5.2 (Theta-only complementary residues). *The algebraic core of the determinant-pair duality is already visible after stripping the Bernoulli contact phases and the universal simple-pole factors. In this paragraph write $\Theta[x] \equiv \Theta(x; \omega)$. For a selected electric subset $\mathcal{I}$ and complementary magnetic subset $\mathcal{J} = \mathcal{I}^c$, define the normalized theta residues*

$$\begin{aligned} \widehat{\mathcal{E}}_{\mathcal{I}} &= \Theta[-S_{\mathcal{I}} - M - \lambda] \Theta[S_{\mathcal{I}} - \widetilde{M} - \lambda] \\ &\times \prod_{i \in \mathcal{I}} \prod_{\ell=1}^{N_f} \frac{1}{\Theta[-\widetilde{m}_i - m_\ell]} \prod_{i \in \mathcal{I}} \prod_{j \in \mathcal{J}} \frac{1}{\Theta[\widetilde{m}_i - \widetilde{m}_j]} \\ &\times \prod_{i \in \mathcal{I}} \prod_{\alpha=1}^n \Theta[-\widetilde{m}_i - b_\alpha] \Theta[\widetilde{m}_i - \widetilde{b}_\alpha], \end{aligned} \quad (5.27)$$

and

$$\begin{aligned} \widehat{\mathcal{M}}_{\mathcal{J}} &= \Theta[-S_{\mathcal{J}} - \lambda] \Theta[S_{\mathcal{J}} - M - \widetilde{M} - \lambda] \\ &\times \prod_{j \in \mathcal{J}} \prod_{\ell=1}^{N_f} \Theta[-\widetilde{m}_j - m_\ell] \prod_{j \in \mathcal{J}} \prod_{i \in \mathcal{I}} \frac{1}{\Theta[\widetilde{m}_i - \widetilde{m}_j]} \\ &\times \prod_{j \in \mathcal{J}} \prod_{\alpha=1}^n \frac{1}{\Theta[-\widetilde{m}_j - b_\alpha] \Theta[\widetilde{m}_j - \widetilde{b}_\alpha]}. \end{aligned} \quad (5.28)$$

Let

$$\widehat{Z}_{\text{sing}} = \prod_{r=1}^{N_f} \prod_{\ell=1}^{N_f} \frac{1}{\Theta[-\widetilde{m}_r - m_\ell]} \prod_{r=1}^{N_f} \prod_{\alpha=1}^n \Theta[-\widetilde{m}_r - b_\alpha] \Theta[\widetilde{m}_r - \widetilde{b}_\alpha]. \quad (5.29)$$

Then the residue matching is sectorwise:

$$\widehat{\mathcal{E}}_{\mathcal{I}} = \widehat{Z}_{\text{sing}} \widehat{\mathcal{M}}_{\mathcal{J}}, \quad \mathcal{J} = \mathcal{I}^c. \quad (5.30)$$

Proof. The determinant factors agree because $S_{\mathcal{I}} + S_{\mathcal{J}} = \widetilde{M}$:

$$-S_{\mathcal{J}} - \lambda = S_{\mathcal{I}} - \widetilde{M} - \lambda, \quad S_{\mathcal{J}} - M - \widetilde{M} - \lambda = -S_{\mathcal{I}} - M - \lambda. \quad (5.31)$$

The off-diagonal factors are also identical after reordering the finite product $\prod_{j \in \mathcal{J}} \prod_{i \in \mathcal{I}} = \prod_{i \in \mathcal{I}} \prod_{j \in \mathcal{J}}$. It remains to check the meson and spectator factors. For each flavor label set

$$X_r := \left(\prod_{\ell=1}^{N_f} \frac{1}{\Theta[-\widetilde{m}_r - m_\ell]} \right) \left(\prod_{\alpha=1}^n \Theta[-\widetilde{m}_r - b_\alpha] \Theta[\widetilde{m}_r - \widetilde{b}_\alpha] \right). \quad (5.32)$$

The theta-only singlet factor is $\widehat{Z}_{\text{sing}} = \prod_{r=1}^{N_f} X_r$, while the meson/spectator part of the magnetic residue contributes $\prod_{j \in \mathcal{J}} X_j^{-1}$. Hence

$$\widehat{Z}_{\text{sing}} \prod_{j \in \mathcal{J}} X_j^{-1} = \left(\prod_{r=1}^{N_f} X_r \right) \left(\prod_{j \in \mathcal{J}} X_j^{-1} \right) = \prod_{i \in \mathcal{I}} X_i, \quad (5.33)$$

which is exactly the meson/spectator part of $\widehat{\mathcal{E}}_{\mathcal{I}}$. This proves the sector identity. $\square$

Remark 5.3 (Theta-only versus elliptic parent integrand). Lemma 5.2 shows that the complementary residue identity itself does not use the spectator balancing condition (5.6); the spectator fugacities enter only through the flavor-factorized variables X_r . The complementary-residue identity is therefore independent of the elliptic properties of Θ : it is a finite algebraic identity. The balancing condition is needed earlier for the parent Bernoulli-decorated cylinder integrand to be single-valued on the auxiliary gauge-fugacity torus and hence to define the JK contour problem in the ω -elliptic presentation used here.

Theorem 5.4 (Determinant-pair cylinder duality). *Let $N_f > N_c$, set $n = N_f - N_c$, and consider the electric and magnetic meromorphic forms Z_{el} and Z_{mag} defined in (5.3) and (5.16), together with the singlet factor $Z_{\text{sing}} = Z_M Z_B$ in (5.22)–(5.23). Assume the determinant shifts are fixed by (5.2), the spectator fugacities obey (5.6), and all flavor and spectator parameters are generic: the selected JK intersections are simple, no non-selected zero or pole collides with them, no resonance hyperplane of the mixed boundary problem passes through them, and no pole lies on the boundary of the chosen ω -elliptic fundamental domain. In the chamber $\eta = (1, \dots, 1)$, with the pole normalization fixed by (4.36), one has*

$$\mathcal{Z}_{\text{el}}^{U(N_c)} = \left(\frac{\varphi_0}{\Theta_0} \right)^{N_c - n} \mathcal{Z}_{\text{mag,full}}^{U(n)}. \quad (5.34)$$

Equivalently, after removing the universal τ, ω -dependent normalization of simple poles,

$$\mathcal{Z}_{\text{el}}^{U(N_c)} \doteq \mathcal{Z}_{\text{mag,full}}^{U(N_f - N_c)}. \quad (5.35)$$

Proof. By Lemma 5.1, the selected magnetic poles are indexed by subsets

$$\mathcal{J} \subset \{1, \dots, N_f\}, \quad |\mathcal{J}| = n, \quad (5.36)$$

with unit multiplicity in the normalized JK measure. We set

$$\mathcal{I} = \mathcal{J}^c, \quad |\mathcal{I}| = N_c. \quad (5.37)$$

At $v_r = -\tilde{m}_j$, $j \in \mathcal{J}$, the determinant-type factors give

$$z_{\text{DD}} \left[-\sum_r v_r + \lambda \right] \longrightarrow z_{\text{DD}} \left[\lambda + \sum_{j \in \mathcal{J}} \tilde{m}_j \right] = z_{\text{DD}} [-S_{\mathcal{I}} + \widetilde{M} + \lambda], \quad (5.38)$$

and

$$z_{\text{DD}} \left[\sum_r v_r + M + \widetilde{M} + \lambda \right] \longrightarrow z_{\text{DD}} \left[\lambda + M + \widetilde{M} - \sum_{j \in \mathcal{J}} \tilde{m}_j \right] = z_{\text{DD}} [S_{\mathcal{I}} + M + \lambda]. \quad (5.39)$$

For $j \in \mathcal{J}$, the meson factors cancel the magnetic DD chirals:

$$z_{\text{DD}}[-v_r + m_\ell] \big|_{v_r = -\tilde{m}_j} = z_{\text{DD}}[\tilde{m}_j + m_\ell], \quad (5.40)$$

and hence

$$z_{\text{RR}}[-m_\ell - \tilde{m}_j] z_{\text{DD}}[\tilde{m}_j + m_\ell] = 1, \quad j \in \mathcal{J}. \quad (5.41)$$

The remaining meson factors are

$$\prod_{i \in \mathcal{I}} \prod_{\ell=1}^{N_f} z_{\text{RR}}[-\tilde{m}_i - m_\ell]. \quad (5.42)$$

Similarly,

$$z_{\text{RR}}[-v_r - \tilde{b}_\alpha] \big|_{v_r = -\tilde{m}_j} = z_{\text{RR}}[\tilde{m}_j - \tilde{b}_\alpha], \quad (5.43)$$

and

$$z_{\text{DD}}[-\tilde{m}_j + \tilde{b}_\alpha] z_{\text{RR}}[\tilde{m}_j - \tilde{b}_\alpha] = 1, \quad j \in \mathcal{J}. \quad (5.44)$$

For the b_α -sector set

$$y_{j\alpha} := \tilde{m}_j + b_\alpha. \quad (5.45)$$

At $v_r = -\tilde{m}_j$, the magnetic factors are

$$z_{\text{RR}}[v_r - b_\alpha] z_{\text{DD}}[v_r - b_\alpha] z_{\text{RR}}[-v_r + b_\alpha] = z_{\text{RR}}[-y_{j\alpha}] z_{\text{DD}}[-y_{j\alpha}] z_{\text{RR}}[y_{j\alpha}]. \quad (5.46)$$

Using the reflected inverse relation $z_{\text{DD}}(x) z_{\text{RR}}(-x) = 1$,

$$z_{\text{DD}}[-y_{j\alpha}] z_{\text{RR}}[y_{j\alpha}] = 1, \quad (5.47)$$

and

$$z_{\text{DD}}[y_{j\alpha}] z_{\text{RR}}[-y_{j\alpha}] = 1. \quad (5.48)$$

After multiplication by Z_B , the surviving b_α -sector is

$$\prod_{i \in \mathcal{I}} \prod_{\alpha=1}^n z_{\text{DD}}[\tilde{m}_i + b_\alpha] z_{\text{DD}}[-\tilde{m}_i + \tilde{b}_\alpha]. \quad (5.49)$$

By Lemma 5.1, the magnetic vector multiplet cancels the $i, j \in \mathcal{J}$, $i \neq j$, RR terms. The remaining magnetic RR chirals give

$$z_{\text{RR}}[v_r + \tilde{m}_i] \longrightarrow z_{\text{RR}}[\tilde{m}_i - \tilde{m}_j]. \quad (5.50)$$

The remaining factor is

$$\prod_{i \in \mathcal{I}} \prod_{j \in \mathcal{J}} z_{\text{RR}}[\tilde{m}_i - \tilde{m}_j], \quad (5.51)$$

Combining (5.42), (5.49), and (5.51) gives

$$Z_{\text{sing}} \mathcal{M}_{\mathcal{J}} = \mathcal{E}_{\mathcal{J}^c}, \quad (5.52)$$

where $\mathcal{M}_{\mathcal{J}}$ denotes the magnetic residue of (5.16). Summing over $\mathcal{J}$ gives

$$\mathrm{JK}_{U(N_c),\eta} Z_{\mathrm{el}} = \left(\frac{\varphi_0}{\Theta_0} \right)^{N_c-n} Z_{\mathrm{sing}} \mathrm{JK}_{U(n),\eta} Z_{\mathrm{mag}}. \quad (5.53)$$

Equivalently, up to an overall normalization depending only on τ and ω ,

$$\mathrm{JK}_{U(N_c),\eta} Z_{\mathrm{el}} \doteq Z_{\mathrm{sing}} \mathrm{JK}_{U(N_f-N_c),\eta} Z_{\mathrm{mag}}. \quad (5.54)$$

This is (5.34) written in the shorthand JK notation. $\square$

For $n = 0$, equivalently $N_f = N_c$, the magnetic gauge group is $U(0)$. The rank-changing identity then becomes a finite-dimensional theta-function evaluation. We write this endpoint explicitly, without using the shorthand cylinder blocks, after the SU projection in Subsection 5.1.3.

For $n > 0$, the dual is the magnetic $U(N_f - N_c)$ theory with singlet sector $Z_M Z_B$, evaluated in the JK presentation (5.16).

5.1.2 $SU(N_c) \leftrightarrow SU(N_f - N_c)$ projection

The projection from $U(N_c)$ and $U(n)$ to $SU(N_c)$ and $SU(n)$ is implemented by inserting periodic Dirac delta factors imposing the vanishing of the corresponding center fugacity. On the electric side one inserts

$$\delta_T \left(\sum_{a=1}^{N_c} u_a \right) \quad (5.55)$$

inside the electric integral, and on the magnetic side

$$\delta_T \left(\sum_{r=1}^n v_r \right) \quad (5.56)$$

inside the magnetic integral. These delta factors introduce no additional meromorphic poles and are evaluated on the JK poles. Equivalently, the projected partition functions are

$$\begin{aligned} \mathcal{Z}_{\mathrm{el}}^{SU(N_c)} &:= \oint_{\mathcal{C}_{\eta}^{\mathrm{el}}} \prod_{a=1}^{N_c} \frac{du_a}{2\pi i} \delta_T \left(\sum_{a=1}^{N_c} u_a \right) Z_{\mathrm{el}}(u), \\ \mathcal{Z}_{\mathrm{mag,full}}^{SU(n)} &:= Z_{\mathrm{sing}} \oint_{\mathcal{C}_{\eta}^{\mathrm{mag}}} \prod_{r=1}^n \frac{dv_r}{2\pi i} \delta_T \left(\sum_{r=1}^n v_r \right) Z_{\mathrm{mag}}(v). \end{aligned} \quad (5.57)$$

In the chamber (5.7) the electric delta gives

$$\delta_T(S_{\mathcal{I}}) = \delta_T \left(\sum_{i \in \mathcal{I}} \tilde{m}_i \right), \quad (5.58)$$

while the magnetic delta gives

$$\delta_T \left(- \sum_{j \in \mathcal{J}} \tilde{m}_j \right). \quad (5.59)$$

Since $\mathcal{I} = \mathcal{J}^c$, the two constraints agree on the flavor $SU(N_f)$ slice

$$\widetilde{M} = \sum_{i=1}^{N_f} \tilde{m}_i = 0 \quad (5.60)$$

modulo the period lattice of the torus. Thus the SU projection is compatible with the complementarity $\mathcal{I} = \mathcal{J}^c$ on the traceless flavor slice.

Equivalently, one may solve the SU constraint explicitly before applying the JK prescription.

5.1.3 Theta-function confining identity

Proposition 5.5 (Theta-function confining endpoint). *The endpoint of the rank-changing identity with magnetic gauge group $U(0)$ gives a pure theta-function evaluation. Set*

$$N_f = N_c =: N, \quad M = \sum_{i=1}^N m_i, \quad \widetilde{M} = \sum_{i=1}^N \widetilde{m}_i. \quad (5.61)$$

There are no $b_\alpha, \widetilde{b}_\alpha$ fugacities in this endpoint. With the mass identification (5.61), the Bernoulli phase in the block expression is independent of the integration variables. The remaining constant is cancelled by the standard normalization of the selected simple poles. The result can therefore be written as the following identity involving theta functions only, in the same special-function family as classical theta and elliptic hypergeometric evaluations [19, 20]:

$$\begin{aligned} & \frac{1}{N!} \oint_{\mathcal{C}_\eta^{(N)}} \prod_{a=1}^N \frac{du_a}{2\pi i} \frac{\Theta\left(-\sum_{a=1}^N u_a - M - \lambda; \omega\right) \Theta\left(\sum_{a=1}^N u_a - \widetilde{M} - \lambda; \omega\right)}{\prod_{a=1}^N \prod_{i=1}^N \Theta(-u_a - m_i; \omega) \Theta(u_a - \widetilde{m}_i; \omega)} \prod_{\substack{a,b=1 \\ a \neq b}}^N \Theta(-u_a + u_b; \omega) \\ &= \frac{1}{(\Theta'_0)^N} \frac{\Theta(-\lambda; \omega) \Theta(-\lambda - M - \widetilde{M}; \omega)}{\prod_{i,j=1}^N \Theta(-m_i - \widetilde{m}_j; \omega)}. \end{aligned} \quad (5.62)$$

Here $\mathcal{C}_\eta^{(N)}$ is the JK contour in the chamber

$$\eta = (1, \dots, 1), \quad (5.63)$$

selecting the poles

$$u_a = \widetilde{m}_{i_a}, \quad \{i_1, \dots, i_N\} = \{1, \dots, N\}, \quad (5.64)$$

while the Vandermonde theta factor eliminates coincident labels. Equation (5.62) is the confining endpoint of the $U(N_c) \leftrightarrow U(N_f - N_c)$ residue identity.

Proof. This is the $n = 0$ specialization of Theorem 5.4. The magnetic gauge integral is empty, the spectator sectors disappear, and the remaining electric residues are labelled by permutations of the N anti-fundamental poles. The Vandermonde theta factor removes coincident labels, while the selected-pole normalization cancels the residual Bernoulli phase, leaving precisely the theta-function evaluation (5.62). $\square$

5.2 Complete intersections on the cylinder

For the GLSM of a complete intersection Calabi–Yau in projective space, the relevant one-loop building block is the regularized determinant

$$z_{\text{RR}}^{2d}(u) = -\frac{e^{\pi i u}}{1 - e^{2\pi i u}} = \frac{1}{2i \sin(\pi u)}. \quad (5.65)$$

The Dirichlet block is normalized with the reflected inverse convention inherited from the four-dimensional blocks:

$$z_{\text{DD}}^{2d}(u) = e^{-\pi i u} (1 - e^{2\pi i u}) = -2i \sin(\pi u). \quad (5.66)$$

Thus

$$z_{\text{DD}}^{2d}(u)z_{\text{RR}}^{2d}(-u) = 1, \quad z_{\text{DD}}^{2d}(u)z_{\text{RR}}^{2d}(u) = -1. \quad (5.67)$$

The elementary monodromies and reflection properties are

$$z_{\text{RR}}^{2d}(u+1) = -z_{\text{RR}}^{2d}(u), \quad z_{\text{DD}}^{2d}(u+1) = -z_{\text{DD}}^{2d}(u), \quad (5.68)$$

and

$$z_{\text{RR}}^{2d}(-u) = -z_{\text{RR}}^{2d}(u), \quad z_{\text{DD}}^{2d}(-u) = -z_{\text{DD}}^{2d}(u). \quad (5.69)$$

The block z_{RR}^{2d} has simple poles at $u \in \mathbb{Z}$, with leading behavior

$$z_{\text{RR}}^{2d}(u) = \frac{1}{2\pi i u} + O(1) \quad \text{as } u \rightarrow 0. \quad (5.70)$$

More generally, for an integer charge $Q \in \mathbb{Z}_{\neq 0}$ and a shift $c \in \mathbb{C}$, the function $z_{\text{RR}}^{2d}(Qu + c)$ has simple poles at the $|Q|$ points

$$u_{*,\ell} = \frac{\ell - c}{Q}, \quad \ell = 0, \dots, |Q| - 1, \quad (5.71)$$

with ordinary residue

$$\text{Res}_{u=u_{*,\ell}} z_{\text{RR}}^{2d}(Qu + c) = \frac{(-1)^\ell}{2\pi i Q}. \quad (5.72)$$

The sign $(-1)^\ell$ reflects the monodromy of z_{RR}^{2d} under integer shifts of its argument. Throughout this subsection we absorb the factor $1/(2\pi i)$ in the contour measure, so that the residue contribution of this pole is

$$\frac{(-1)^\ell}{Q}. \quad (5.73)$$

We use the oriented rank-one JK convention

$$\text{JK}_+ = \sum_{\text{positive-charge poles}} \text{Res}, \quad \text{JK}_- = - \sum_{\text{negative-charge poles}} \text{Res}. \quad (5.74)$$

With this convention, the two chambers compute the same partition function whenever the residue theorem applies:

$$\text{JK}_+ Z = \text{JK}_- Z. \quad (5.75)$$

We consider a $U(1)$ GLSM with N chiral multiplets X_i , $i = 1, \dots, N$, of gauge charge $+1$, and r chiral multiplets P_A , $A = 1, \dots, r$, of gauge charge $-\mathbf{n}_A$, with

$$\mathbf{n}_A \in \mathbb{Z}_{>0}. \quad (5.76)$$

The superpotential is

$$W = \sum_{A=1}^r P_A G_A(X_1, \dots, X_N), \quad (5.77)$$

where each G_A is homogeneous of degree $\mathbf{n}_A$. The Calabi–Yau condition is

$$\sum_{A=1}^r \mathbf{n}_A = N. \quad (5.78)$$

Geometrically, the positive JK phase realizes the complete intersection

$$X_{\mathbf{n}_1, \dots, \mathbf{n}_r} = \{G_1 = \dots = G_r = 0\} \subset \mathbb{CP}^{N-1}. \quad (5.79)$$

Denoting by u the $U(1)$ gauge fugacity and by $x_i, \mathbf{p}_A$ the flavor fugacities of X_i, P_A , the cylinder one-loop integrand is

$$Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(u) = \prod_{A=1}^r z_{\text{RR}}^{2d}[-\mathbf{n}_A u + \mathbf{p}_A] \prod_{i=1}^N z_{\text{RR}}^{2d}[u + x_i]. \quad (5.80)$$

Under $u \mapsto u + 1$, the full integrand transforms as

$$Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(u + 1) = (-1)^{N + \sum_{A=1}^r \mathbf{n}_A} Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(u). \quad (5.81)$$

The Calabi–Yau condition (5.78) gives

$$N + \sum_{A=1}^r \mathbf{n}_A = 2N, \quad (5.82)$$

and therefore

$$Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(u + 1) = Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(u). \quad (5.83)$$

Equivalently, in the multiplicative variable

$$z = e^{2\pi i u}, \quad (5.84)$$

the Calabi–Yau condition cancels the net gauge-dependent half-power of z . The meromorphic one-form

$$\frac{dz}{2\pi i z} Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(z) \quad (5.85)$$

has no extra residue at $z = 0$ or at $z = \infty$. This is the multiplicative form of the same single-valuedness condition.

The superpotential constrains the flavor fugacities. If the monomial

$$P_A X_1^{k_1} \dots X_N^{k_N}, \quad \sum_{i=1}^N k_i = \mathbf{n}_A, \quad (5.86)$$

appears in $P_A G_A$, flavor neutrality requires

$$\mathbf{p}_A + \sum_{i=1}^N k_i x_i = 0. \quad (5.87)$$

For a generic polynomial G_A , this fixes most flavor symmetries. In residue computations it is convenient to keep the x_i distinct as an equivariant regulator and impose the desired superpotential constraints only after the residues have been evaluated.

The positive chamber receives contributions from the poles of the positively charged fields X_i , namely

$$u_i^+ = -x_i, \quad i = 1, \dots, N. \quad (5.88)$$

Assuming the x_i are distinct, these poles are simple. Each pole satisfies $u_i^+ + x_i = 0$, so there is no additional monodromy sign. The positive-chamber JK residue is

$$\text{JK}_+ Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d} = \sum_{i=1}^N \left[\prod_{A=1}^r z_{\text{RR}}^{2d} [\mathbf{n}_A x_i + \mathbf{p}_A] \right] \prod_{\substack{j=1 \\ j \neq i}}^N z_{\text{RR}}^{2d} [x_j - x_i]. \quad (5.89)$$

This is the cylinder expression for the geometric phase, localized at the poles of the fields X_i parametrizing the ambient projective space $\mathbb{CP}^{N-1}$.

The negative chamber receives contributions from the poles of the negatively charged fields P_A . For fixed A , the pole equation is

$$-\mathbf{n}_A u + \mathbf{p}_A \in \mathbb{Z}. \quad (5.90)$$

Modulo the periodicity $u \sim u + 1$, this gives $\mathbf{n}_A$ solutions,

$$u_{A,\ell}^- = \frac{\mathbf{p}_A - \ell}{\mathbf{n}_A}, \quad \ell = 0, \dots, \mathbf{n}_A - 1. \quad (5.91)$$

At such a pole, the local residue contribution of $z_{\text{RR}}^{2d} [-\mathbf{n}_A u + \mathbf{p}_A]$ is

$$-\frac{(-1)^\ell}{\mathbf{n}_A}. \quad (5.92)$$

The oriented negative-chamber prescription (5.74) includes an additional minus sign, yielding a net contribution $(+1)(-1)^\ell / \mathbf{n}_A$. Hence

$$\begin{aligned} \text{JK}_- Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d} &= \sum_{A=1}^r \frac{1}{\mathbf{n}_A} \sum_{\ell=0}^{\mathbf{n}_A-1} (-1)^\ell \left[\prod_{i=1}^N z_{\text{RR}}^{2d} \left[x_i + \frac{\mathbf{p}_A - \ell}{\mathbf{n}_A} \right] \right] \\ &\quad \times \left[\prod_{\substack{B=1 \\ B \neq A}}^r z_{\text{RR}}^{2d} \left[\mathbf{p}_B - \frac{\mathbf{n}_B}{\mathbf{n}_A} (\mathbf{p}_A - \ell) \right] \right]. \end{aligned} \quad (5.93)$$

The signs $(-1)^\ell$ are the one-cycle cylinder analogue of the orbifold projection phases that appear in Landau–Ginzburg orbifold expansions of GLSM partition functions [37].

Proposition 5.6 (Two-dimensional chamber equality). *Assume that the two-dimensional complete intersection cylinder integrand is single-valued on the gauge circle. In the complete-intersection family (5.80), this follows from the Calabi–Yau relation (5.78); equivalently, it is the two-dimensional degeneration of the cancellation of the gauge-dependent Bernoulli anomaly, schematically $\sum_s \varepsilon_s q_s^2 = 0$, together with the corresponding linear phase condition in the higher-dimensional block. Assume also that all poles are simple and away from the contour at infinity. Then the positive and negative oriented JK chambers compute the same partition function,*

$$\text{JK}_+ Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d} = \text{JK}_- Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}. \quad (5.94)$$

Proof. Single-valuedness, ensured in the examples above by (5.78), makes the integrand a meromorphic one-form on the gauge circle. The residue theorem gives

$$\sum_{\text{positive-charge poles}} \text{Res} + \sum_{\text{negative-charge poles}} \text{Res} = 0. \quad (5.95)$$

The oriented JK convention (5.74) includes the compensating sign for the negative chamber. Therefore the equality of ordinary residues is equivalent to (5.94). $\square$

The formulas above reproduce explicitly examples such as the cubic curve in $\mathbb{CP}^2$, the quintic threefold, and the intersection of two quadrics in $\mathbb{CP}^3$. In these examples the equality of chambers requires the signs $(-1)^\ell$ in (5.93).

For $r = 1$, one obtains a hypersurface of degree $\mathbf{n} = N$ in $\mathbb{CP}^{N-1} = \mathbb{CP}^{\mathbf{n}-1}$. The integrand reduces to

$$Z_{\mathbf{n}}^{2d}(u) = z_{\text{RR}}^{2d}[-\mathbf{n}u + \mathbf{p}] \prod_{i=1}^{\mathbf{n}} z_{\text{RR}}^{2d}[u + x_i]. \quad (5.96)$$

Imposing, for instance, the flavor constraint associated with the monomial $PX_1 \cdots X_{\mathbf{n}}$,

$$\mathbf{p} + \sum_{i=1}^{\mathbf{n}} x_i = 0, \quad (5.97)$$

the positive-chamber residue becomes

$$\text{JK}_+ Z_{\mathbf{n}}^{2d} = \sum_{i=1}^{\mathbf{n}} z_{\text{RR}}^{2d} \left[\mathbf{n}x_i - \sum_{k=1}^{\mathbf{n}} x_k \right] \prod_{\substack{j=1 \\ j \neq i}}^{\mathbf{n}} z_{\text{RR}}^{2d}[x_j - x_i], \quad (5.98)$$

while the oriented negative-chamber residue is

$$\text{JK}_- Z_{\mathbf{n}}^{2d} = \frac{1}{\mathbf{n}} \sum_{\ell=0}^{\mathbf{n}-1} (-1)^\ell \prod_{i=1}^{\mathbf{n}} z_{\text{RR}}^{2d} \left[x_i - \frac{1}{\mathbf{n}} \sum_{k=1}^{\mathbf{n}} x_k - \frac{\ell}{\mathbf{n}} \right]. \quad (5.99)$$

For $\mathbf{n} = 5$, this is the cylinder counterpart of the quintic GLSM. The positive chamber computes the cylinder partition function in the geometric phase of the quintic threefold

$$X_5 \subset \mathbb{CP}^4, \quad (5.100)$$

localized on the five torus fixed points of $\mathbb{CP}^4$. The negative chamber computes the same cylinder partition function in the Landau–Ginzburg orbifold phase, with orbifold group $\mathbb{Z}_{\mathbf{n}}$ and $\mathbf{n}$ twisted sectors labelled by $\ell = 0, \dots, \mathbf{n} - 1$. Since the cylinder has only one periodic holonomy direction, the orbifold sum has $\mathbf{n}$ sectors rather than the $\mathbf{n}^2$ elliptic sectors that arise on T^2 in the elliptic genus.

For $r > 1$, the negative chamber is naturally interpreted as a hybrid phase. In the negative JK chamber the fields P_A cannot all vanish, and after quotienting by the $U(1)$ gauge symmetry they parametrize the weighted projective stack

$$\mathbb{WCP}_{[\mathbf{n}_1, \dots, \mathbf{n}_r]}^{r-1} := [(\mathbb{C}^r \setminus \{0\})/\mathbb{C}^*], \quad (5.101)$$

where $\mathbb{C}^*$ acts with weights $(\mathbf{n}_1, \dots, \mathbf{n}_r)$. The sign of the gauge charges does not affect the resulting stack, since one may invert the $\mathbb{C}^*$ parameter. The integers $\mathbf{n}_A$ need not be pairwise coprime. If

$$\gcd(\mathbf{n}_1, \dots, \mathbf{n}_r) = g > 1, \quad (5.102)$$

then the stack has a generic stabilizer $\mathbb{Z}_g$. A stronger condition, often called well-formedness, requires

$$\gcd(\mathbf{n}_1, \dots, \widehat{\mathbf{n}}_A, \dots, \mathbf{n}_r) = 1 \quad (5.103)$$

for every A . This removes generic codimension-one stabilizers, but is not needed for the GLSM nor for the residue computation. Thus the negative chamber is a hybrid Landau–Ginzburg phase fibered over $\mathbb{WCP}_{[\mathbf{n}_1, \dots, \mathbf{n}_r]}^{r-1}$, with Landau–Ginzburg fiber given by the fields X_i and superpotential $W = \sum_A P_A G_A(X)$. For $r = 1$, the base $\mathbb{WCP}_{[\mathbf{n}]}^0$ is a stacky point with stabilizer $\mathbb{Z}_{\mathbf{n}}$, recovering the familiar Landau–Ginzburg orbifold phase.

The cylinder determinant z_{RR}^{2d} gives a direct $\text{Cyl} \simeq I \times S^1$ analogue of the GLSM phase structure, compatible with the exact boundary framework for two-dimensional 2, 2 gauge theories [1]. The equality (5.94) provides a stringent consistency check of the cylinder partition function. This two-dimensional construction should be viewed as the Cyl counterpart of the four-dimensional $\text{Cyl} \times T^2$ one-loop blocks discussed above. Starting from the full elliptic-gamma representation, the first degeneration from $\text{Cyl} \times T^2$ to $\text{Cyl} \times S^1$ is

$$\text{Im } \tau \rightarrow +\infty, \quad p = e^{2\pi i \tau} \rightarrow 0. \quad (5.104)$$

The remaining theta block has elliptic nome

$$q = e^{2\pi i \omega}. \quad (5.105)$$

Its cylinder degeneration is

$$\text{Im } \omega \rightarrow +\infty, \quad q = e^{2\pi i \omega} \rightarrow 0. \quad (5.106)$$

In this limit

$$\Theta(u; \omega) = (e^{2\pi i u}; q)_\infty (q e^{-2\pi i u}; q)_\infty \longrightarrow 1 - e^{2\pi i u}. \quad (5.107)$$

Thus the four-dimensional one-loop block reduces, up to Bernoulli/contact terms, to the two-dimensional block $z_{\text{RR}}^{2d}(u)$. The chamber equality (5.94) is the two-dimensional shadow of the electric–magnetic residue identities discussed in Section 5.1.

5.3 Factorization and cap wavefunctions

The factorization viewpoint follows the holomorphic-block philosophy of Refs. [6, 7, 9, 66, 67] and its four-dimensional elliptic realization on geometries such as $D^2 \times T^2$ [8].

The cylinder $\text{Cyl} \times T^2$ has two boundary components, and its partition function is naturally a boundary-to-boundary kernel. At the level of one-loop determinants this kernel factorizes into two cap wavefunctions, equivalently the holomorphic blocks associated with the two caps of the cylinder. We do not repeat the three-dimensional factorization analysis: the factorization of $3d$ interval partition functions on $I \times T^2 \simeq \text{Cyl} \times S^1$ into hemisphere blocks has been worked out recently in Ref. [12]. We focus instead on the four-dimensional elliptic cylinder kernel and on its two-dimensional degeneration on $\text{Cyl} \simeq I \times S^1$.

The words “holomorphic” and “anti-holomorphic” below refer to the two opposite cap orientations in the gluing. They should not be interpreted as complex conjugation.

5.3.1 Four-dimensional factorization on $\text{Cyl} \times T^2$

We first discuss the factorization of the four-dimensional cylinder blocks on $\text{Cyl} \times T^2$. We use the full cylinder blocks defined in (4.34), with

$$B_3(u) := B_{3,3}(u; 1, \tau, \omega). \quad (5.108)$$

With the elliptic-Gamma conventions of Refs. [28–30], the elliptic Gamma function satisfies the gluing identity

$$\Gamma_e(u; \tau, \omega) \Gamma_e(\omega - u; \tau, \omega) = \frac{1}{\Theta(u; \omega)}. \quad (5.109)$$

Thus the theta factor appearing in the cylinder determinant can be regarded as the inverse product of two elliptic-Gamma cap wavefunctions. These cap wavefunctions are the $D^2 \times T^2$ holomorphic blocks of Refs. [6–9, 66, 67], written in the Shintani–Barnes normalization used in this paper. Compared with the common holomorphic-block conventions, we keep the Bernoulli contact factors explicit and use a reflected Dirichlet argument so that the elementary inverse relation is exactly $\mathcal{B}_D(u) \mathcal{B}_R(-u) = 1$.

We denote the $D^2 \times T^2$ cap wavefunctions by

$$\mathcal{B}_D(u) := \frac{\exp\left[-\frac{i\pi}{3} B_3(-u)\right]}{\Gamma_e(-u; \tau, \omega)}, \quad \mathcal{B}_R(u) := \exp\left[+\frac{i\pi}{3} B_3(u)\right] \Gamma_e(u; \tau, \omega). \quad (5.110)$$

The wavefunctions associated with the oppositely oriented cap are

$$\mathcal{B}_D^\vee(u) := \frac{\exp\left[-\frac{i\pi}{3} B_3(\omega + u)\right]}{\Gamma_e(\omega + u; \tau, \omega)}, \quad \mathcal{B}_R^\vee(u) := \exp\left[+\frac{i\pi}{3} B_3(\omega - u)\right] \Gamma_e(\omega - u; \tau, \omega). \quad (5.111)$$

The superscript $\vee$ refers to the opposite cap orientation in the gluing, not to Hermitian conjugation.

Using (5.109), one obtains the exact factorization

$$z_{DD}(u) = \mathcal{B}_D(u) \mathcal{B}_D^\vee(u), \quad z_{RR}(u) = \mathcal{B}_R(u) \mathcal{B}_R^\vee(u). \quad (5.112)$$

Moreover, the reflected inverse relation already holds separately at the level of the two caps:

$$\mathcal{B}_D(u) \mathcal{B}_R(-u) = 1, \quad \mathcal{B}_D^\vee(u) \mathcal{B}_R^\vee(-u) = 1. \quad (5.113)$$

Thus the elementary cancellation

$$z_{DD}(u) z_{RR}(-u) = 1 \quad (5.114)$$

is refined to two independent cap-wavefunction cancellations.

This immediately factorizes any integrand built out of z_{DD} and z_{RR} into cap and opposite-cap contributions. For example, the electric integrand of the $U(N_c) \leftrightarrow U(N_f - N_c)$ duality can be written as

$$Z_{\text{el}}(u) = Z_{\text{el}}^{\text{hol}}(u) Z_{\text{el}}^{\text{anti}}(u), \quad (5.115)$$

where

$$\begin{aligned}
Z_{\text{el}}^{\text{hol}}(u) &= \frac{1}{N_c!} \mathcal{B}_D \left[\sum_{a=1}^{N_c} u_a + M + \lambda \right] \mathcal{B}_D \left[- \sum_{a=1}^{N_c} u_a + \widetilde{M} + \lambda \right] \\
&\times \prod_{\substack{a,b=1 \\ a \neq b}}^{N_c} \mathcal{B}_D[u_a - u_b] \\
&\times \prod_{a=1}^{N_c} \prod_{i=1}^{N_f} \mathcal{B}_R[-u_a - m_i] \mathcal{B}_R[u_a - \widetilde{m}_i] \\
&\times \prod_{a=1}^{N_c} \prod_{\alpha=1}^n \mathcal{B}_D[u_a + b_\alpha] \mathcal{B}_D[-u_a + \widetilde{b}_\alpha].
\end{aligned} \tag{5.116}$$

The factor $Z_{\text{el}}^{\text{anti}}(u)$ is obtained from (5.116) by replacing $\mathcal{B}_D, \mathcal{B}_R$ with $\mathcal{B}_D^\vee, \mathcal{B}_R^\vee$.

Similarly, the magnetic integrand (5.16) factorizes as

$$Z_{\text{mag}}(v) = Z_{\text{mag}}^{\text{hol}}(v) Z_{\text{mag}}^{\text{anti}}(v). \tag{5.117}$$

Explicitly,

$$\begin{aligned}
Z_{\text{mag}}^{\text{hol}}(v) &= \frac{1}{n!} \mathcal{B}_D \left[- \sum_{r=1}^n v_r + \lambda \right] \mathcal{B}_D \left[\sum_{r=1}^n v_r + M + \widetilde{M} + \lambda \right] \\
&\times \prod_{\substack{r,s=1 \\ r \neq s}}^n \mathcal{B}_D[v_r - v_s] \\
&\times \prod_{r=1}^n \prod_{i=1}^{N_f} \mathcal{B}_R[v_r + \widetilde{m}_i] \mathcal{B}_D[-v_r + m_i] \\
&\times \prod_{r=1}^n \prod_{\alpha=1}^n \mathcal{B}_R[v_r - b_\alpha] \mathcal{B}_D[v_r - b_\alpha] \mathcal{B}_R[-v_r + b_\alpha] \mathcal{B}_R[-v_r - \widetilde{b}_\alpha].
\end{aligned} \tag{5.118}$$

The factor $Z_{\text{mag}}^{\text{anti}}(v)$ is obtained from (5.118) by replacing $\mathcal{B}_D, \mathcal{B}_R$ with $\mathcal{B}_D^\vee, \mathcal{B}_R^\vee$.

The identity inserted in (5.16) factorizes separately on the two caps:

$$\mathcal{B}_D[v_r - b_\alpha] \mathcal{B}_R[-v_r + b_\alpha] = 1, \quad \mathcal{B}_D^\vee[v_r - b_\alpha] \mathcal{B}_R^\vee[-v_r + b_\alpha] = 1. \tag{5.119}$$

Thus the cap-wavefunction factorization is compatible with the JK presentation without introducing any separate local factor to be split between the two caps.

The singlet sector factorizes in the same way:

$$Z_{\text{sing}} = Z_{\text{sing}}^{\text{hol}} Z_{\text{sing}}^{\text{anti}}. \tag{5.120}$$

For instance,

$$Z_M^{\text{hol}} = \prod_{i,j=1}^{N_f} \mathcal{B}_R[-m_i - \widetilde{m}_j], \quad Z_M^{\text{anti}} = \prod_{i,j=1}^{N_f} \mathcal{B}_R^\vee[-m_i - \widetilde{m}_j], \tag{5.121}$$

and similarly for the boundary singlet sector Z_B .

The residue expansion is compatible with this factorization. At an electric pole labelled by

$$\mathcal{I} \subset \{1, \dots, N_f\}, \quad |\mathcal{I}| = N_c, \tag{5.122}$$

the residue factorizes as

$$\mathcal{E}_{\mathcal{I}} = \mathcal{E}_{\mathcal{I}}^{\text{hol}} \mathcal{E}_{\mathcal{I}}^{\text{anti}}. \quad (5.123)$$

Likewise, at a magnetic pole labelled by

$$\mathcal{J} \subset \{1, \dots, N_f\}, \quad |\mathcal{J}| = n, \quad (5.124)$$

one has

$$\mathcal{M}_{\mathcal{J}} = \mathcal{M}_{\mathcal{J}}^{\text{hol}} \mathcal{M}_{\mathcal{J}}^{\text{anti}}. \quad (5.125)$$

In the magnetic b_a -sector, the completed DD/RR presentation separates the same-argument contact factor from the RR factors which lie in the opposite chamber. The reflected inverse pairs in (5.119) remain equal to one separately on each cap, while the same-argument DD/RR products contribute only holomorphic contact factors.

The term-by-term matching of the full cylinder duality,

$$Z_{\text{sing}} \mathcal{M}_{\mathcal{J}} = \mathcal{E}_{\mathcal{J}^c}, \quad (5.126)$$

is therefore refined to a matching of factorized kernels:

$$Z_{\text{sing}}^{\text{hol}} \mathcal{M}_{\mathcal{J}}^{\text{hol}} Z_{\text{sing}}^{\text{anti}} \mathcal{M}_{\mathcal{J}}^{\text{anti}} = \mathcal{E}_{\mathcal{J}^c}^{\text{hol}} \mathcal{E}_{\mathcal{J}^c}^{\text{anti}}, \quad (5.127)$$

with the completed DD/RR pairs inserted separately on the two caps.

5.3.2 The elliptic kernel interpretation

We now make the boundary-to-boundary interpretation of the cylinder block more explicit. The following bra-ket notation should be read as a formal boundary-state calculus for the localized wavefunctions on the elliptic fugacity torus. Its content is the set of exact identities among the cap wavefunctions introduced in (5.110) and (5.111); no additional Hilbert-space completion is assumed. Accordingly, this subsection should be viewed as a formal calculus for the elementary D/R matrix elements computed in this paper, rather than as a construction of a complete boundary-state basis. A more intrinsic kernel would require allowing general boundary interactions or defects, for instance boundary operators lifting Wilson-loop bases, and is not attempted here.

We denote a localized boundary wavefunction by

$$|u; \mathbf{q}\rangle, \quad \mathbf{q} \in \{D, R\}, \quad (5.128)$$

where u is the boundary fugacity and $\mathbf{q}$ is the boundary polarization. The cap state has components $\mathcal{B}_D, \mathcal{B}_R$, while the opposite cap has components $\mathcal{B}_D^\vee, \mathcal{B}_R^\vee$. As above, $\vee$ indicates the opposite orientation of the cap, not Hermitian conjugation. Equations (5.110) and (5.111) imply the elementary pairings

$$\mathcal{B}_R^\vee(u) \mathcal{B}_D(u - \omega) = 1, \quad \mathcal{B}_D^\vee(u) \mathcal{B}_R(u + \omega) = 1. \quad (5.129)$$

Equivalently, the affine shifts by ω can be absorbed into the formal gluing pairings

$$\langle u; R^\vee | v; D \rangle = \delta_E(v - u + \omega), \quad \langle u; D^\vee | v; R \rangle = \delta_E(v - u - \omega), \quad (5.130)$$

where δ_E is the formal delta function on the elliptic fugacity torus. These shifts are part of the gluing rule on the boundary torus.

The cylinder partition function is the localized matrix element of a boundary-to-boundary operator. For the four elementary polarizations we write

$$\mathcal{K}_{\mathbf{q}_+\mathbf{q}_-}^{\text{Cyl} \times T^2} = \int_E d\mu(u) |u; \mathbf{q}_+\rangle z_{\mathbf{q}_+\mathbf{q}_-}^{\text{Cyl} \times T^2}(u) \langle u + \mu_{\mathbf{q}_+\mathbf{q}_-}; \mathbf{q}_-^\vee|, \quad (5.131)$$

with

$$\mu_{RR} = 0, \quad \mu_{DD} = 0, \quad \mu_{DR} = \omega, \quad \mu_{RD} = -\omega. \quad (5.132)$$

In components this says

$$z_{\mathbf{q}_+\mathbf{q}_-}^{\text{Cyl} \times T^2}(u) = \mathcal{B}_{\mathbf{q}_+}(u) \mathcal{B}_{\mathbf{q}_-}^\vee(u + \mu_{\mathbf{q}_+\mathbf{q}_-}). \quad (5.133)$$

Using (5.129), one obtains the sewing identities

$$z_{RR}^{\text{Cyl} \times T^2}(u) \mathcal{B}_D(u - \omega) = \mathcal{B}_R(u), \quad z_{DD}^{\text{Cyl} \times T^2}(u) \mathcal{B}_R(u + \omega) = \mathcal{B}_D(u), \quad (5.134)$$

and

$$z_{DR}^{\text{Cyl} \times T^2}(u) \mathcal{B}_D(u) = \mathcal{B}_D(u), \quad z_{RD}^{\text{Cyl} \times T^2}(u) \mathcal{B}_R(u) = \mathcal{B}_R(u). \quad (5.135)$$

Equivalently,

$$\mathcal{K}_{RR}^{\text{Cyl} \times T^2} |\mathcal{B}_D\rangle = |\mathcal{B}_R\rangle, \quad \mathcal{K}_{DD}^{\text{Cyl} \times T^2} |\mathcal{B}_R\rangle = |\mathcal{B}_D\rangle, \quad (5.136)$$

while

$$\mathcal{K}_{DR}^{\text{Cyl} \times T^2} |\mathcal{B}_D\rangle = |\mathcal{B}_D\rangle, \quad \mathcal{K}_{RD}^{\text{Cyl} \times T^2} |\mathcal{B}_R\rangle = |\mathcal{B}_R\rangle. \quad (5.137)$$

Thus the mixed blocks are identity kernels in the two polarized sectors, whereas the unmixed blocks implement elementary changes of polarization.

This interpretation is compatible with radial slicing of the cap and belongs to the same broad family of gluing and cigar constructions studied in [13–15, 17]. Cutting the disk into annuli gives

$$D^2 \times T^2 = (\text{Cyl} \times T^2)_n \cup_{T^3} \cdots \cup_{T^3} (\text{Cyl} \times T^2)_1 \cup_{T^3} D^{2'} \times T^2. \quad (5.138)$$

In the boundary-state calculus this becomes a composition of the kernels (5.131). For instance,

$$\mathcal{K}_{DD}^{\text{Cyl} \times T^2} \mathcal{K}_{RR}^{\text{Cyl} \times T^2} = \mathcal{K}_{DR}^{\text{Cyl} \times T^2}, \quad \mathcal{K}_{RR}^{\text{Cyl} \times T^2} \mathcal{K}_{DD}^{\text{Cyl} \times T^2} = \mathcal{K}_{RD}^{\text{Cyl} \times T^2}, \quad (5.139)$$

where the shifts of the fugacity are fixed by the pairings (5.130). In components this is

$$z_{DD}^{\text{Cyl} \times T^2}(u) z_{RR}^{\text{Cyl} \times T^2}(u + \omega) = z_{DR}^{\text{Cyl} \times T^2}(u) = 1, \quad z_{RR}^{\text{Cyl} \times T^2}(u) z_{DD}^{\text{Cyl} \times T^2}(u - \omega) = z_{RD}^{\text{Cyl} \times T^2}(u) = 1. \quad (5.140)$$

More generally, any admissible word in the four elementary kernels is a formal radial slicing. Closed words in the two polarizations refine the same cap, while open words describe radial interfaces changing the boundary polarization.

The same identities also give the expected behavior of a massive chiral pair. Consider two chiral multiplets Q_a, Q_b with a superpotential mass term

$$W = mQ_a Q_b, \quad r_a + r_b = 2, \quad (5.141)$$

and with opposite gauge and flavor charges. If the first chiral has cylinder argument

$$x = \rho(u) + \nu + r_a \gamma_R \quad (5.142)$$

in an RR block, then the second chiral, in the complementary DD polarization, has argument

$$-\rho(u) - \nu + (r_b - 2)\gamma_R = -x. \quad (5.143)$$

Therefore the massive pair contributes the identity

$$z_{RR}(x)z_{DD}(-x) = 1. \quad (5.144)$$

Similarly,

$$z_{DD}(x)z_{RR}(-x) = 1, \quad z_{DR}(x)z_{RD}(-x) = 1, \quad z_{RD}(x)z_{DR}(-x) = 1. \quad (5.145)$$

Thus the usual statement that a massive chiral pair with a single supersymmetric vacuum contributes one to the protected index is realized pointwise by the reflected inverse relation of the cylinder blocks, provided the two chirals are assigned complementary boundary polarizations at the two ends. Non-complementary assignments require additional boundary terms or boundary degrees of freedom in order to make the massive superpotential compatible with the supersymmetric variational problem.

The same affine gluing calculus also reproduces the compact $S^2 \times T^2$ chiral determinant. Let γ_Φ denote the total additive fugacity appearing in the chiral determinant, including the gauge, flavor and R-symmetry contributions appropriate to the chiral multiplet under consideration. For an integer S^2 flux b , set

$$u_+ = -\gamma_\Phi + \frac{1-b}{2}\omega, \quad u_- = -\gamma_\Phi + \frac{1+b}{2}\omega, \quad (5.146)$$

so that $u_- - u_+ = b\omega$. The standard S^2 fusion with flux b pairs the Robin cap with the oppositely oriented Dirichlet cap as

$$Z_{S^2 \times T^2}^\Phi(\gamma_\Phi, b) = \langle \mathcal{B}_D^\vee | \mathbf{G}_{S^2}^{(b)} | \mathcal{B}_R \rangle = \mathcal{B}_R(u_+) \mathcal{B}_D^\vee(u_- - \omega). \quad (5.147)$$

Using (5.110) and (5.111), this gives

$$Z_{S^2 \times T^2}^\Phi(\gamma_\Phi, b) = \exp \left[\frac{i\pi}{3} (B_3(u_+) - B_3(u_-)) \right] \frac{\Gamma_e(u_+; \tau, \omega)}{\Gamma_e(u_-; \tau, \omega)}. \quad (5.148)$$

The match with the standard $T^2 \times S^2$ chiral determinant is obtained by using the elliptic-Gamma difference equation along the ω -direction,

$$\Gamma_e(u + \omega; \tau, \omega) = \Theta(u; \tau) \Gamma_e(u; \tau, \omega). \quad (5.149)$$

Since $u_- = u_+ + b\omega$, (5.148) becomes the finite theta-product formula

$$Z_{S^2 \times T^2}^\Phi(\gamma_\Phi, b) = \exp\left[\frac{i\pi}{3}(B_3(u_+) - B_3(u_-))\right] \begin{cases} \prod_{j=0}^{b-1} \Theta(u_+ + j\omega; \tau)^{-1}, & b > 0, \\ 1, & b = 0, \\ \prod_{j=1}^{-b} \Theta(u_+ - j\omega; \tau), & b < 0. \end{cases} \quad (5.150)$$

Equivalently, for $b > 0$,

$$\prod_{j=0}^{b-1} \Theta(u_+ + j\omega; \tau)^{-1} = \prod_{j=0}^{b-1} \Theta\left(-\gamma_\Phi + \frac{1-b+2j}{2}\omega; \tau\right)^{-1}. \quad (5.151)$$

The exponential prefactor in (5.150) is the local Bernoulli contact term associated with our Shintani–Barnes normalization of the cap wavefunctions. After the corresponding local counterterm normalization, the non-local determinant is precisely the finite theta-product form of the $T^2 \times S^2$ chiral index of Closset and Shamir [68]. Thus both the open cylinder block and the closed $S^2 \times T^2$ determinant are built from the same cap wavefunctions and the same affine gluing shifts on the boundary elliptic curve. This is the sense in which the localized partition function on $\text{Cyl} \times T^2$ defines an elliptic kernel: it is the matrix element of a boundary-to-boundary operator whose entries are meromorphic elliptic functions of the boundary fugacity and whose sewing with $D^2 \times T^2$ holomorphic-block cap states reproduces the expected cap and compact-space blocks.

5.3.3 Two-dimensional factorization on Cyl

The same structure appears in the two-dimensional cylinder degeneration. The elementary blocks are

$$z_{\text{RR}}^{2d}(u) = \frac{1}{2i \sin(\pi u)}, \quad z_{\text{DD}}^{2d}(u) = -2i \sin(\pi u). \quad (5.152)$$

Using Euler’s reflection formula,

$$\Gamma(u)\Gamma(1-u) = \frac{\pi}{\sin(\pi u)}, \quad (5.153)$$

one obtains

$$z_{\text{RR}}^{2d}(u) = \frac{\Gamma(u)\Gamma(1-u)}{2\pi i}, \quad z_{\text{DD}}^{2d}(u) = \frac{2\pi i}{\Gamma(-u)\Gamma(1+u)}. \quad (5.154)$$

Choosing a branch of $(2\pi i)^{1/2}$, we define the two-dimensional cap wavefunctions by

$$\mathcal{B}_{\text{R}}^{2d}(u) := \frac{\Gamma(u)}{(2\pi i)^{1/2}}, \quad \mathcal{B}_{\text{D}}^{2d}(u) := \frac{(2\pi i)^{1/2}}{\Gamma(-u)}. \quad (5.155)$$

The opposite-cap wavefunctions are

$$\mathcal{B}_{\text{R}}^{2d,\vee}(u) := \frac{\Gamma(1-u)}{(2\pi i)^{1/2}}, \quad \mathcal{B}_{\text{D}}^{2d,\vee}(u) := \frac{(2\pi i)^{1/2}}{\Gamma(1+u)}. \quad (5.156)$$

The superscript $\vee$ again denotes the opposite orientation of the cap, not Hermitian conjugation. With these definitions,

$$z_{\text{RR}}^{2d}(u) = \mathcal{B}_{\text{R}}^{2d}(u) \mathcal{B}_{\text{R}}^{2d,\vee}(u), \quad z_{\text{DD}}^{2d}(u) = \mathcal{B}_{\text{D}}^{2d}(u) \mathcal{B}_{\text{D}}^{2d,\vee}(u), \quad (5.157)$$

and the reflected inverse relation already holds at the level of the cap wavefunctions:

$$\mathcal{B}_{\text{D}}^{2d}(u) \mathcal{B}_{\text{R}}^{2d}(-u) = 1, \quad \mathcal{B}_{\text{D}}^{2d,\vee}(u) \mathcal{B}_{\text{R}}^{2d,\vee}(-u) = 1. \quad (5.158)$$

The gluing identities are the two-dimensional degeneration of (5.129), with the unit shift replacing the elliptic shift ω :

$$\mathcal{B}_{\text{R}}^{2d,\vee}(u) \mathcal{B}_{\text{D}}^{2d}(u-1) = 1, \quad \mathcal{B}_{\text{D}}^{2d,\vee}(u) \mathcal{B}_{\text{R}}^{2d}(u+1) = 1. \quad (5.159)$$

Thus the boundary-to-boundary interpretation also applies to the two-dimensional cylinder $\text{Cyl} \simeq I \times S^1$. In a formal localized boundary basis $|u; \mathbf{q}\rangle$, with $u \sim u+1$, the elementary two-dimensional kernels can be written as

$$\mathcal{K}_{\mathbf{q}_+\mathbf{q}_-}^{2d} = \int_{\mathbb{C}/\mathbb{Z}} d\mu(u) |u; \mathbf{q}_+\rangle z_{\mathbf{q}_+\mathbf{q}_-}^{2d}(u) \langle u + \mu_{\mathbf{q}_+\mathbf{q}_-}^{2d}; \mathbf{q}_-^\vee|, \quad (5.160)$$

where

$$\mu_{\text{RR}}^{2d} = 0, \quad \mu_{\text{DD}}^{2d} = 0, \quad \mu_{\text{DR}}^{2d} = 1, \quad \mu_{\text{RD}}^{2d} = -1. \quad (5.161)$$

The component form is

$$z_{\mathbf{q}_+\mathbf{q}_-}^{2d}(u) = \mathcal{B}_{\mathbf{q}_+}^{2d}(u) \mathcal{B}_{\mathbf{q}_-}^{2d,\vee}(u + \mu_{\mathbf{q}_+\mathbf{q}_-}^{2d}). \quad (5.162)$$

Consequently,

$$z_{\text{DR}}^{2d}(u) = \mathcal{B}_{\text{D}}^{2d}(u) \mathcal{B}_{\text{R}}^{2d,\vee}(u+1) = 1, \quad z_{\text{RD}}^{2d}(u) = \mathcal{B}_{\text{R}}^{2d}(u) \mathcal{B}_{\text{D}}^{2d,\vee}(u-1) = 1, \quad (5.163)$$

while

$$z_{\text{DD}}^{2d}(u) z_{\text{RR}}^{2d}(u+1) = 1, \quad z_{\text{RR}}^{2d}(u) z_{\text{DD}}^{2d}(u-1) = 1. \quad (5.164)$$

These equations are the two-dimensional analogues of (5.135) and (5.140). They show that further slicing of the two-dimensional cylinder is again represented by composition of identity kernels and polarization-changing interfaces. The massive-pair identity degenerates in the same way:

$$z_{\text{RR}}^{2d}(u) z_{\text{DD}}^{2d}(-u) = 1, \quad z_{\text{DD}}^{2d}(u) z_{\text{RR}}^{2d}(-u) = 1. \quad (5.165)$$

For the complete-intersection GLSM, the cylinder integrand

$$Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(u) = \prod_{A=1}^r z_{\text{RR}}^{2d}[-\mathbf{n}_A u + \mathbf{p}_A] \prod_{i=1}^N z_{\text{RR}}^{2d}[u + x_i] \quad (5.166)$$

factorizes as

$$Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d}(u) = Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d, \text{hol}}(u) Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d, \text{anti}}(u), \quad (5.167)$$

where the two factors are now written directly in terms of the cap wavefunctions,

$$Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d, \text{hol}}(u) = \prod_{A=1}^r \mathcal{B}_{\text{R}}^{2d}[-\mathbf{n}_A u + \mathbf{p}_A] \prod_{i=1}^N \mathcal{B}_{\text{R}}^{2d}[u + x_i], \quad (5.168)$$

and

$$Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d, \text{anti}}(u) = \prod_{A=1}^r \mathcal{B}_R^{2d, \vee}[-\mathbf{n}_A u + \mathbf{p}_A] \prod_{i=1}^N \mathcal{B}_R^{2d, \vee}[u + x_i]. \quad (5.169)$$

The chamber equality

$$\text{JK}_+ Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d} = \text{JK}_- Z_{\mathbf{n}_1, \dots, \mathbf{n}_r}^{2d} \quad (5.170)$$

can therefore be interpreted as an equality between two residue decompositions of the same factorized cylinder kernel. In the positive chamber, the poles are

$$u_i^+ = -x_i, \quad i = 1, \dots, N, \quad (5.171)$$

and give the geometric-phase expansion. In the negative chamber, the poles are

$$u_{A, \ell}^- = \frac{\mathbf{p}_A - \ell}{\mathbf{n}_A}, \quad \ell = 0, \dots, \mathbf{n}_A - 1, \quad (5.172)$$

and give the hybrid or Landau–Ginzburg expansion.

Thus the two-dimensional identity has the same kernel structure.

The cylinder partition function is a two-boundary kernel. Its residue expansion is a sum of factorized contributions,

$$Z_{\text{Cyl}} \sim \sum_{\alpha} B_{\alpha}^{\text{hol}} B_{\alpha}^{\text{anti}}. \quad (5.173)$$

For $\text{Cyl} \times T^2$, the cap wavefunctions are built from elliptic Gamma functions. For $\text{Cyl} \simeq I \times S^1$, the cap wavefunctions are built from Euler Gamma functions.

5.4 Holographic theories on $\text{Cyl} \times T^2$

Two standard four-dimensional matter contents provide useful benchmarks for theories that also have holographic incarnations: $\mathcal{N} = 4$ SYM with gauge group $SU(N)$ and the Klebanov–Witten conifold theory [69] with gauge group $SU(N) \times SU(N)$. The point we emphasize is that the naive assignment in which all chiral multiplets are placed in RR polarization does not define a ω -elliptic meromorphic integrand on the ω -elliptic gauge-fugacity torus of $\text{Cyl} \times T^2$. However, by changing the boundary polarization of a suitable subset of chiral multiplets, equivalently by replacing some z_{RR} factors by z_{DD} factors, one obtains ω -elliptic cylinder integrands.

We assign a sign

$$\sigma_{\text{ch}} = \begin{cases} +1, & \text{chiral in } RR \text{ polarization,} \\ -1, & \text{chiral in } DD \text{ polarization.} \end{cases} \quad (5.174)$$

The gauge-dependent part of the ω -elliptic monodromy is controlled by the quadratic terms in the Bernoulli phases together with the theta-function quasi-periodicities. In the examples below, cancellation of this gauge-dependent monodromy imposes simple linear constraints on the signs σ_{ch} .

5.4.1 $\mathcal{N} = 4$ SYM with gauge group $SU(N)$

We first consider $\mathcal{N} = 4$ SYM in $\mathcal{N} = 1$ language, one of the canonical gauge-theory examples in the AdS/CFT correspondence [70]. The theory contains one vector multiplet and three adjoint chiral multiplets

$$X_I \in \text{Adj}_{SU(N)}, \quad I = 1, 2, 3, \quad (5.175)$$

with superpotential

$$W_{\mathcal{N}=4} = \text{Tr } X_1[X_2, X_3]. \quad (5.176)$$

Let

$$u_a, \quad a = 1, \dots, N, \quad (5.177)$$

be the $SU(N)$ gauge fugacities, subject to

$$\sum_{a=1}^N u_a = 0. \quad (5.178)$$

The naive cylinder integrand, with all three adjoint chirals in RR polarization, is

$$\begin{aligned} Z_{\mathcal{N}=4}^{\text{naive}} &= \frac{1}{N!} [z_{RR}(\Delta_1) z_{RR}(\Delta_2) z_{RR}(\Delta_3)]^{N-1} \\ &\times \prod_{\substack{a,b=1 \\ a \neq b}}^N z_{DD}(u_{ab}) \prod_{I=1}^3 z_{RR}(u_a - u_b + \Delta_I). \end{aligned} \quad (5.179)$$

This expression is not ω -elliptic on the $SU(N)$ gauge fugacity torus. Indeed, the three adjoint chiral multiplets have polarization signs

$$\sigma_1 + \sigma_2 + \sigma_3 = 1 + 1 + 1 = 3, \quad (5.180)$$

whereas cancellation of the gauge-dependent ω -elliptic monodromy of the vector multiplet requires

$$\sigma_1 + \sigma_2 + \sigma_3 = 1. \quad (5.181)$$

Thus the all- RR assignment over-cancels the vector contribution and leaves a nontrivial ω -elliptic multiplier.

An ω -elliptic assignment is obtained by taking two adjoint chirals in RR polarization and one in DD polarization. We choose

$$X_1, X_2 \in RR, \quad X_3 \in DD. \quad (5.182)$$

Then

$$\sigma_1 + \sigma_2 + \sigma_3 = 1 + 1 - 1 = 1, \quad (5.183)$$

and the condition (5.181) is satisfied. The corresponding ω -elliptic $SU(N)$ contour integral is the one displayed in (1.8). The factor raised to the power $N - 1$ there is the Cartan contribution of the three adjoint chirals of $SU(N)$, while the product over $a \neq b$ gives the contribution of the roots. The superpotential imposes the usual flavor balancing condition

$$\Delta_1 + \Delta_2 + \Delta_3 = \Delta_W \quad \text{mod } \mathbb{Z} + \omega\mathbb{Z}. \quad (5.184)$$

This condition is independent of the cancellation of gauge ω -elliptic monodromies; it expresses the neutrality of the superpotential (5.176).

5.4.2 Klebanov–Witten theory

We now consider the Klebanov–Witten conifold quiver [69]. The gauge group is

$$SU(N)_u \times SU(N)_v, \quad (5.185)$$

with gauge fugacities

$$u_a, \quad v_b, \quad a, b = 1, \dots, N, \quad (5.186)$$

subject to

$$\sum_{a=1}^N u_a = 0, \quad \sum_{b=1}^N v_b = 0. \quad (5.187)$$

The matter fields are

$$A_i \in (\mathbf{N}_u, \overline{\mathbf{N}}_v), \quad i = 1, 2, \quad (5.188)$$

and

$$B_j \in (\overline{\mathbf{N}}_u, \mathbf{N}_v), \quad j = 1, 2. \quad (5.189)$$

The superpotential is

$$W_{\text{KW}} = \text{Tr} (A_1 B_1 A_2 B_2 - A_1 B_2 A_2 B_1). \quad (5.190)$$

The naive all- RR integrand is

$$\begin{aligned} Z_{\text{KW}}^{\text{naive}} &= \prod_{\substack{a,b=1 \\ a \neq b}}^N z_{\text{DD}}(u_{ab}) \prod_{\substack{a,b=1 \\ a \neq b}}^N z_{\text{DD}}(v_a - v_b) \\ &\times \prod_{a,b=1}^N z_{\text{RR}}(u_a - v_b + \alpha_1) z_{\text{RR}}(u_a - v_b + \alpha_2) \\ &\times \prod_{a,b=1}^N z_{\text{RR}}(-u_a + v_b + \beta_1) z_{\text{RR}}(-u_a + v_b + \beta_2). \end{aligned} \quad (5.191)$$

This expression is not ω -elliptic. For the conifold quiver with equal ranks $N = N_u = N_v$, cancellation of the gauge-dependent monodromy of the two vector multiplets requires

$$\sigma_{A_1} + \sigma_{A_2} + \sigma_{B_1} + \sigma_{B_2} = 2. \quad (5.192)$$

The naive all- RR assignment gives instead

$$1 + 1 + 1 + 1 = 4, \quad (5.193)$$

and hence leaves a nontrivial ω -elliptic multiplier.

An ω -elliptic polarization is obtained by taking three bifundamentals in RR polarization and one in DD polarization. For definiteness we choose

$$A_1, A_2, B_1 \in RR, \quad B_2 \in DD. \quad (5.194)$$

Then

$$\sigma_{A_1} + \sigma_{A_2} + \sigma_{B_1} + \sigma_{B_2} = 1 + 1 + 1 - 1 = 2, \quad (5.195)$$

and (5.192) is satisfied. The corresponding ω -elliptic cylinder integrand is

$$\begin{aligned} Z_{\text{KW}}^{SU(N) \times SU(N)} &= \prod_{\substack{a,b=1 \\ a \neq b}}^N z_{\text{DD}}(u_{ab}) \prod_{\substack{a,b=1 \\ a \neq b}}^N z_{\text{DD}}(v_a - v_b) \\ &\times \prod_{a,b=1}^N z_{\text{RR}}(u_a - v_b + \alpha_1) z_{\text{RR}}(u_a - v_b + \alpha_2) \\ &\times \prod_{a,b=1}^N z_{\text{RR}}(-u_a + v_b + \beta_1) z_{\text{DD}}(-u_a + v_b + \beta_2), \end{aligned} \quad (5.196)$$

with the $SU(N) \times SU(N)$ constraints (5.187). The superpotential imposes the balancing condition

$$\alpha_1 + \alpha_2 + \beta_1 + \beta_2 = \Delta_W \mod \mathbb{Z} + \omega\mathbb{Z}. \quad (5.197)$$

The choice (5.194) is not unique. Any assignment with three bifundamentals in RR polarization and one in DD polarization satisfies (5.192). Different choices make different subgroups of the flavor symmetry manifest, but they are equivalent from the viewpoint of ω -ellipticity.

A Conventions and Auxiliary Formulae

This appendix collects conventions and auxiliary formulae used in the main text.

A.1 Spinors and sigma matrices

We use two-component Euclidean spinors. Undotted spinors, such as ζ_α , transform in the $(\mathbf{2}, \mathbf{1})$ representation of $\text{Spin}(4) = SU(2)_+ \times SU(2)_-$, while dotted spinors, such as $\tilde{\zeta}_{\dot{\alpha}}$, transform in $(\mathbf{1}, \mathbf{2})$. Hermitian conjugation gives spinors with the conjugate index structure, and the norms are

$$|\zeta|^2 = \zeta_\alpha^\dagger \zeta^\alpha, \quad |\tilde{\zeta}|^2 = \tilde{\zeta}_{\dot{\alpha}}^\dagger \tilde{\zeta}^{\dot{\alpha}}. \quad (A.1)$$

Spinor indices are raised and lowered with antisymmetric tensors $\epsilon_{\alpha\beta}$, $\epsilon^{\alpha\beta}$, $\epsilon_{\dot{\alpha}\dot{\beta}}$, $\epsilon^{\dot{\alpha}\dot{\beta}}$. We use

$$\epsilon^{12} = +1, \quad \epsilon_{12} = -1, \quad \epsilon^{\alpha\beta} \epsilon_{\beta\gamma} = \delta^\alpha_\gamma, \quad \epsilon_{\alpha\beta} \epsilon^{\beta\gamma} = \delta_\alpha^\gamma, \quad (A.2)$$

and similarly for dotted indices. These are the raising and lowering conventions used in the Wolfram Mathematica checks of the background. We contract spinors as $\zeta\chi = \zeta^\alpha \chi_\alpha$ and $\tilde{\zeta}\tilde{\chi} = \tilde{\zeta}_{\dot{\alpha}} \tilde{\chi}^{\dot{\alpha}}$.

In an orthonormal frame the Euclidean sigma matrices are

$$\sigma^a = (\sigma^1, \sigma^2, \sigma^3, -i\mathbf{1}_2), \quad \tilde{\sigma}^a = (-\sigma^1, -\sigma^2, -\sigma^3, -i\mathbf{1}_2), \quad (A.3)$$

where the first three matrices on the right-hand side are the Pauli matrices. Curved sigma matrices are obtained by contraction with the frame and inverse frame:

$$\sigma^\mu = E_a^\mu \sigma^a, \quad \tilde{\sigma}^\mu = E_a^\mu \tilde{\sigma}^a, \quad \sigma_\mu = e_\mu^a \sigma_a, \quad \tilde{\sigma}_\mu = e_\mu^a \tilde{\sigma}_a. \quad (A.4)$$

The generators acting on the two chiralities are

$$\sigma^{ab} := \frac{1}{4}(\sigma^a \tilde{\sigma}^b - \sigma^b \tilde{\sigma}^a), \quad \tilde{\sigma}^{ab} := \frac{1}{4}(\tilde{\sigma}^a \sigma^b - \tilde{\sigma}^b \sigma^a). \quad (\text{A.5})$$

With $\varepsilon_{1234} = +1$, they obey

$$\frac{1}{2}\varepsilon_{abcd}\sigma^{cd} = \sigma_{ab}, \quad \frac{1}{2}\varepsilon_{abcd}\tilde{\sigma}^{cd} = -\tilde{\sigma}_{ab}. \quad (\text{A.6})$$

The Clifford relations and hermiticity properties are

$$\sigma^a \tilde{\sigma}^b + \sigma^b \tilde{\sigma}^a = -2\delta^{ab}\mathbf{1}_2, \quad \tilde{\sigma}^a \sigma^b + \tilde{\sigma}^b \sigma^a = -2\delta^{ab}\mathbf{1}_2, \quad (\text{A.7})$$

$$(\sigma^a)^\dagger = -\tilde{\sigma}^a, \quad (\sigma^{ab})^\dagger = -\sigma^{ab}, \quad (\tilde{\sigma}^{ab})^\dagger = -\tilde{\sigma}^{ab}. \quad (\text{A.8})$$

$$\sigma_{\alpha\dot{\alpha}}^a \tilde{\sigma}_a^{\dot{\beta}\beta} = -2\delta_\alpha^\beta \delta_{\dot{\alpha}}^{\dot{\beta}}, \quad (\text{A.9})$$

$$\begin{aligned} \sigma^a \tilde{\sigma}^b \sigma^c &= -\delta^{ab}\sigma^c + \delta^{ac}\sigma^b - \delta^{bc}\sigma^a + \varepsilon^{abcd}\sigma_d, \\ \tilde{\sigma}^a \sigma^b \tilde{\sigma}^c &= -\delta^{ab}\tilde{\sigma}^c + \delta^{ac}\tilde{\sigma}^b - \delta^{bc}\tilde{\sigma}^a - \varepsilon^{abcd}\tilde{\sigma}_d. \end{aligned} \quad (\text{A.10})$$

For commuting supersymmetry parameters one has the exchange rules

$$\zeta\chi = -\chi\zeta, \quad \tilde{\zeta}\tilde{\chi} = -\tilde{\chi}\tilde{\zeta}, \quad \zeta\sigma^a\tilde{\chi} = \tilde{\chi}\tilde{\sigma}^a\zeta, \quad \zeta\sigma^{ab}\chi = \chi\sigma^{ab}\zeta. \quad (\text{A.11})$$

If two dynamical fermions are exchanged, the additional minus sign from their Grassmann parity must also be included. A useful Fierz identity, in these conventions, is

$$(\zeta\chi)(\tilde{\zeta}\tilde{\chi}) = -\frac{1}{2}(\zeta\sigma^a\tilde{\chi})(\chi\sigma_a\tilde{\zeta}). \quad (\text{A.12})$$

The spinor covariant derivatives are

$$\nabla_\mu\zeta = \partial_\mu\zeta - \frac{1}{2}\omega_{\mu ab}\sigma^{ab}\zeta, \quad \nabla_\mu\tilde{\zeta} = \partial_\mu\tilde{\zeta} - \frac{1}{2}\omega_{\mu ab}\tilde{\sigma}^{ab}\tilde{\zeta}. \quad (\text{A.13})$$

The spin connection is determined by the torsion-free Cartan equation

$$de^a + \omega^a_b \wedge e^b = 0, \quad (\text{A.14})$$

Equivalently, in components, with E_a^μ the inverse frame,

$$\omega_{\mu ab} = -e_{b\nu} \left(\partial_\mu E_a^\nu + \Gamma_{\mu\lambda}^\nu E_a^\lambda \right), \quad (\text{A.15})$$

which is the component convention used in the symbolic checks. The curvature convention is

$$R_{\mu\nu}{}^{ab} = \partial_\mu\omega_\nu{}^{ab} - \partial_\nu\omega_\mu{}^{ab} + \omega_\mu{}^{ac}\omega_{\nu c}{}^b - \omega_\nu{}^{ac}\omega_{\mu c}{}^b. \quad (\text{A.16})$$

A.2 Shintani–Barnes regularization

We define the q -Pochhammer symbol by

$$(z; q)_\infty := \prod_{k=0}^{\infty} (1 - zq^k). \quad (\text{A.17})$$

For additive variables we use throughout the theta function

$$\Theta(u; \omega) := (e^{2\pi i u}; e^{2\pi i \omega})_\infty (e^{2\pi i(\omega - u)}; e^{2\pi i \omega})_\infty. \quad (\text{A.18})$$

and similarly for the double q -Pochhammer symbol and elliptic Gamma function

$$(z; q, p)_\infty := \prod_{k,j=0}^{\infty} (1 - zq^k p^j), \quad \Gamma(z; q, p) := \frac{(pqz^{-1}; q, p)_\infty}{(z; q, p)_\infty}. \quad (\text{A.19})$$

A useful property is

$$\frac{\Gamma(pz; q, p)}{\Gamma(z; q, p)} = \Theta(u; \omega), \quad z = e^{2\pi i u}, \quad q = e^{2\pi i \omega}. \quad (\text{A.20})$$

Define the fugacities

$$z := e^{2\pi i u}, \quad q := e^{2\pi i \omega}, \quad p := e^{2\pi i \tau}. \quad (\text{A.21})$$

The product representations in (A.17)–(A.18) converge for $|q|, |p| < 1$ (equivalently $\Im(\omega), \Im(\tau) > 0$), while the final expressions below define the analytically continued regularization for generic ω, τ . The regularization is ordered: we first regularize the torus lattice and then the half-cylinder product. This ordering fixes the Bernoulli phase, and a different ordering would shift the answer by a holomorphic non-vanishing contact factor, equivalently by a choice of local counterterm. The symbols $C_{4d}(\omega, \tau)$, $C_{3d}(\omega)$ and C_{2d} below denote such scheme-dependent normalizations independent of the holonomy variable. We also use the additive elliptic-Gamma notation

$$\Gamma_e(u; \tau, \omega) := \Gamma(z; p, q). \quad (\text{A.22})$$

The fundamental unregularized product we encounter in the computation of 1-loop determinants on $D^2 \times T^2$ or $\text{Cyl} \times T^2$ is

$$\prod_{m_\varphi \geq 0} \prod_{(m_x, m_y) \in \mathbb{Z}^2} (\omega m_\varphi + \tau m_x - m_y + u). \quad (\text{A.23})$$

This product is to be understood in the Shintani–Barnes sense, i.e. by analytic continuation of the Hurwitz multiple zeta function and its associated multiple Shintani-Barnes Gamma functions [30, 64, 65]. The elliptic-Gamma form and the accompanying Bernoulli phase are consistent with the modular and product formulae of Refs. [28, 29]. Namely,

$$\prod_{m_\varphi \geq 0} \prod_{(m_x, m_y) \in \mathbb{Z}^2} (\omega m_\varphi + \tau m_x - m_y + \mu) \stackrel{\text{SB}}{=} C_{4d}(\omega, \tau) \frac{e^{-i\frac{\pi}{3} B_{3,3}(\mu|1, \tau, \omega)}}{\Gamma(e^{2\pi i \mu}; e^{2\pi i \tau}, e^{2\pi i \omega})}, \quad (\text{A.24})$$

where $C_{4d}(\omega, \tau)$ depends only on ω, τ (and on the chosen normalization of the Barnes double Gamma) and is independent of μ . Here we defined the cubic Bernoulli polynomial

$$B_{3,3}(u; \omega_1, \omega_2, \omega_3) = \frac{u^3}{\omega_1 \omega_2 \omega_3} - \frac{3u^2 (\omega_1 + \omega_2 + \omega_3)}{2\omega_1 \omega_2 \omega_3} + \frac{u (\omega_1^2 + \omega_2^2 + \omega_3^2 + 3(\omega_1 \omega_2 + \omega_3 \omega_2 + \omega_1 \omega_3))}{2\omega_1 \omega_2 \omega_3} - \frac{(\omega_1 + \omega_2 + \omega_3) (\omega_1 \omega_2 + \omega_3 \omega_2 + \omega_1 \omega_3)}{4\omega_1 \omega_2 \omega_3} \quad (\text{A.25})$$

A.3 Jeffrey–Kirwan residue conventions

This appendix fixes the residue convention used throughout the paper. The purpose is only to make the normalization of the contour measure explicit. In particular, every occurrence of $\text{JK}_{U(r), \eta} Z$ in the main text is normalized with the measure

$$\prod_{a=1}^r \frac{du_a}{2\pi i}. \quad (\text{A.26})$$

Thus no extra factor of $(2\pi i)^r$ is missing from the residue formulae in the duality identities.

Let $u = (u_1, \dots, u_r)$ be local coordinates near a pole u_* , and suppose that the polar hyperplanes passing through u_* are locally given by

$$Q_j(u - u_*) = 0, \quad j = 1, \dots, r, \quad (\text{A.27})$$

where the charge covectors $Q_j \in (\mathbb{R}^r)^*$ are linearly independent. Near such a non-degenerate pole, a meromorphic r -form can be written as

$$\omega(u) = \frac{f(u) du_1 \wedge \dots \wedge du_r}{Q_1(u - u_*) \dots Q_r(u - u_*)} + \text{less singular terms}, \quad (\text{A.28})$$

with f holomorphic at u_* . The Jeffrey–Kirwan residue in chamber $\eta \in \mathbb{R}^r$ is

$$\text{JK-Res}_{u=u_*, \eta} \omega = \begin{cases} \frac{f(u_*)}{|\det(Q_1, \dots, Q_r)|}, & \eta \in \text{Cone}(Q_1, \dots, Q_r), \\ 0, & \eta \notin \text{Cone}(Q_1, \dots, Q_r), \end{cases} \quad (\text{A.29})$$

up to the orientation convention chosen for the ordered set of charges. In the convention used in the main text, the orientation is fixed so that a simple positive-charge pole of the one-dimensional factor $z_{\text{RR}}(u)$ contributes positively.

Equivalently, for an isolated pole selected by the chamber, the JK residue can be represented by the normalized contour integral

$$\text{JK-Res}_{u=u_*, \eta} \omega = \int_{\Gamma_{u_*, \eta}} \prod_{a=1}^r \frac{du_a}{2\pi i} \frac{f(u)}{Q_1(u - u_*) \dots Q_r(u - u_*)}, \quad (\text{A.30})$$

where $\Gamma_{u_*, \eta}$ is the small real r -torus encircling the selected hyperplanes with the orientation compatible with η . If one wrote the same contour integral using $\prod_a du_a$ instead, the answer would be multiplied by $(2\pi i)^r$.

For the cylinder blocks, the relevant simple poles come from the RR factor

$$z_{\text{RR}}(u) = \frac{\exp\left[\frac{i\pi}{3}P_3(u)\right]}{\Theta(u;\omega)}. \quad (\text{A.31})$$

Since

$$\Theta(u;\omega) = \Theta'_0 u + O(u^2), \quad \Theta'_0 = \left. \frac{\partial}{\partial u} \Theta(u;\omega) \right|_{u=0}, \quad (\text{A.32})$$

one has

$$z_{\text{RR}}(u) = \frac{\varphi_0}{\Theta'_0} \frac{1}{u} + O(1), \quad \varphi_0 := \exp\left[\frac{i\pi}{3}P_3(0)\right]. \quad (\text{A.33})$$

Therefore a selected simple pole of a positive-charge RR hyperplane contributes

$$\frac{\varphi_0}{\Theta'_0} \quad (\text{A.34})$$

in the normalized JK convention. This is the origin of the powers of φ_0/Θ'_0 in the residue formulae in the main text.

The theta-function confining identity in Subsection 5.1.3 is written directly as a contour integral rather than as the abstract symbol JK. For this reason its measure is displayed explicitly as

$$\prod_{a=1}^N \frac{du_a}{2\pi i}. \quad (\text{A.35})$$

With this normalization the theta identity has the right-hand side shown in the main text. With the unnormalized measure $\prod_a du_a$, the left-hand side would be multiplied by $(2\pi i)^N$.

B Anomaly Equations

In this appendix the periods of the cubic Bernoulli polynomial are kept arbitrary. We use the same symbol as in the main text, namely

$$B_3(u) := B_{3,3}(u;\omega_1,\omega_2,\omega_3), \quad P_3(x) := B_3(x) + B_3(\omega_3 - x). \quad (\text{B.1})$$

A Robin block with argument u contributes $P_3(u)$, while a Dirichlet block with argument u contributes the negative of the same expression evaluated at $-u$. Whenever a DD multiplet appears, its contribution is obtained by applying this reflection to its actual block argument. This is the anomaly-polynomial version of the inverse relation $z_{\text{DD}}(u)z_{\text{RR}}(-u) = 1$. For the $U(N_c)$ vector contribution we get the polynomial

$$\sum_{i,j=1}^{N_c} P_3(x_i - x_j) = 2A \left[N_c \sum_i x_i^2 - \left(\sum_i x_i \right)^2 \right] + N_c^2 C \quad (\text{B.2})$$

where

$$A := -\frac{3(\omega_1 + \omega_2)}{\omega_1\omega_2\omega_3}, \quad C := -\frac{(\omega_1 + \omega_2)(\omega_1\omega_2 + \omega_3^2)}{2\omega_1\omega_2\omega_3}. \quad (\text{B.3})$$

For an adjoint DD/RR multiplet we get

$$\pm \sum_{i,j=1}^{N_c} P_3(u_i - u_j + t) = \pm 2A \left[N_c \sum_i u_i^2 - \left(\sum_i u_i \right)^2 \right] \pm N_c^2 [At(t - \omega_3) + C]. \quad (\text{B.4})$$

For a DD/RR determinant multiplet we get

$$P_3\left(\sum_{i=1}^{N_c} u_i + M\right) = \pm A \left[\left(\sum_i u_i\right)^2 + (2M - \omega_3) \sum_i u_i \right] \pm [AM(M - \omega_3) + C] , \quad (\text{B.5})$$

and similarly for a DD/RR anti-determinant multiplet

$$\pm P_3\left(-\sum_{i=1}^{N_c} u_i + \bar{M}\right) = \pm A \left[\left(\sum_i u_i\right)^2 - (2\bar{M} - \omega_3) \sum_i u_i \right] \pm [A\bar{M}(\bar{M} - \omega_3) + C] . \quad (\text{B.6})$$

For a fundamental DD/RR multiplet we get the contribution

$$\pm \sum_i P_3(u_i + m) = \pm A \left[\sum_i u_i^2 + (2m - \omega_3) \sum_i u_i \right] \pm N_c [Am(m - \omega_3) + C] , \quad (\text{B.7})$$

and similarly for an anti-fundamental DD/RR multiplet

$$\pm \sum_i P_3(-u_i + \bar{m}) = \pm A \left[\sum_i u_i^2 - (2\bar{m} - \omega_3) \sum_i u_i \right] \pm N_c [A\bar{m}(\bar{m} - \omega_3) + C] . \quad (\text{B.8})$$

Consider a vector, adjoints (N_{ad}), (anti-)determinants (N_{d} , $\bar{N}_{\text{d}}$) and (anti-)fundamentals (N_{f} , $\bar{N}_{\text{f}}$), contributing with signs (i.e. DD/RR conditions) $\epsilon_a, \eta_d, \bar{\eta}_{\bar{d}}, \rho_f, \bar{\rho}_{\bar{f}}$ respectively. In order to cancel the quadratic anomaly we need the conditions

$$2N_c(1 + DD_{\text{ad}} - RR_{\text{ad}}) + (DD_{\text{f}} - RR_{\text{f}}) + (\bar{D}\bar{D}_{\text{f}} - \bar{R}\bar{R}_{\text{f}}) = 0 , \quad (\text{B.9})$$

$$2(1 + DD_{\text{ad}} - RR_{\text{ad}}) = (DD_{\text{d}} - RR_{\text{d}}) + (\bar{D}\bar{D}_{\text{d}} - \bar{R}\bar{R}_{\text{d}}) . \quad (\text{B.10})$$

For the linear anomaly we need to fulfill the condition

$$\sum_d \eta_d M_d - \sum_{\bar{d}} \bar{\eta}_{\bar{d}} \bar{M}_{\bar{d}} + \sum_f \rho_f m_f - \sum_{\bar{f}} \bar{\rho}_{\bar{f}} \bar{m}_{\bar{f}} = 0 , \quad (\text{B.11})$$

where we shifted all the masses by $\omega_3/2$ for convenience. An option is 2 RR and 1 DD adjoints (e.g. $\mathcal{N} = 4$). A related flavor-adjoint option requires one RR adjoint, not one DD adjoint, together with N RR fundamentals and N DD anti-fundamentals. Another possibility is the theory without adjoints, 1 DD determinant and 1 DD anti-determinant and fundamentals/anti-fundamentals satisfying

$$RR_{\text{f}} + \bar{R}\bar{R}_{\text{f}} - DD_{\text{f}} - \bar{D}\bar{D}_{\text{f}} = 2N_c . \quad (\text{B.12})$$

and the linear constraint

$$M - \bar{M} + \sum_{f=1}^{DD_{\text{f}}} m_f^{DD} - \sum_{f=1}^{RR_{\text{f}}} m_f^{RR} - \sum_{f=1}^{\bar{D}\bar{D}_{\text{f}}} \bar{m}_f^{DD} + \sum_{f=1}^{\bar{R}\bar{R}_{\text{f}}} \bar{m}_f^{RR} = 0 . \quad (\text{B.13})$$

The determinant-pair duality used in the main text is one concrete solution of these anomaly equations. In that example the determinant masses are identified with the total fundamental and anti-fundamental flavor masses, and the remaining spectator fugacities obey the balancing condition displayed in Section 5.1. The full duality statement and its JK-residue proof are given in the main text, so we do not repeat them in this appendix.

C Lower Dimensional Limits

C.1 Three-dimensional limit

Given the form of the determinant, the degeneration from $\text{Cyl} \times T^2$ to $\text{Cyl} \times S^1$ is obtained by taking $\text{Im } \tau \rightarrow +\infty$, equivalently $p = e^{2\pi i \tau} \rightarrow 0$. The corresponding 3d 1-loops then are, for the Dirichlet polarization,

$$Z_{1\text{-loop}}^{\text{CM}}|_{DD} = \prod_{(m_\varphi, m_y) \in \mathbb{Z}^2} \left(\omega m_\varphi + m_y - q_G \gamma_G^{(3d)} - (r-2) \gamma_R^{(3d)} \right). \quad (\text{C.1})$$

and, for the Robin polarization, the raw product reads

$$Z_{1\text{-loop}}^{\text{CM}}|_{RR} = \prod_{(m_\varphi, m_y) \in \mathbb{Z}^2} \left(\omega m_\varphi + m_y + q_G \gamma_G^{(3d)} + r \gamma_R^{(3d)} \right)^{-1}, \quad (\text{C.2})$$

with

$$\gamma_R^{(3d)} = \frac{1}{2}(\alpha_\varphi \omega + \alpha_y) \quad (\text{C.3})$$

and

$$\gamma_G^{(3d)} = \mathcal{A}_0 + \beta_y + \omega \beta_\varphi. \quad (\text{C.4})$$

The products in (C.1)–(C.2) are over the full lattice $\mathbb{Z}^2$ and are understood in the Shintani–Barnes sense, i.e. by analytic continuation of the Barnes multiple zeta function and its associated multiple Gamma functions.

Introduce now the shifts

$$\mu_D^{(3d)} := q_G \gamma_G^{(3d)} + (r-2) \gamma_R^{(3d)}, \quad \mu_R^{(3d)} := q_G \gamma_G^{(3d)} + r \gamma_R^{(3d)}. \quad (\text{C.5})$$

The Robin raw product is the reflected product with shift $-\mu_R^{(3d)}$. Its literal Shintani–Barnes evaluation is therefore obtained by applying the formula below at $\mu = -\mu_R^{(3d)}$ and then inverting the result. In the displayed RR block we instead use the same contact-term normalization as in the four-dimensional block convention and write the answer with argument $\mu_R^{(3d)}$. The two expressions differ by a holomorphic non-vanishing contact factor, so we denote the normalized equality by $\doteq$ rather than by $\stackrel{\text{SB}}{=}$. The canonical factorization of the $\mathbb{Z}^2$ product is

$$\prod_{(m_\varphi, m_y) \in \mathbb{Z}^2} (\omega m_\varphi + m_y - \mu) \stackrel{\text{SB}}{=} C_{3d}(\omega) \exp\left(-\frac{\pi i}{2} B_{2,2}(\mu | \omega, 1)\right) \Theta(\mu; \omega), \quad (\text{C.6})$$

where $C_{3d}(\omega)$ depends only on ω (and on the chosen normalization of the Barnes double Gamma) and is independent of μ . Here we defined the quadratic Bernoulli polynomial

$$B_{2,2}(u | \omega_1, \omega_2) = \frac{u^2}{\omega_1 \omega_2} - \frac{\omega_1 + \omega_2}{\omega_1 \omega_2} u + \frac{\omega_1^2 + 3\omega_1 \omega_2 + \omega_2^2}{6 \omega_1 \omega_2}. \quad (\text{C.7})$$

Applying (C.6) to (C.1)–(C.2) gives

$$Z_{1\text{-loop}}^{\text{CM}}|_{DD}^{(3d)} \stackrel{\text{SB}}{=} C_{3d}(\omega) \exp\left(-\frac{\pi i}{2} B_{2,2}(\mu_D^{(3d)} | \omega, 1)\right) \Theta(\mu_D^{(3d)}; \omega), \quad (\text{C.8})$$

$$Z_{1\text{-loop}}^{\text{CM}}|_{RR}^{(3d)} \doteq C_{3d}(\omega)^{-1} \exp\left(+\frac{\pi i}{2} B_{2,2}(\mu_R^{(3d)} | \omega, 1)\right) \Theta(\mu_R^{(3d)}; \omega)^{-1}. \quad (\text{C.9})$$

The prefactor $C_{3d}(\omega)$ cancels in determinant ratios and in polarization changes where only μ varies. With the normalization displayed above, the three-dimensional DD and RR blocks are inverse at the same argument; the reflected product differs by a holomorphic non-vanishing contact factor. The functional form of these 3d one-loop determinants agrees with the bulk $I \times T^2$ interval-index blocks of Ref. [10], once the contact-term normalization and boundary-matter sectors are matched. In this precise sense, the bulk one-loop sector of the construction of Ref. [10] is contained in the present framework as the $\text{Im } \tau \rightarrow +\infty$ degeneration of the four-dimensional cylinder kernel. It is also consistent with the lower-dimensional hyperbolic-block limits studied in Ref. [57].

C.2 Two-dimensional limit

The further degeneration from $\text{Cyl} \times S^1$ to Cyl is obtained by taking $\text{Im } \omega \rightarrow +\infty$, equivalently $q = e^{2\pi i \omega} \rightarrow 0$, or by setting $\alpha_y = \beta_y = 0$ and keeping only the corresponding zero modes in the infinite products. For instance, if we take the $\text{Im } \omega \rightarrow +\infty$ limit we obtain, for the Dirichlet polarization,

$$Z_{1\text{-loop}}^{\text{CM}}|_{DD} = \prod_{m_y \in \mathbb{Z}} \left(m_y - q_G \gamma_G^{(2d)} - (r-2) \gamma_R^{(2d)} \right). \quad (\text{C.10})$$

and, for the Robin polarization, the raw product reads

$$Z_{1\text{-loop}}^{\text{CM}}|_{RR} = \prod_{m_y \in \mathbb{Z}} \left(m_y + q_G \gamma_G^{(2d)} + r \gamma_R^{(2d)} \right)^{-1}, \quad (\text{C.11})$$

with

$$\gamma_R^{(2d)} = \frac{1}{2} \alpha_y \quad (\text{C.12})$$

and

$$\gamma_G^{(2d)} = \mathcal{A}_0 + \beta_y. \quad (\text{C.13})$$

Instead, if we just set all y -components to zero, we obtain for the Dirichlet polarization,

$$Z_{1\text{-loop}}^{\text{CM}}|_{DD} = \prod_{m_\varphi \in \mathbb{Z}} \left(m_\varphi - q_G \gamma_G^{(2d)} - (r-2) \gamma_R^{(2d)} \right). \quad (\text{C.14})$$

and, for the Robin polarization, the raw product reads

$$Z_{1\text{-loop}}^{\text{CM}}|_{RR} = \prod_{m_\varphi \in \mathbb{Z}} \left(m_\varphi + q_G \gamma_G^{(2d)} + r \gamma_R^{(2d)} \right)^{-1}, \quad (\text{C.15})$$

with

$$\gamma_R^{(2d)} = \frac{1}{2} \alpha_\varphi \quad (\text{C.16})$$

and

$$\gamma_G^{(2d)} = \frac{\mathcal{A}_0}{\omega} + \beta_\varphi = \hat{\mathcal{A}}_0 + \beta_\varphi. \quad (\text{C.17})$$

C.3 The $\mathbb{Z}$ -product and the cylinder determinant

The $\mathbb{Z}$ -products in (C.10)–(C.11) and in (C.14)–(C.15) are regularized by the Barnes one-Gamma function (equivalently, by the Hurwitz zeta prescription).

For $\mu \in \mathbb{C}$ one has the canonical Shintani–Barnes identity

$$\prod_{m \in \mathbb{Z}} (m - \mu) \stackrel{\text{SB}}{=} C_{2d} \exp(-\pi i B_{1,1}(\mu | 1)) (1 - e^{2\pi i \mu}), \quad (\text{C.18})$$

where C_{2d} is a μ -independent constant fixed by the normalization of Γ_1 (equivalently by the choice of zeta subtraction). In particular, (C.18) is equivalent to the familiar form $2 \sin(\pi \mu)$ up to a μ -independent factor, and exhibits explicitly the phase defined by the linear Bernoulli polynomial

$$B_{1,1}(u | \omega) = \frac{u}{\omega} - \frac{1}{2}. \quad (\text{C.19})$$

Define now the shifts

$$\mu_D^{(2d)} := q_G \gamma_G^{(2d)} + (r - 2) \gamma_R^{(2d)}, \quad \mu_R^{(2d)} := q_G \gamma_G^{(2d)} + r \gamma_R^{(2d)}. \quad (\text{C.20})$$

As in the three-dimensional limit, the Robin product is evaluated in the contact normalization compatible with $z_{\text{DD}}^{2d}(u) z_{\text{RR}}^{2d}(-u) = 1$. Applying (C.18) yields, for the $m_y \in \mathbb{Z}$ reduction (C.10)–(C.11),

$$Z_{1\text{-loop}}^{\text{CM}}|_{DD}^{(2d)} \stackrel{\text{SB}}{=} C_{2d} \exp(-\pi i B_{1,1}(\mu_D^{(2d)} | 1)) (1 - e^{2\pi i \mu_D^{(2d)}}), \quad (\text{C.21})$$

$$Z_{1\text{-loop}}^{\text{CM}}|_{RR}^{(2d)} \stackrel{\text{SB}}{=} C_{2d}^{-1} \exp(+\pi i B_{1,1}(\mu_R^{(2d)} | 1)) (1 - e^{2\pi i \mu_R^{(2d)}})^{-1}. \quad (\text{C.22})$$

The same formulas apply verbatim to the alternative reduction (C.14)–(C.15), with $\mu_D^{(2d)}, \mu_R^{(2d)}$ computed using the corresponding $(\gamma_R^{(2d)}, \gamma_G^{(2d)})$ in that branch. As in three dimensions, the constants $C_{2d}^{\pm 1}$ cancel in determinant ratios.

References

- [1] Kentaro Hori and Mauricio Romo. Exact Results In Two-Dimensional (2,2) Supersymmetric Gauge Theories With Boundary. 8 2013. [arXiv:1308.2438](#).
- [2] Daigo Honda and Takuya Okuda. Exact results for boundaries and domain walls in 2d supersymmetric theories. *JHEP*, 09:140, 2015. [arXiv:1308.2217](#), [doi:10.1007/JHEP09\(2015\)140](#).
- [3] Vasily Pestun. Localization of gauge theory on a four-sphere and supersymmetric Wilson loops. *Commun. Math. Phys.*, 313:71–129, 2012. [arXiv:0712.2824](#), [doi:10.1007/s00220-012-1485-0](#).
- [4] Guido Festuccia and Nathan Seiberg. Rigid Supersymmetric Theories in Curved Super-space. *JHEP*, 06:114, 2011. [arXiv:1105.0689](#), [doi:10.1007/JHEP06\(2011\)114](#).

- [5] Thomas T. Dumitrescu, Guido Festuccia, and Nathan Seiberg. Exploring Curved Superspace. *JHEP*, 08:141, 2012. [arXiv:1205.1115](#), [doi:10.1007/JHEP08\(2012\)141](#).
- [6] Sara Pasquetti. Factorisation of $N = 2$ Theories on the Squashed 3-Sphere. *JHEP*, 04:120, 2012. [arXiv:1111.6905](#), [doi:10.1007/JHEP04\(2012\)120](#).
- [7] Christopher Beem, Tudor Dimofte, and Sara Pasquetti. Holomorphic Blocks in Three Dimensions. *JHEP*, 12:177, 2014. [arXiv:1211.1986](#), [doi:10.1007/JHEP12\(2014\)177](#).
- [8] Pietro Longhi, Fabrizio Nieri, and Antonio Pittelli. Localization of 4d $\mathcal{N} = 1$ theories on $\mathbb{D}^2 \times \mathbb{T}^2$. *JHEP*, 12:147, 2019. [arXiv:1906.02051](#), [doi:10.1007/JHEP12\(2019\)147](#).
- [9] Fabrizio Nieri and Sara Pasquetti. Factorisation and holomorphic blocks in 4d. *JHEP*, 11:155, 2015. [arXiv:1507.00261](#), [doi:10.1007/JHEP11\(2015\)155](#).
- [10] Katsuyuki Sugiyama and Yutaka Yoshida. Supersymmetric indices on $I \times T^2$, elliptic genera and dualities with boundaries. *Nucl. Phys. B*, 960:115168, 2020. [arXiv:2007.07664](#), [doi:10.1016/j.nuclphysb.2020.115168](#).
- [11] Yutaka Yoshida and Katsuyuki Sugiyama. Localization of three-dimensional $\mathcal{N} = 2$ supersymmetric theories on $S^1 \times D^2$. *PTEP*, 2020(11):113B02, 2020. [arXiv:1409.6713](#), [doi:10.1093/ptep/ptaa136](#).
- [12] Boan Zhao, Panos Betzios, and Paul Luis Roehl. Factorizations of 3d Interval Partition Functions. 9 2025. [arXiv:2509.23902](#).
- [13] Mykola Dedushenko. Gluing. Part I. Integrals and symmetries. *JHEP*, 04:175, 2020. [arXiv:1807.04274](#), [doi:10.1007/JHEP04\(2020\)175](#).
- [14] Mykola Dedushenko. Gluing II: boundary localization and gluing formulas. *Lett. Math. Phys.*, 111(1):18, 2021. [arXiv:1807.04278](#), [doi:10.1007/s11005-021-01355-8](#).
- [15] Mykola Dedushenko and Nikita Nekrasov. Interfaces and quantum algebras, I: Stable envelopes. *J. Geom. Phys.*, 194:104991, 2023. [arXiv:2109.10941](#), [doi:10.1016/j.geomphys.2023.104991](#).
- [16] Mykola Dedushenko and Mikhail Litvinov. Supersymmetric reduction of 3D theories on a line segment. *Phys. Rev. D*, 108(4):045016, 2023. [arXiv:2212.07455](#), [doi:10.1103/PhysRevD.108.045016](#).
- [17] Mykola Dedushenko and Nikita Nekrasov. Interfaces and Quantum Algebras, II: Cigar Partition Function. 6 2023. [arXiv:2306.16434](#).
- [18] Mikhail Litvinov, Sergey Alekseev, and Mykola Dedushenko. Chiral life on a slab. *Lett. Math. Phys.*, 116(1):7, 2026. [arXiv:2301.00038](#), [doi:10.1007/s11005-025-02040-w](#).

- [19] Vyacheslav P. Spiridonov. On the elliptic beta function. *Russ. Math. Surveys*, 56:185–186, 2001. [arXiv:math/0011001](#), [doi:10.1070/RM2001v056n01ABEH000371](#).
- [20] Eric M. Rains. Transformations of elliptic hypergeometric integrals. *Annals of Mathematics*, 171:169–243, 2010. [arXiv:math/0309252](#), [doi:10.4007/annals.2010.171.169](#).
- [21] Lea E. Bottini, Chiung Hwang, Sara Pasquetti, and Matteo Sacchi. 4d S-duality wall and $SL(2, \mathbb{Z})$ relations. *JHEP*, 03:035, 2022. [arXiv:2110.08001](#), [doi:10.1007/JHEP03\(2022\)035](#).
- [22] Sergio Benvenuti, Riccardo Comi, and Sara Pasquetti. Star-triangle dualities and supersymmetric improved bifundamentals. *JHEP*, 09:171, 2025. [arXiv:2410.19049](#), [doi:10.1007/JHEP09\(2025\)171](#).
- [23] Simone Giacomelli, Chiung Hwang, Fabio Marino, Sara Pasquetti, and Matteo Sacchi. Probing bad theories with the dualization algorithm. Part I. *JHEP*, 04:008, 2024. [arXiv:2309.05326](#), [doi:10.1007/JHEP04\(2024\)008](#).
- [24] Simone Giacomelli, Chiung Hwang, Fabio Marino, Sara Pasquetti, and Matteo Sacchi. Probing bad theories with the dualization algorithm. Part II. *JHEP*, 07:165, 2024. [arXiv:2401.14456](#), [doi:10.1007/JHEP07\(2024\)165](#).
- [25] Daniele Iannotti and Antonio Pittelli. Twisted Index on Hyperbolic Four-Manifolds. *Lett. Math. Phys.*, 114(2):39, 2024. [arXiv:2307.11634](#), [doi:10.1007/s11005-024-01788-x](#).
- [26] Tudor Dimofte, Davide Gaiotto, and Natalie M. Paquette. Dual boundary conditions in 3d SCFT's. *JHEP*, 05:060, 2018. [arXiv:1712.07654](#), [doi:10.1007/JHEP05\(2018\)060](#).
- [27] Mykola Dedushenko. On the 4d/3d/2d view of the SCFT/VOA correspondence. 12 2023. [arXiv:2312.17747](#).
- [28] Giovanni Felder and Alexander Varchenko. The elliptic gamma function and $SL(3, F) \ltimes F^3$. *Advances in Mathematics*, 156(1):44–76, 2000. [arXiv:math/9907061](#), [doi:10.1006/aima.2000.1942](#).
- [29] Atsushi Narukawa. The modular properties and the integral representations of the multiple elliptic gamma functions. *Advances in Mathematics*, 189(2):247–267, 2004. [arXiv:math/0306164](#), [doi:10.1016/j.aim.2003.11.009](#).
- [30] Eduardo Friedman and Simon Ruijsenaars. Shintani–barnes zeta and gamma functions. *Advances in Mathematics*, 187(2):362–395, 2004. [doi:10.1016/j.aim.2003.07.020](#).
- [31] Nikita A. Nekrasov and Samson L. Shatashvili. Supersymmetric vacua and Bethe ansatz. *Nucl. Phys. B Proc. Suppl.*, 192-193:91–112, 2009. [arXiv:0901.4744](#), [doi:10.1016/j.nuclphysbps.2009.07.047](#).

- [32] Lisa C. Jeffrey and Frances C. Kirwan. Localization for nonabelian group actions. *Topology*, 34:291–327, 1995. [arXiv:alg-geom/9307001](#), [doi:10.1016/0040-9383\(94\)00028-J](#).
- [33] Francesco Benini, Richard Eager, Kentaro Hori, and Yuji Tachikawa. Elliptic genera of two-dimensional $\mathcal{N} = 2$ gauge theories with rank-one gauge groups. *Lett. Math. Phys.*, 104:465–493, 2014. [arXiv:1305.0533](#), [doi:10.1007/s11005-013-0673-y](#).
- [34] Francesco Benini, Richard Eager, Kentaro Hori, and Yuji Tachikawa. Elliptic Genera of 2d $\mathcal{N} = 2$ Gauge Theories. *Commun. Math. Phys.*, 333(3):1241–1286, 2015. [arXiv:1308.4896](#), [doi:10.1007/s00220-014-2210-y](#).
- [35] Cyril Closset, Heeyeon Kim, and Brian Willett. Supersymmetric partition functions and the three-dimensional A-twist. *JHEP*, 03:074, 2017. [arXiv:1701.03171](#), [doi:10.1007/JHEP03\(2017\)074](#).
- [36] Nathan Seiberg. Electric–magnetic duality in supersymmetric non-abelian gauge theories. *Nucl. Phys. B*, 435:129–146, 1995. [arXiv:hep-th/9411149](#), [doi:10.1016/0550-3213\(94\)00023-8](#).
- [37] Edward Witten. Phases of $n=2$ theories in two dimensions. *Nuclear Physics B*, 403:159–222, 1993. [arXiv:hep-th/9301042](#), [doi:10.1016/0550-3213\(93\)90033-L](#).
- [38] Seyed Morteza Hosseini, Anton Nedelin, and Alberto Zaffaroni. The cardy limit of the topologically twisted index and black strings in AdS_5 . *Journal of High Energy Physics*, 04:014, 2017. [arXiv:1611.09374](#), [doi:10.1007/JHEP04\(2017\)014](#).
- [39] Seyed Morteza Hosseini, Kiril Hristov, and Alberto Zaffaroni. Microstates of rotating AdS_5 strings. *Journal of High Energy Physics*, 11:090, 2019. [arXiv:1909.08000](#), [doi:10.1007/JHEP11\(2019\)090](#).
- [40] Junho Hong, James T. Liu, and Leopoldo A. Pando Zayas. The topologically twisted index of $\mathfrak{su}(4)$ super-yang–mills on $t^2 \times s^2$ and the elliptic genus. *Journal of High Energy Physics*, 07:018, 2018. [arXiv:1804.04592](#), [doi:10.1007/JHEP07\(2018\)018](#).
- [41] Matteo Inglese, Dario Martelli, and Antonio Pittelli. The spindle index from localization. *J. Phys. A*, 57(8):085401, 2024. [arXiv:2303.14199](#), [doi:10.1088/1751-8121/ad2225](#).
- [42] Edoardo Colombo, Seyed Morteza Hosseini, Dario Martelli, Antonio Pittelli, and Alberto Zaffaroni. Microstates of Accelerating and Supersymmetric AdS_4 Black Holes from the Spindle Index. *Phys. Rev. Lett.*, 133(3):031603, 2024. [arXiv:2404.07173](#), [doi:10.1103/PhysRevLett.133.031603](#).
- [43] Curtis G. Jr. Callan and Jeffrey A. Harvey. Anomalies and Fermion Zero Modes on Strings and Domain Walls. *Nucl. Phys. B*, 250:427–436, 1985. [doi:10.1016/0550-3213\(85\)90489-4](#).

- [44] Davide Gaiotto and Edward Witten. Supersymmetric Boundary Conditions in N=4 Super Yang-Mills Theory. *J. Statist. Phys.*, 135:789–855, 2009. [arXiv:0804.2902](#), [doi:10.1007/s10955-009-9687-3](#).
- [45] Michael R. Douglas. D-branes, categories and N=1 supersymmetry. *Journal of Mathematical Physics*, 42:2818–2843, 2001. [arXiv:hep-th/0011017](#), [doi:10.1063/1.1374448](#).
- [46] Anton Kapustin and Yi Li. D-branes in landau–ginzburg models and algebraic geometry. *Journal of High Energy Physics*, 12:005, 2003. [arXiv:hep-th/0210296](#), [doi:10.1088/1126-6708/2003/12/005](#).
- [47] Manfred Herbst, Kentaro Hori, and David Page. Phases of N=2 theories in 1+1 dimensions with boundary. 3 2008. [arXiv:0803.2045](#).
- [48] Sergio Cecotti and Cumrun Vafa. Topological–anti-topological fusion. *Nuclear Physics B*, 367(2):359–461, 1991. [doi:10.1016/0550-3213\(91\)90021-0](#).
- [49] Sergio Cecotti, Davide Gaiotto, and Cumrun Vafa. tt^* geometry in 3 and 4 dimensions. *JHEP*, 05:055, 2014. [arXiv:1312.1008](#), [doi:10.1007/JHEP05\(2014\)055](#).
- [50] Gregory W. Moore and Nathan Seiberg. Classical and quantum conformal field theory. *Communications in Mathematical Physics*, 123:177–254, 1989. [doi:10.1007/BF01238857](#).
- [51] Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert. TFT construction of RCFT correlators i: partition functions. *Nuclear Physics B*, 646:353–497, 2002. [arXiv:hep-th/0204148](#), [doi:10.1016/S0550-3213\(02\)00744-7](#).
- [52] Federico Faedo, Nicolo Petri, and Alessia Segati. Holographic reconstruction for defect CFTs from $AdS_p \times S^q$ spacetimes. 6 2026. [arXiv:2606.03106](#).
- [53] Giuseppe Dibitetto and Nicolo Petri. $AdS_D \times I$ solutions in axio-dilaton gravity. 6 2026. [arXiv:2606.04933](#).
- [54] Giuseppe Dibitetto and Nicolo Petri. AdS_9 solutions in type II supergravities. 6 2026. [arXiv:2606.04938](#).
- [55] Seyed Morteza Hosseini, Kiril Hristov, and Alberto Zaffaroni. Gluing gravitational blocks for AdS black holes. *JHEP*, 12:168, 2019. [arXiv:1909.10550](#), [doi:10.1007/JHEP12\(2019\)168](#).
- [56] Francesco Benini and Nikolay Bobev. Two-dimensional SCFTs from wrapped branes and c-extremization. *JHEP*, 06:005, 2013. [arXiv:1302.4451](#), [doi:10.1007/JHEP06\(2013\)005](#).
- [57] Antonio Pittelli. Supersymmetric localization of refined chiral multiplets on topologically twisted $H^2 \times S^1$. *Phys. Lett. B*, 801:135154, 2020. [arXiv:1812.11151](#), [doi:10.1016/j.physletb.2019.135154](#).

- [58] Matteo Inglese, Dario Martelli, and Antonio Pittelli. Supersymmetry and Localization on Three-Dimensional Orbifolds. 12 2023. [arXiv:2312.17086](#).
- [59] Antonio Pittelli. Orbifold Indices in Four Dimensions. 3 2024. [arXiv:2403.12318](#).
- [60] Martin F. Sohnius and Peter C. West. An alternative minimal off-shell version of N=1 supergravity. *Phys. Lett. B*, 105:353–357, 1981. [doi:10.1016/0370-2693\(81\)90778-4](#).
- [61] Guido Festuccia, Anastasios Gorantis, Antonio Pittelli, Konstantina Polydorou, and Lorenzo Ruggeri. Cohomological localization of N=2 gauge theories with matter. *JHEP*, 09:133, 2020. [arXiv:2005.12944](#), [doi:10.1007/JHEP09\(2020\)133](#).
- [62] Rodolfo Panerai, Antonio Pittelli, and Konstantina Polydorou. Topological Correlators and Surface Defects from Equivariant Cohomology. *JHEP*, 09:185, 2020. [arXiv:2006.06692](#), [doi:10.1007/JHEP09\(2020\)185](#).
- [63] Mykola Dedushenko, Silviu S. Pufu, and Ran Yacoby. A one-dimensional theory for Higgs branch operators. *JHEP*, 03:138, 2018. [arXiv:1610.00740](#), [doi:10.1007/JHEP03\(2018\)138](#).
- [64] E. W. Barnes. On the theory of the multiple gamma function. *Transactions of the Cambridge Philosophical Society*, 19:374–425, 1904.
- [65] Takuro Shintani. On a kronecker limit formula for real quadratic fields. *Journal of the Faculty of Science, University of Tokyo. Section IA, Mathematics*, 24:167–199, 1977.
- [66] Fabrizio Nieri, Sara Pasquetti, and Filippo Passerini. 3d and 5d Gauge Theory Partition Functions as q -deformed CFT Correlators. *Lett. Math. Phys.*, 105(1):109–148, 2015. [arXiv:1303.2626](#), [doi:10.1007/s11005-014-0727-9](#).
- [67] Fabrizio Nieri, Sara Pasquetti, Filippo Passerini, and Alessandro Torrielli. 5D partition functions, q -Virasoro systems and integrable spin-chains. *JHEP*, 12:040, 2014. [arXiv:1312.1294](#), [doi:10.1007/JHEP12\(2014\)040](#).
- [68] Cyril Closset and Itamar Shamir. The N=1 Chiral Multiplet on $T^2 \times S^2$ and Supersymmetric Localization. 11 2013. [arXiv:1311.2430](#).
- [69] Igor R. Klebanov and Edward Witten. Superconformal field theory on threebranes at a Calabi-Yau singularity. *Nucl. Phys. B*, 536:199–218, 1998. [arXiv:hep-th/9807080](#), [doi:10.1016/S0550-3213\(98\)00654-3](#).
- [70] Juan Martin Maldacena. The Large N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998. [arXiv:hep-th/9711200](#), [doi:10.4310/ATMP.1998.v2.n2.a1](#).