Patient Record — ID 1734

Age	29	**Sex**	F
Race/Ethnicity	Caucasian	**Insurance**	Blue Cross Blue Shield (PPO)
Total Encounters	8	**Generation Seed**	6497

Primary Diagnoses

• Disseminated intravascular coagulation
• Physiologic dyspnea of pregnancy
• Traumatic hemorrhagic shock with acute coagulopathy
• Bilateral adrenal hemorrhage (Waterhouse-Friderichsen syndrome)
• Pelvic Girdle Pain (Pregnancy-related)
• Acute fatty liver of pregnancy
• Umbilical cord compression causing recurrent variable decelerations

Comorbidities

• Pregnancy
• Acute posthemorrhagic anemia

Patient Profile

Sex	F
Age at First Encounter	29
Race/Ethnicity	Caucasian
Insurance	Blue Cross Blue Shield (PPO)
Occupation	Elementary school teacher
Smoking Status	Never smoker
Alcohol Use	Rare social use, none during pregnancies

Allergies:

• NKDA

Chronic Conditions:

• Chronic hypertension
• History of postpartum hemorrhage

Home Medications:

• Labetalol - 200 mg PO twice daily (adherence reported)
• Prenatal vitamin with iron - 1 tablet PO daily

Surgical History:

• Manual removal of retained placenta (age 27)
• Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

Family History:

• Mother with chronic hypertension
• Maternal aunt with severe preeclampsia
• Previous child with transient hypoglycemia (resolved)
• No known bleeding disorders

Encounters (8)

Encounter 1 — 2020-01-15 — Obstetrics & Gynecology

Encounter ID	7167	**Type**	icu
Attending	Dr. Sarah Chen	**Order**	0
Department	Obstetrics & Gynecology	**Gen. Method**	template

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Generalized weakness and excessive vaginal bleeding 12 hours after delivering first child

HPI

12 hours after delivering her first child. The delivery was complicated by a retained placenta, which required manual removal. Her prenatal course was unremarkable

PMH (modified)

Active Problem List:
- Chronic hypertension
- History of postpartum hemorrhage

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

Home Medications:
- Labetalol 200 mg PO twice daily (adherence reported)
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY (modified)

Smoking: Never smoker. Alcohol: Rare social use, none during pregnancies. Occupation: Elementary school teacher.

VITALS

Blood pressure is 90/60 mmHg, heart rate is 120 beats per minute, and temperature is 37.2°C (99.0°F)

PHYSICAL_EXAM

She appears pale and diaphoretic. Pelvic examination reveals moderate vaginal bleeding.

LABS

- Hemoglobin: 8 g/dL (Normal: 12-16 g/dL)
- Platelet count: 50,000/mm³ (Normal: 150,000-400,000/mm³)
- Prothrombin time (PT): 18 seconds (Normal: 12-14 seconds)
- Partial thromboplastin time (PTT): 45 seconds (Normal: 25-35 seconds)
- Fibrinogen: 150 mg/dL (Normal: 200-400 mg/dL)
- D-dimer: Elevated

Encounter 2 — 2022-01-15 — Obstetrics & Gynecology

Encounter ID	7163	**Type**	outpatient
Attending	Dr. Maria Rodriguez	**Order**	1
Department	Obstetrics & Gynecology	**Gen. Method**	hybrid

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Shortness of breath, especially when lying down, during second pregnancy

HPI (modified)

This 31-year-old G2P1 with chronic hypertension and prior postpartum hemorrhage presents for second pregnancy with complaints of shortness of breath, especially when lying down. She denies any cough, fever, or wheezing.

PMH (modified)

Active Problem List:
- Disseminated intravascular coagulation (diagnosed 2020-01-15)
- Chronic hypertension
- History of postpartum hemorrhage

She has had an uncomplicated prenatal course thus far.

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

Home Medications:
- Labetalol 200 mg PO twice daily (adherence reported)
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY (modified)

Smoking: Never smoker. Alcohol: Rare social use, none during pregnancies. Occupation: Elementary school teacher.

VITALS

On examination, her respiratory rate is 18 breaths per minute, and oxygen saturation is 97% on room air.

PHYSICAL_EXAM

Lung auscultation reveals clear breath sounds bilaterally. Her abdomen is gravid, and fundal height is appropriate for gestational age.

OTHER_STUDIES

A spirometry test yields the following results:

- Forced vital capacity (FVC): 4.0 L (normal)
- Forced expiratory volume in 1 second (FEV1): 3.2 L (normal)
- FEV1/FVC ratio: 80% (normal)
- Residual volume (RV): Decreased
- Functional residual capacity (FRC): Decreased

Encounter 3 — 2022-04-15 — Obstetrics & Gynecology

Encounter ID	7166	**Type**	outpatient
Attending	Dr. Maria Rodriguez	**Order**	2
Department	Obstetrics & Gynecology	**Gen. Method**	hybrid

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Progressive pain over pubic bone and bilateral groin areas worsened by walking or turning in bed

HPI (modified)

This 31-year-old with chronic hypertension and postpartum hemorrhage returns for prenatal care following her January visit. She reports that the pain is worse when walking, climbing stairs, or turning in bed. She denies leakage of fluid, vaginal bleeding, or contractions.

PMH (modified)

Active Problem List:
- Disseminated intravascular coagulation (diagnosed 2020-01-15)
- Physiologic dyspnea of pregnancy (diagnosed 2022-01-15)
- Chronic hypertension
- History of postpartum hemorrhage

Her pregnancy has been uncomplicated.

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

Home Medications:
- Labetalol 200 mg PO twice daily (adherence reported)
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY (modified)

Smoking: Never smoker. Alcohol: Rare social use, none during pregnancies. Occupation: Elementary school teacher.

VITALS

Her vital signs are within normal limits.

PHYSICAL_EXAM

On physical examination, she has tenderness over the pubic symphysis and pain with passive abduction of her hips. There is no erythema or edema over the joints, and her skin is intact.

LABS

Urinalysis is negative for protein, glucose, and infection.

Encounter 4 — 2022-07-15 — Obstetrics & Gynecology

Encounter ID	7165	**Type**	outpatient
Attending	Dr. James Park	**Order**	3
Department	Obstetrics & Gynecology	**Gen. Method**	hybrid

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Routine prenatal visit with reports of decreased fetal movements over past two days

HPI (modified)

The patient is a 31-year-old G2P1 female with chronic hypertension and a history of postpartum hemorrhage, now returning for prenatal care following prior visits for physiologic dyspnea and pelvic girdle pain. She has a history of chronic hypertension, which has been poorly controlled during pregnancy despite adherence to labetalol therapy. She reports decreased fetal movements over the past two days. Her prenatal course has been otherwise unremarkable.

PMH (modified)

Active Problem List:
- Disseminated intravascular coagulation (diagnosed 2020-01-15)
- Physiologic dyspnea of pregnancy (diagnosed 2022-01-15)
- Pelvic Girdle Pain (Pregnancy-related) (diagnosed 2022-04-15)
- Chronic hypertension
- History of postpartum hemorrhage

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

She takes labetalol therapy.

Home Medications:
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY (modified)

Smoking: Never smoker. Alcohol: Rare social use, none during pregnancies. Occupation: Elementary school teacher.

VITALS

Her blood pressure is 152/96 mmHg

PHYSICAL_EXAM

Fundal height is measuring 30 cm, which is less than expected for gestational age

IMAGING

An ultrasound is performed, which shows an estimated fetal weight in the 8th percentile for gestational age, but with a normal head circumference. Doppler flow studies of the umbilical artery reveal increased resistance to blood flow.

Encounter 5 — 2022-09-15 — Emergency Medicine

Encounter ID	7168	**Type**	ed
Attending	Dr. Aisha Patel	**Order**	4
Department	Emergency Medicine	**Gen. Method**	hybrid

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Persistent nausea, vomiting, and severe epigastric pain radiating to right upper quadrant

HPI (modified)

This 31-year-old female with chronic hypertension, postpartum hemorrhage, and recent bilateral adrenal hemorrhage presents to the emergency department with a 24-hour history of persistent nausea, vomiting, and severe epigastric pain. She reports that the pain radiates to her right upper quadrant.

PMH (modified)

Active Problem List:
- Disseminated intravascular coagulation (diagnosed 2020-01-15)
- Physiologic dyspnea of pregnancy (diagnosed 2022-01-15)
- Pelvic Girdle Pain (Pregnancy-related) (diagnosed 2022-04-15)
- Bilateral adrenal hemorrhage (Waterhouse-Friderichsen syndrome) (diagnosed 2022-07-15)
- Chronic hypertension
- History of postpartum hemorrhage

Her pregnancy has been uncomplicated until now.

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

Home Medications:
- Labetalol 200 mg PO twice daily (adherence reported)
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY (modified)

Smoking: Never smoker. Alcohol: Rare social use, none during pregnancies. Occupation: Elementary school teacher.

VITALS

On examination, her blood pressure is 145/95 mmHg, heart rate is 102 beats per minute, and she is afebrile.

PHYSICAL_EXAM

Her skin and sclera are anicteric. Abdominal examination reveals right upper quadrant tenderness without rebound or guarding.

LABS

Laboratory studies are as follows:

- Hemoglobin: 12.1 g/dL
- Platelet count: 155,000/mm^3
- Aspartate aminotransferase (AST): 250 U/L
- Alanine aminotransferase (ALT): 230 U/L
- Total bilirubin: 1.2 mg/dL
- Alkaline phosphatase: 120 U/L
- Serum glucose: 45 mg/dL
- Prothrombin time (PT): 16 seconds (control 12 seconds)
- Partial thromboplastin time (PTT): 40 seconds (control 30 seconds)
- Fibrinogen: 150 mg/dL

OTHER_STUDIES

Fetal monitoring shows a reactive fetal heart tracing.

PLAN

The obstetric team is consulted.

Encounter 6 — 2022-11-15 — Obstetrics & Gynecology

Encounter ID	7169	**Type**	inpatient
Attending	Dr. Sarah Chen	**Order**	5
Department	Obstetrics & Gynecology	**Gen. Method**	hybrid

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Sudden onset irregular abdominal pain at term gestation

HPI (modified)

The patient is a 31-year-old G2P1 with chronic hypertension and prior postpartum hemorrhage who presents to the labor and delivery unit at term following recent acute fatty liver of pregnancy and adrenal hemorrhage. She reports a 1-hour history of sudden onset, irregular abdominal pain. She denies any vaginal bleeding or leakage of fluid. Her pregnancy has been uncomplicated, and she has received regular prenatal care.

PMH (modified)

Active Problem List:
- Disseminated intravascular coagulation (diagnosed 2020-01-15)
- Physiologic dyspnea of pregnancy (diagnosed 2022-01-15)
- Pelvic Girdle Pain (Pregnancy-related) (diagnosed 2022-04-15)
- Bilateral adrenal hemorrhage (Waterhouse-Friderichsen syndrome) (diagnosed 2022-07-15)
- Acute fatty liver of pregnancy (diagnosed 2022-09-15)
- Chronic hypertension
- History of postpartum hemorrhage

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

Home Medications:
- Labetalol 200 mg PO twice daily (adherence reported)
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY (modified)

Smoking: Never smoker. Alcohol: Rare social use, none during pregnancies. Occupation: Elementary school teacher.

VITALS

On examination, her blood pressure is 120/75 mmHg, heart rate is 88 beats per minute, and temperature is 37.0°C (98.6°F)

PHYSICAL_EXAM

Fetal heart monitoring shows a baseline fetal heart rate of 140 beats per minute with recurrent variable decelerations that decrease to 90 beats per minute with >50% of uterine contractions. The decelerations are noted to have a rapid descent and return to baseline.
There is no evidence of maternal fever, tachycardia, or uterine tenderness.

PLAN

The patient is placed in the left lateral decubitus position, and oxygen is administered via face mask. Despite these measures, the fetal heart tracing does not improve.

Encounter 7 — 2022-12-15 — Obstetrics & Gynecology

Encounter ID	7162	**Type**	icu
Attending	Dr. James Park	**Order**	6
Department	Obstetrics & Gynecology	**Gen. Method**	hybrid

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Sudden-onset severe abdominal pain and vaginal bleeding with significantly decreased fetal movements

HPI (modified)

This 31-year-old G2P1 female with chronic hypertension has a history of postpartum hemorrhage complicated by disseminated intravascular coagulation and antepartum complications including adrenal hemorrhage. She presents to the emergency department with sudden-onset severe abdominal pain and vaginal bleeding that began 3 hours ago. She reports that the pain is constant and not associated with uterine contractions. Fetal movements have significantly decreased since the onset of symptoms. Shortly after arrival, the patient develops oozing of blood from her intravenous catheter site and persistent vaginal bleeding.

PMH (modified)

Active Problem List:
- Disseminated intravascular coagulation (diagnosed 2020-01-15)
- Physiologic dyspnea of pregnancy (diagnosed 2022-01-15)
- Pelvic Girdle Pain (Pregnancy-related) (diagnosed 2022-04-15)
- Bilateral adrenal hemorrhage (Waterhouse-Friderichsen syndrome) (diagnosed 2022-07-15)
- Acute fatty liver of pregnancy (diagnosed 2022-09-15)
- Umbilical cord compression causing recurrent variable decelerations (diagnosed 2022-11-15)
- Chronic hypertension
- History of postpartum hemorrhage

Her past pregnancies were uneventful.

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

Home Medications:
- Labetalol 200 mg PO twice daily (adherence reported)
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY

She does not smoke or use drugs.

VITALS

Blood pressure is 90/58 mm Hg, pulse is 118/min

PHYSICAL_EXAM

On examination, her uterus is firm and tender to palpation, and there is moderate vaginal bleeding. She appears pale and diaphoretic. There is no external trauma.

LABS

Hemoglobin 8.3 g/dL
Platelet count 44,000/mm^3
Prothrombin time (PT) 24 seconds
Activated PTT 60 seconds
Fibrinogen 90 mg/dL
Fibrin degradation products Elevated

Encounter 8 — 2023-07-15 — Emergency Medicine

Encounter ID	7164	**Type**	ed
Attending	Dr. Aisha Patel	**Order**	7
Department	Emergency Medicine	**Gen. Method**	hybrid

Structured EHR Sections:

CHIEF_COMPLAINT (modified)

Dizziness and fatigue following motor vehicle collision

HPI (modified)

The patient is a 32-year-old female with chronic hypertension on labetalol and a history of postpartum hemorrhage complicated by disseminated intravascular coagulation, who presents after a motor vehicle collision. She reports that she was the restrained driver in a head-on collision at moderate speed. Airbags were deployed, and she did not lose consciousness. She complains of mild abdominal pain but no other specific symptoms.

PMH (modified)

Active Problem List:
- Disseminated intravascular coagulation (diagnosed 2020-01-15)
- Physiologic dyspnea of pregnancy (diagnosed 2022-01-15)
- Pelvic Girdle Pain (Pregnancy-related) (diagnosed 2022-04-15)
- Bilateral adrenal hemorrhage (Waterhouse-Friderichsen syndrome) (diagnosed 2022-07-15)
- Acute fatty liver of pregnancy (diagnosed 2022-09-15)
- Umbilical cord compression causing recurrent variable decelerations (diagnosed 2022-11-15)
- Chronic hypertension
- History of postpartum hemorrhage

PSH (modified)

- Manual removal of retained placenta (age 27)
- Cesarean section (implied by possible repeat surgeries, but not explicitly stated - will leave as manual removal only)

MEDICATIONS (modified)

Home Medications:
- Labetalol 200 mg PO twice daily (adherence reported)
- Prenatal vitamin with iron 1 tablet PO daily

ALLERGIES (modified)

NKDA

FAMILY_HISTORY (modified)

- Mother with chronic hypertension
- Maternal aunt with severe preeclampsia
- Previous child with transient hypoglycemia (resolved)
- No known bleeding disorders

SOCIAL_HISTORY (modified)

Smoking: Never smoker. Alcohol: Rare social use, none during pregnancies. Occupation: Elementary school teacher.

VITALS

Her blood pressure is 90/60 mmHg, heart rate is 120 beats per minute, respiratory rate is 22 breaths per minute, and temperature is 37.0°C (98.6°F). The patient's blood pressure drops to 85/55 mmHg despite receiving 2 liters of normal saline.

PHYSICAL_EXAM

On examination, she appears pale and anxious. Her skin is cool and clammy. Abdominal examination reveals mild tenderness in the left upper quadrant with no rebound or guarding. Bowel sounds are normal. Pelvic examination is unremarkable. No external injuries are noted.

LABS

Laboratory studies show:

- Hemoglobin: 9.5 g/dL (normal: 12-16 g/dL)
- Hematocrit: 28% (normal: 36-46%)
- Platelets: 250,000/mm³ (normal: 150,000-400,000/mm³)
- White blood cells: 12,000/mm³ (normal: 4,500-11,000/mm³)
- Serum lactate: 4 mmol/L (normal: 0.5-2.2 mmol/L)

IMAGING

Focused Assessment with Sonography for Trauma (FAST) exam is negative for free fluid. Chest X-ray is unremarkable.