

Today's topic

- Derivatives of vector valued functions (Sec 10.7)

Practice 1: Find the maximum rate of change of f and direction in which it occurs.

$$f(x, y, z) = \log(xy^2z^2), \quad \text{at } (1, -2, -3)$$

Space Curves: We have discussed earlier. We also call them parametric curves

$$\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k} \quad a \leq t \leq b$$

Examples: circle, ellipse, parabola, helix, straight line

DERIVATIVES

The **derivative** \mathbf{r}' of a vector function \mathbf{r} is defined in much the same way as for real-valued functions:


3

$$\frac{d\mathbf{r}}{dt} = \mathbf{r}'(t) = \lim_{h \rightarrow 0} \frac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$$

the vector $\mathbf{r}'(t)$ is called the **tangent vector** to the curve

unit tangent vector

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}$$



The following theorem gives us a convenient method for computing the derivative of a vector function \mathbf{r} : just differentiate each component of \mathbf{r} .

4 THEOREM If $\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle = f(t) \mathbf{i} + g(t) \mathbf{j} + h(t) \mathbf{k}$, where f , g , and h are differentiable functions, then

$$\mathbf{r}'(t) = \langle f'(t), g'(t), h'(t) \rangle = f'(t) \mathbf{i} + g'(t) \mathbf{j} + h'(t) \mathbf{k}$$

V EXAMPLE 8


- (a) Find the derivative of $\mathbf{r}(t) = (1 + t^3)\mathbf{i} + te^{-t}\mathbf{j} + \sin 2t\mathbf{k}$.
- (b) Find the unit tangent vector at the point where $t = 0$.

EXAMPLE 9 For the curve $\mathbf{r}(t) = \sqrt{t} \mathbf{i} + (2 - t) \mathbf{j}$, find $\mathbf{r}'(t)$ and sketch the position vector $\mathbf{r}(1)$ and the tangent vector $\mathbf{r}'(1)$.

In fact, plot the curve on octave and all unit tangent vectors.

A curve given by a vector function $\mathbf{r}(t)$ on an interval I is called **smooth** if \mathbf{r}' is continuous and $\mathbf{r}'(t) \neq \mathbf{0}$ (except possibly at any endpoints of I). For instance, the helix in Example 10 is smooth because $\mathbf{r}'(t)$ is never $\mathbf{0}$.

EXAMPLE 11 Determine whether the semicubical parabola $\mathbf{r}(t) = \langle 1 + t^3, t^2 \rangle$ is smooth.



5 THEOREM Suppose \mathbf{u} and \mathbf{v} are differentiable vector functions, c is a scalar, and f is a real-valued function. Then

$$1. \frac{d}{dt} [\mathbf{u}(t) + \mathbf{v}(t)] = \mathbf{u}'(t) + \mathbf{v}'(t)$$

$$2. \frac{d}{dt} [c\mathbf{u}(t)] = c\mathbf{u}'(t)$$

$$3. \frac{d}{dt} [f(t)\mathbf{u}(t)] = f'(t)\mathbf{u}(t) + f(t)\mathbf{u}'(t)$$

$$4. \frac{d}{dt} [\mathbf{u}(t) \cdot \mathbf{v}(t)] = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$$

$$5. \frac{d}{dt} [\mathbf{u}(t) \times \mathbf{v}(t)] = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$$

$$6. \frac{d}{dt} [\mathbf{u}(f(t))] = f'(t)\mathbf{u}'(f(t)) \quad (\text{Chain Rule})$$

V EXAMPLE 12 Show that if $|\mathbf{r}(t)| = c$ (a constant), then $\mathbf{r}'(t)$ is orthogonal to $\mathbf{r}(t)$ for all t .

- 55.** The curves $\mathbf{r}_1(t) = \langle t, t^2, t^3 \rangle$ and $\mathbf{r}_2(t) = \langle \sin t, \sin 2t, t \rangle$ intersect at the origin. Find their angle of intersection correct to the nearest degree.

Find $\mathbf{r}(t)$ if $\mathbf{r}'(t) = 2t\mathbf{i} + 3t^2\mathbf{j} + \sqrt{t}\mathbf{k}$ and $\mathbf{r}(1) = \mathbf{i} + \mathbf{j}$.

If two objects travel through space along two different curves, it's often important to know whether they will collide. (Will a missile hit its moving target? Will two aircraft collide?) The curves might intersect, but we need to know whether the objects are in the same position *at the same time*. Suppose the trajectories of two particles are given by the vector functions

$$\mathbf{r}_1(t) = \langle t^2, 7t - 12, t^2 \rangle \quad \mathbf{r}_2(t) = \langle 4t - 3, t^2, 5t - 6 \rangle$$

for $t \geq 0$. Do the particles collide?

