Response: OpenAI Hilbert–Smith preprint

Sherlock reverse-verification notes

Alan J. Coppola (AI-assisted / AI-generated using Cursor, with Grok, Anthropic, and OpenAI models)

2026-10-07

Independent read of the OpenAI Math Release preprint ***The Hilbert--Smith conjecture in every finite dimension***

Upstream: `github.com/openai/math` SHA: `adc7f12…`

Repo: `github.com/ajjcoppola/hilberts-5th-oai`

This work is AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the direction of Alan J. Coppola.

Original questions

1. Overview: how faithful p-adic actions are excluded near the identity homeomorphism.
2. Geometric argument, or algebraic invariants in every Euclidean neighborhood?
3. Public / internet responses to this AI-generated proof?
4. Categorize the proof; what does it say about the structure of $\mathbb{R}^n$?
5. Hold/break under Menger-type sponges as $\mathbb{R}^n$ replacements (where the theorem is false)?
6. Is the contradiction that $\mathbb{R}^n$ cannot factor through an orbit space with "wonky" local sphere maps?

TL;DR

OpenAI's preprint claims a proof of the Hilbert–Smith conjecture in every finite dimension by ruling out faithful continuous actions of the p-adic integers on topological manifolds. The argument is not the classical Yang orbit-dimension obstruction. Instead it is an integrality-versus-divisibility contradiction for a sheaf-theoretic Witt/signature invariant: a small open subgroup $G \cong \mathbb{Z}_p$ acts on a Euclidean chart, the action is stabilized into an odd-dimensional sphere S^d, and averaging near the identity produces a degree-one test through the orbit space. Character symmetry then forces that orientation class to split into arbitrarily fine equal rational pieces, which cannot fit inside a fixed denominator lattice on $E_d(S^d)$. A targeted audit found no outright failure in the load-bearing lemmas, though several steps remain weak pending expert or Lean review; on Menger-type sponges, where faithful $\mathbb{Z}_p$ actions do exist, the proof correctly refuses to run because Newman and Euclidean-chart hypotheses fail.

Proof overview (Q1, Q2, Q4, Q6)

Proof overview — OpenAI Hilbert–Smith preprint (2026-09-23)

AI disclaimer: This audit is AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the direction of Alan J. Coppola.

Answers to reverse-verification questions Q1, Q2, Q4, Q6. Source: vendored LaTeX in `paper/`.


Statement

Theorem (Hilbert–Smith, all finite dimensions). A locally compact second-countable Hausdorff group acting faithfully and jointly continuously on a connected Hausdorff second-countable topological n-manifold without boundary is a Lie group.

Reduction. Classical structure theory $\Rightarrow$ it suffices to exclude faithful continuous actions of the additive p-adic integers $\mathbb{Z}_p$ on such manifolds (`thm:main`).

Mechanism in one paragraph

Assume a faithful $\mathbb{Z}_p$-action on M^n. Newman's theorem produces a small open subgroup $G \cong \mathbb{Z}_p$ acting faithfully and orientation-preservingly on a connected open chart set $O \subset \mathbb{R}^n$. Stabilize to odd dimension $d > n$ by $W = O \times \mathbb{R}^{d-n} \subset \mathbb{R}^d \subset S^d$. Pass to the one-point compactification W^+ and orbit space $Y = W^+/G$. Average a degree-one pinch over a sufficiently small open subgroup so a degree-one test $S^d \to Y \to S^d$ factors through Y. Build an integral Witt invariant $E_d(S^d)$ of sheaf forms (continuous polyhedral generators). Character projectors on finite partitions of p-power roots of unity produce p^k integral classes z_i that sum to $4u_d$. Finite quotient sheets plus an odd-sphere concentration homotopy force all rational images $\overline{z}_i$ equal. Hence each is $(4/p^k)\overline{u}_d$. But a prior comparison with the semialgebraic subcategory shows all integral classes lie in the fixed lattice $L_d^{-1}\mathbb{Z}\,\overline{u}_d$ with $L_d = 2^{e_d}$ independent of k. Choose $p^k > 4L_d$: contradiction.


Faithful Zp on M^n
Newman: faithful G on O in R^n
W = O x R^(d-n) in S^d
Average pinch: deg-1 test through Y
Odd-sphere: concentrate D_s o a on V
Region V with p^k sheets
beta permutes character classes
Lattice: denom bound L_d = 2^e
Sum = 4 u_d so each = 4/p^k u_d

Q1 — Why no p-adic homeomorphisms near id?

The point is not that $\mathbb{Z}_p$ is "too large," but that it has **arbitrarily small open subgroups** in the compact-open topology on $\mathrm{Homeo}(M)$.

1. **Chart reduction** (`prop:chart-reduction`, §5): if every open subgroup fixed a neighborhood of every point, Newman's theorem plus connectedness would force the action trivial, contradicting faithfulness. So some point has no neighborhood fixed by a whole open subgroup; a small open $G \cong \mathbb{Z}_p$ acts faithfully on an invariant chart open O.
2. **Averaging near id** (`lem:averaged-test`): take a degree-one pinch supported in a ball in W. Because G is profinite, a small open $H \subset G$ moves that pinch by Euclidean distance $< 1/2$ uniformly. Haar averaging produces an H-invariant map still of degree one, which descends through $Y = W^+/H$.
3. **Finite groups cannot do this:** for a free/near-free $\mathbb{Z}/p$ action, any invariant map through the orbit space has degree divisible by p (transfer). So there is no degree-one averaged test. The arithmetic contradiction needs $p^k > 4L_d$; for odd p even $k = 1$ can suffice, but only after a degree-one (or coprime-to-p) test exists — which requires a **small** subgroup.

So the disproof is: a faithful action near the identity would manufacture arbitrarily fine equal integral Witt pieces of a fixed orientation class on S^d, contradicting the denominator lattice.

Q2 — Geometric or algebraic?

Mostly algebraic-invariant, with a thin geometric shell.

Layer	Nature	Content
Chart / Newman / orientation	Geometric	Local Euclidean charts, orientation preservation
Averaging / sheets / odd-sphere	Topological	Haar average, Serre on $\pi_*(S^{\mathrm{odd}})$, Čech
Duality / Witt / lattice	Algebraic	Verdier duality on generated categories; Balmer triangular Witt; integral comparison $\mathcal{P} \to \mathcal{T}$
Contradiction	Arithmetic	Equal parts vs bounded denominators

The obstruction is **not** Yang/Bredon–Raymond–Williams cohomological-dimension raising of the orbit space. The paper explicitly avoids any dimension bound on Y. The invariant that must hold in every Euclidean neighborhood is the **integer signature** of a degree-d form on a small oriented ball (`prop:generic-signature`), transferred to a fixed lattice on $E_d(S^d)$.

Q4 — What the proof says about the structure of $\mathbb{R}^n$

Open sets in $\mathbb{R}^n$ (after odd stabilization into S^d) carry:

1. **Local Poincaré duality** with real coefficients, packaged as Verdier dualizing complexes and orientation forms.
2. **Integral signature** for continuous (not merely PL) proper maps from polyhedra — after comparison with a semialgebraic generating subcategory and a uniform 2-power denominator.

3. **No infinite p-divisible equal splitting** of that orientation class through a continuous $\mathbb{Z}_p$-orbit factorization.

In slogan form: $\mathbb{R}^n$ (and finite-dimensional topological manifolds) are **signature-rigid** under continuous p-adic orbit quotients. The rigidity is measured on the ambient sphere, not on the possibly wild orbit space.

Q6 — Is the contradiction "orbit space factoring + wonky local sphere maps"?

Partly yes, but the wonky maps are controlled.

- Yes: the degree-one test **always factors** $S^d \xrightarrow{c} W^+ \xrightarrow{q} Y \xrightarrow{a} S^d$ (`lem:averaged-test`). That is the orbit-space factorization.
- The "wonky" part is concentrating a positive-degree postcomposition into a region with p^k labeled sheets (`prop:move-test` + `lem:finite-sheets`), then using a character multiplier β to cyclically permute Witt character classes (`prop:equal-parts`).
- The **basic contradiction** is arithmetic on S^d:

$$\sum_{i=0}^{p^k-1} \overline{z}_i(a) = 4\overline{u}_d, \qquad \overline{z}_i(a) = \frac{4}{p^k}\overline{u}_d \notin L_d^{-1}\mathbb{Z}\,\overline{u}_d$$

  for $p^k > 4L_d$.

So: factoring through an orbit space is necessary but not sufficient for the paradox. The paradox is that such a factorization, combined with faithfulness (arbitrarily fine finite sheets) and odd-sphere homotopy, would force unbounded denominators in a group proven to have bounded denominators.

Section map

Section	Role
§1 Intro	Statement, predecessors, numerical sketch
§2 Categories	$\mathcal{T}(X)$, $\mathcal{P}(X)$, duality, homotopy invariance
§3 Localization	Germs, Witt localization, dense doubling
§4 Lattice	Bounded $\mathcal{P} \to \mathcal{T}$ comparison; L_d; degree; contractible quotient
§5 Local tests	Chart, average, sheets, move test
§6 Characters	Projectors, equal parts, contradiction, Lie conclusion
App.	Duality signs for cylinder

Verdict pointer

Targeted audit grades and sponge stress tests: `02_audit_ledger.md`, `03_sponge_stress_test.md`.

Audit ledger (Targets A–C)

Targeted audit ledger (Sherlock targets A–C)

AI disclaimer: This audit is AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the direction of Alan J. Coppola.

Scope: line-by-line on load-bearing lemmas only (plan: targeted audit).
Grades: `PASS` = argument appears complete and correctly cited; `WEAK` = plausible but citation gap, omitted routine detail, or hypothesis that needs expert check; `FAIL` = identified break.
Overall targeted verdict: No FAIL found. Several `WEAK` items remain the natural places for independent expert / Lean scrutiny. The proof does **not** "prove too much" on known sponge models (see `03_sponge_stress_test.md`).

Target A — Homotopy invariance + bounded lattice

`thm:homotopy` — Homotopy invariance

File: `paper/sections/02-categories.tex` ~701–739
Grade: PASS (with WEAK dependence on cylinder signs)

- Builds on `cat:endpoint-forms` (neutral cylinder form with opposite endpoint signs) and `prop:test-pushforward`.
- Correctly notes that the homotopy need not be proper: objects have compact support in $I \times K$.
- Implements Woolf's interval argument inside the *generated* continuous category $\mathcal{T}$, without requiring constructible images.
- **Residual risk:** absolute normalization of the cylinder scalar c is deferred to the appendix (`app:duality-signs`). Sign bookkeeping is historically delicate; appendix exists and is cited, so not FAIL.

`thm:bounded-comparison` — Bounded $\mathcal{P} \to \mathcal{T}$ comparison on spheres

File: `paper/sections/04-lattice.tex` ~125–185
Grade: PASS

- Inducts on sphere dimension via pole covers U, V, localization sequences, `prop:open-cover` density, and `thm:dense-doubling` (exponent 2).
- Finite diagram chases (`lat:finite-chases`) give a uniform 2-power annihilating ker/coker in every degree.
- Exponent bookkeeping (`lat:exponent-bookkeeping`) is explicit ($e_j = 3(9^j - 1)/4$); only existence of some fixed e_d is used later.
- **Why this is load-bearing:** without a *uniform* denominator independent of later k, the equal-parts arithmetic cannot contradict.

`prop:sphere-rank` — Rational sphere calculation

File: `04-lattice.tex` ~203–251
Grade: PASS

- Mayer–Vietoris / localization induction shows $G_r(S^j) \otimes \mathbb{Q} \simeq (G_r(\mathrm{pt}) \oplus G_{r-j}(\mathrm{pt})) \otimes \mathbb{Q}$ for both $F = W(\mathcal{P})$ and $E = W(\mathcal{T})$.
- For odd d, $E_d(S^d) \otimes \mathbb{Q}$ is 1-dimensional.
- Dense doubling supplies rational isomorphism of relative groups; no claim of integral isomorphism $\mathcal{P} \to \mathcal{T}$ is needed here.

`thm:sphere-lattice` — **Fixed sphere lattice**

File: `04-lattice.tex` ~370–398
Grade: PASS

- Combines rational 1-dimensionality with `prop:generic-signature` (semialgebraic local signature $\mathbb{Z}\overline{u}_{F,d}$) and the 2^{e_d} cokernel bound.
- Result: $\mathrm{im}(E_d(S^d) \to E_d(S^d) \otimes \mathbb{Q}) \subseteq L_d^{-1}\mathbb{Z}\overline{u}_d$ with $L_d = 2^{e_d}$ **independent of action, p, and k.**
- **Key meta-point:** rational equivalence alone would lose the denominator bound (remarked in §1); the paper's integral comparison is essential.

`prop:generic-signature` **(supporting)**

Grade: PASS

- Hardt triviality ⇒ dense open where cohomology is finite locally constant; local signature on small balls is integral and respects Witt relations.
- Standard semialgebraic topology; correctly limited to $\mathcal{P}$, then transferred.

Target B — Germs, localization, dense doubling

`thm:germs` — **Germ description of quotient Homs**

File: `paper/sections/03-localization.tex` ~165–219
Grade: PASS

- Cutoff (`lem:cutoff`) + compact-support characterization (`prop:compact-support`) produce roofs representing germs.
- Fullness/faithfulness for summand sources handled carefully.
- This is the device that lets character projectors live on $X\backslash\{b\}$ while classes live in $E_d(X)$.

`thm:witt-localization` — **Balmer–Walter LES**

File: `03-localization.tex` ~227–293
Grade: WEAK (citation-correct, hypothesis-heavy)

- Invokes Balmer–Walter 2002, Theorem 2.1.
- Checks: thick duality-stable kernel, essential smallness, 2 invertible in Hom groups (real vector spaces), TR4$^+$, duality preserves S.
- **Why WEAK not PASS:** correctness reduces to whether Balmer–Walter applies verbatim to these *generated* subcategories of $D^b(\mathrm{Sh}_{\mathbb{R}}(X))$. The checklist looks complete; an independent reader should still verify against the published hypotheses (the paper notes BW removes the earlier weak-cancellation assumption).

`prop:open-cover` — **Dense inclusion from open covers**

File: `03-localization.tex` ~301–334
Grade: PASS

- Germs identify the two quotient Hom groups; cutoff proves every object is a summand of an extended object.
- Feeds the sphere induction in Target A.

`thm:dense-doubling` — **Explicit** d **with** $\iota d = d\iota = 2$

File: `03-localization.tex` ~344–460
Grade: WEAK

- Constructs $D(q) = q \perp q \perp H(x[1])$ in the smaller category via $y \oplus y[1] \in \mathcal{A}$.
- Neutrality descent: builds a large lagrangian L and argues $L \in \mathcal{A}$ by successive cones/split triangles.
- Parallel to Hornbostel–Schlichting cofinality, but self-contained.
- **Why WEAK:** the lagrangian membership argument (`L` via cones of $a \oplus 0$, etc.) is the longest purely formal stretch in §3; easy to hide a category-membership slip. No concrete contradiction found on reading; recommend Lean or second-human check.

`prop:contractible-quotient` — **Integral iso after killing contractible open**

File: `04-lattice.tex` ~487–529
Grade: PASS

- For odd $d \geq 3$, $E_d(U) = 0$ on contractible U; $E_{d-1}(U) \to E_{d-1}(X)$ split-injective via point projection (handles both $d \equiv 1, 3 \pmod 4$).
- Localization LES $\Rightarrow$ integral isomorphism $E_d(X) \xrightarrow{\sim} W_d(\mathcal{T}(X)/\mathcal{T}(U))$.
- Used to lift character involution classes without splitting quotient idempotents.

Target C — Character classes and equal parts

`lem:character-coproduct`

File: `paper/sections/06-character-classes.tex` ~37–99
Grade: WEAK

- $Rq_*\underline{\mathbb{C}} \simeq \bigoplus_\lambda \mathcal{H}_\lambda$ via vanishing of higher cohomology on zero-dimensional orbits + eigenspace decomposition of finite cyclic representations.
- Derived pushforward of finite character subsum isomorphic to sum of pushforwards, using soft resolutions and compactness of fibers.
- **Why WEAK:** infinite character support / commuting of Rt_* with infinite sums is subtle; the paper addresses it via soft sheaves on compact Hausdorff spaces. Expert sheaf check recommended.

`lem:orthogonality`

File: `06-character-classes.tex` ~150–203
Grade: PASS

- G-action by orientation-preserving isometries; complex structure J with $J^\dagger = -J$; cross pairings between distinct character summands vanish.
- Implies self-adjointness of finite partition projectors on the punctured target.

`def:character-classes` / `prop:sum-classes`

File: `06-character-classes.tex` ~225–291
Grade: PASS

- Classes defined as $\ell_U(z_i) = [A, \alpha] + [A, \alpha \circ (2e_i - \mathrm{id})]$ in the quotient Witt group — **without** requiring images of idempotents in $\mathcal{T}(X)$.
- Matrix isometry S proves $\sum z_i = 2[A, \alpha]$ integrally.
- Elegant and appears correct; this is where the paper avoids a common false step (splitting continuous projectors over b).

`prop:equal-parts`

File: `06-character-classes.tex` ~411–443
Grade: WEAK

- On a concentrated test, locally constant β of modulus 1 multiplies coefficients and cyclically permutes character summands; induces quotient isometries conjugating $e_i \mapsto e_{i-1}$.
- **Why WEAK:** must check that M_β is an isometry of the *restricted manifold* pairing α_h, that it matches under $A_h|_{X^\circ} \simeq Rt_*\mathcal{H}|_{X^\circ}$, and that the germ theorem places the conjugation in $\mathcal{Q}_U(X)$. The sketch is coherent; this is the sharpest geometric–algebraic interface in §6.

`prop:move-test`

File: `paper/sections/05-local-tests.tex` ~478–517
Grade: PASS (topology); WEAK dependence on `lem:orbit-injectivity`

- Pullback of orientation class vanishes on $F = Y \backslash V$ by moving the pinch into $q^{-1}(V)$ and using orbit-injectivity of rational Čech.
- Odd-sphere lemma (`lem:odd-sphere-null`) + Tietze neighborhood extension concentrate $D_s a$.
- Odd-sphere telescope $\to K(\mathbb{Q}, d)$ is classical (Serre; also used by Dranishnikov–Ferry–Weinberger).
- **`lem:orbit-injectivity` WEAK residual:** claims injectivity of $r^*\colon \check{H}^*(E/G; \mathbb{Q}) \to \check{H}^*(E; \mathbb{Q})$ with **no covering-dimension bound**, via vanishing of higher cohomology on zero-dimensional orbits and Haar averaging of locally constant $\mathbb{Q}$-functions. Argument looks carefully written; still a place specialists may probe because Y can be infinite-dimensional.

Cross-cutting residual risks (not graded FAIL)

1. **Generated continuous category** $\mathcal{T}$ is larger than constructible/Woolf settings; duality restriction (`prop:test-duality`) is argued via simplex generators — appears sound, not fully re-audited line-by-line outside Targets A–C.
2. **No Lean formalization** of this preprint found in the Oct 2026 release materials (see `04_public_responses.md`).
3. **Homology-manifold edge case** is outside the theorem's hypotheses; see sponge doc.

Suggested expert / Lean priorities

Priority	Object	Why
1	`thm:dense-doubling` neutrality descent	Long formal Witt argument
2	`prop:equal-parts` + germ isometry of M_β	Action enters here
3	`thm:witt-localization` vs Balmer–Walter hypotheses	Citation bridge
4	`lem:orbit-injectivity` on wild compacta	No dim bound
5	Cylinder signs appendix	Historical foot-gun

Sanity check (finite $\mathbb{Z}/p$)

See `03_sponge_stress_test.md` §Sanity: failure is exactly at `lem:averaged-test` (no degree-one invariant pinch). The arithmetic steps are not reached. **Sanity PASS.**

Sponge stress test (Q5)

Sponge / non-manifold stress test

AI disclaimer: This audit is AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the direction of Alan J. Coppola.

Goal (plan): Can the argument be run on spaces where faithful continuous $\mathbb{Z}_p$-actions **exist**, thereby "proving too much"? If every lemma's hypotheses hold on such a space, that is a candidate gap. If a manifold-specific hypothesis fails early, the proof is correctly scoped.

Framing: A Menger sponge where Hilbert–Smith is false does **not** refute a theorem about manifolds. It is a probe for overclaiming.

Sanity: finite $\mathbb{Z}/p$ on $\mathbb{R}^2$

Step	Status
Chart / Newman	OK for finite cyclic rotation (or fails faithfulness differently)
`lem:averaged-test` degree-one through orbit space	**FAILS**
Lattice / equal parts	Not reached

Why averaging fails. For $G = \mathbb{Z}/p$ acting (nearly) freely, any G-invariant continuous map $W^+ \to S^d$ that descends through $Y = W^+/G$ has degree divisible by p (transfer / covering degree of the regular orbits). There is no open subgroup smaller than G to average over while preserving degree 1.
Conclusion: The proof does not incorrectly rule out finite cyclic actions. Break is exactly at `lem:averaged-test`. **Sanity PASS.**

Break-point matrix

For each model: first lemma/hypothesis that fails when one tries to copy the proof.
- means the step still makes sense; the chain stops at the first **BREAK**.

Model	Faithful $\mathbb{Z}_p$?	First BREAK	Later notes
Topological n-manifold (theorem setting)	Conjecturally no; paper claims no	*(none — target of proof)*	Full chain
Finite $\mathbb{Z}/p$ on $\mathbb{R}^2$	N/A (finite)	`lem:averaged-test`	Degree divisible by p

Model	Faithful $\mathbb{Z}_p$?	First BREAK	Later notes
$\mathbb{Z}_p \times \mathbb{R}^n$ (left translation on $\mathbb{Z}_p$)	Yes	**Not a manifold** / no Euclidean chart (`prop:chart-reduction`)	Locally $\mathbb{Z}_p \times \mathbb{R}^n \not\simeq \mathbb{R}^{n+k}$
p-adic solenoid Σ_p	Yes (standard free action; orbit S^1)	**Not a manifold**; no open $O \subset \mathbb{R}^n$; `lem:collapse` needs $W \subset \mathbb{R}^d$	Averaging on solenoid can produce maps to S^1; orientation form u_d of chart type absent
Menger compactum μ^n (Dranishnikov 1989)	Yes — free action of any zero-dimensional compact group	**Not a manifold**; Newman fails; no chart in $\mathbb{R}^n$; collapse into S^d fails	Orbit dim can equal n; cohomological dim may jump — irrelevant here because chart already fails
Menger sponge / curve μ^1 ("Munger sponge")	Yes (special case of Dranishnikov)	Same as μ^n	Classic Cantor-like 1-dimensional universal curve
Raymond–Williams compactum (Ann. of Math. 1963; arXiv:1909.12660)	Yes; $\dim X = n$, $\dim X/\mathbb{Z}_p = n+2$	**Not a manifold**; chart/collapse fail	Historically motivated by Yang's dim-raising obstruction for *manifolds*

Model	Faithful $\mathbb{Z}_p$?	First BREAK	Later notes
ENR homology n-manifold (e.g. Bryant–Ferry–Mio–Weinberger; Bing dogbone $\times\mathbb{R}$)	**Unknown** whether faithful $\mathbb{Z}_p$ can exist	**OPEN EDGE**	Local duality / orientation may partially survive; Newman-type results are subtler. Paper's hypotheses require topological manifolds (charts to $\mathbb{R}^n$). If a free $\mathbb{Z}_p$ action on an ENR homology manifold were found, one must check whether `prop:chart-reduction` + orientation pairing still apply — potential stress region, **not** a present counterexample to the written theorem

---

Detailed sponge notes

1. Dranishnikov Menger actions (including the sponge)

Fact: Every Menger manifold (in particular μ^n) admits a free continuous action of $\mathbb{Z}_p$ (Dranishnikov, *Math. USSR Izv.* 32 (1989)).

Why the OpenAI argument stops:

1. μ^n is not locally Euclidean → `prop:chart-reduction` hypotheses false.
2. There is no open embedding $W \hookrightarrow \mathbb{R}^d \to$ continuous collapse $c : S^d \to W^+$ of §5 is undefined in the paper's form.
3. Newman's theorem is a **manifold** theorem; fixed-point-open ⇒ identity fails for sponges.
4. No orientation class of an open set in $\mathbb{R}^d$ to push into $E_d(S^d)$ via *tqc*.

The lattice theory on S^d itself remains fine — it never sees the sponge. The action never produces the character classes on S^d through a degree-one chart test.

"Proves too much?" No. Manifold hypotheses fail at the first geometric step.

2. Solenoid

The mapping torus / inverse-limit circle $\Sigma_p = \varprojlim(S^1 \xrightarrow{\times p} S^1)$ carries a free $\mathbb{Z}_p$-action with orbit space S^1.

- Orbit space is a manifold, but the **total space** is not.
- One can average maps to spheres, but the paper's degree-one test is built from a pinch in an open Euclidean W, not from abstract averaging on a solenoid.
- Break: chart + collapse + orientation of $W \subset \mathbb{R}^d$.

3. Raymond–Williams

These examples show dim-raising for $\mathbb{Z}_p$ actions on **compacta**, historically tied to: *if* a manifold admitted a faithful $\mathbb{Z}_p$ action then Yang's cohomological dimension relation would force $\mathrm{cd}_{\mathbb{Z}}(M/\mathbb{Z}_p) = \dim M + 2$.

The OpenAI proof **does not use** that obstruction. It uses signature lattices on S^d. On Raymond–Williams spaces the signature-chart pipeline never starts. So RW examples neither confirm nor refute the algebraic steps; they only show that "wild orbit spaces exist" — which the paper already allows for manifolds' orbit spaces without bounding their dimension.

4. Homology manifolds (edge)

Topological manifolds ⇒ ENR homology manifolds, but not conversely in high dimensions (with exotic examples). Local Čech duality and orientation sheaves can exist without Euclidean charts.

Paper ingredient	Homology manifold?
Newman	Partial analogues exist in literature; not automatic
Open $O \subset \mathbb{R}^n$	**Fails** without resolution/chart
Orientation pairing on $W \subset \mathbb{R}^d$	Needs Euclidean (or at least locally dualizable oriented) open set inside S^d
Averaging / sheets / characters	Could make sense abstractly

Verdict: The written proof uses **Euclidean charts** essentially (`prop:chart-reduction`, $W \subset \mathbb{R}^d$, collapse from S^d). It does not claim Hilbert–Smith for homology manifolds. No known counterexample on that class was used to break the argument. Flag for future work, not as a current FAIL.

Q5 answer (sponges as $\mathbb{R}^n$ replacements)

Through the Menger/Cantor-sponge lens:

1. Hilbert–Smith is **false** if "manifold" is replaced by "Menger compactum / sponge."
2. The OpenAI proof **correctly refuses** to run there: it needs Newman + Euclidean chart + orientation collapse into S^d.
3. The algebraic lattice on $E_d(S^d)$ is a statement about spheres/polyhedra, not about $\mathbb{R}^n$ alone; the manifold enters by manufacturing a degree-one test through a $\mathbb{Z}_p$-orbit space from a chart action.
4. Therefore sponges are **consistency checks**, not disproofs. They show the geometric hypotheses are sharp.
5. The only alarming outcome would have been: a sponge satisfying every cited lemma while admitting free $\mathbb{Z}_p$. **That did not occur.**

"Proves too much" summary

Test	Result
Finite $\mathbb{Z}/p$	Breaks at averaging — good
Menger / sponge / solenoid / RW	Break at manifold/chart/collapse — good
Homology manifolds	Outside scope; open edge
Candidate gap from sponge pass-through	**None found**

Combined with the audit ledger: **no sponge-based disproof of the preprint; no FAIL in targeted algebraic steps; WEAK items remain for expert review.**

Public responses (Q3)

Public responses to the OpenAI Hilbert–Smith preprint (Q3)

AI disclaimer: These notes are AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the direction of Alan J. Coppola.

Search date: 2026-10-07
Paper: OpenAI, *The Hilbert–Smith conjecture in every finite dimension*, 2026-09-23, in openai/math.

Executive finding

As of 2026-10-07 there is **no published technical rebuttal or confirmation** of this specific proof's correctness. Coverage is almost entirely about the **release process** of OpenAI's large math dump (hundreds of manuscripts), verification norms (Lean / Leiden / Palomar / Hexagon), and community reaction — not a line-by-line reading of the Hilbert–Smith argument.

Lean status for this preprint: No Lean formalization of the full Hilbert–Smith proof was found in the OpenAI math release materials. Related Lean activity exists only for *stating* the conjecture or Pardon's 3D theorem (DeepMind `formal-conjectures`), and for other OpenAI results (`openai/ten-proofs`), not this manuscript.

News / commentary (process, not proof audit)

Source	Date (approx.)	Substance re: Hilbert–Smith
New Scientist	~2026-10	Lists Hilbert–Smith among claimed results; Buzzard caution on which number-theory items look impressive / Lean-verified; notes AGMAI guidelines largely not followed
Scientific American	~2026-10	Dump will take months to parse; many results Lean-checked but **not all**; company mathematicians may not yet understand all releases
It Does What Now?	2026-10-06	Explicitly names Hilbert–Smith among claimed settlements; early reaction focuses on release ethics (Kra, Guillen, Litt, Bubeck); "nobody had yet had time to judge" correctness
OfficeChai	~2026-10	Topology row lists Hilbert–Smith; stresses verification as the open question; Tao warning on superficially flawless AI proofs
AGI Hunt / aggregator posts	2026-10-07	Mentions Mahler, Unique Games, Hilbert–Smith, etc., as claimed in the batch

Expert / community technical venues

Venue	Finding
MathOverflow	No hit on this OpenAI Hilbert–Smith preprint / Witt-sheaf argument (searches returned unrelated Witt-vector / prismatic posts)
arXiv	Paper appears to live on GitHub `openai/math`, not as a standard arXiv submission under that title (as of search)
Tao / Buzzard blogs	Commentary on AI math release practices in general; no dedicated Hilbert–Smith erratum/endorsement found in search
Palomar / Hexagon / Leiden	News notes OpenAI was encouraged to use these and largely did not for this dump

Lean / formalization adjacency

Item	Relation
google-deepmind/formal-conjectures #2188	Issue to **formalize the statement** of Hilbert–Smith (pre-dates / independent of OpenAI proof)
formal-conjectures PR #4599	Draft formalization of **Pardon's 3D** p-adic exclusion statement
openai/ten-proofs	Lean for a *different* OpenAI package (sphere packing, codes, etc.) — not Hilbert–Smith

What to watch next

1. Expert notes from geometric topology / transformation groups (Pardon, Dranishnikov school, Raymond–Williams lineage).
2. Sheaf / Witt specialists reading §§2–4 (Woolf, Balmer–Walter users).
3. Any Lean port of `thm:dense-doubling`, `thm:sphere-lattice`, or `thm:contradiction`.
4. Journal submission / referee reports if the manuscript is submitted outside the GitHub dump.

Implication for this Sherlock session

Public silence on *technical* content means this repo's targeted audit and sponge stress test are **independent first-pass verification**, not a digest of an existing consensus. Grades in `02_audit_ledger.md` should be read as provisional expert-prep, not as community acceptance.


Lemma dependency ledger

Lemma dependency ledger

AI disclaimer: This ledger is AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the direction of Alan J. Coppola.

Extracted from vendored LaTeX labels and the proof's load-bearing citations. File paths are relative to `paper/`.

Core dependency chain (contradiction)



Targeted audit nodes (by Sherlock target)

Target	Labels	Location
A	`thm:homotopy`, `thm:bounded-comparison`, `prop:sphere-rank`, `thm:sphere-lattice`	`sections/02-categories.tex:701`, `04-lattice.tex:125,203,370`
B	`thm:germs`, `thm:witt-localization`, `prop:open-cover`, `thm:dense-doubling`, `prop:contractible-quotient`	`03-localization.tex:165,227,301,344`; `04-lattice.tex:487`
C	`lem:character-coproduct`, `lem:orthogonality`, `def:character-classes`, `prop:equal-parts`, `prop:move-test`	`06-character-classes.tex`; `05-local-tests.tex:478`

Target	Labels	Location
Sanity	`lem:averaged-test`	`05-local-tests.tex:135`

External citations used as black boxes

Citation	Role in proof
Newman 1931 / Pardon 2019	Periodic homeomorphism fixing open set is identity
Balmer–Walter 2002, Thm 2.1	Integral Witt localization sequence
Hornbostel–Schlichting 2004, App. A	Cofinality / dense doubling precedent
Woolf 2008	Cylinder homotopy for Witt of sheaves (antecedent)
Serre 1951, 1953	Finiteness of $\pi_j(S\hat{}d)$ (j>d); degree maps kill torsion on odd spheres
Hardt 1980	Semialgebraic triviality for generic signature
Stacks Project	Proper base change, soft sheaves, derived sums
Pardon 2013	Classical reduction: non-Lie ⇒ embeds Z_p

Full labeled environment inventory

See automated extract in session notes; 45 labeled theorem/lemma/proposition/definition/remark environments across sections 1–6 and the duality-signs appendix.

Sherlock session log

Sherlock session — Hilbert–Smith OpenAI preprint

AI disclaimer: This session log is AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the direction of Alan J. Coppola.

Date: 2026-10-07
Workspace: `/Users/ajjc/proj/Hilberts-5th-OAI`
Protocol: Sherlock Mode (6 steps) + plan `hilbert-smith_reverse_audit`

Outcome

Targeted reverse-verification of OpenAI's Sept 23, 2026 Hilbert–Smith preprint.

- **Mechanism:** integrality vs p^k-divisibility of sheaf Witt / signature classes on S^d, not Yang dim-raising.
- **Audit (Targets A–C):** 0 FAIL, ~4 WEAK (dense-doubling, witt-localization/BW, character-coproduct, equal-parts).
- **Sanity:** finite $\mathbb{Z}/p$ breaks at `lem:averaged-test` (good).
- **Sponges:** Menger/solenoid/RW break at chart/Newman; no "proves too much" gap. Homology manifolds flagged OPEN EDGE.
- **Public:** no technical rebuttal as of 2026-10-07; no Lean for this proof.

Deliverables

- `paper/` vendored (SHA `adc7f12…`)
- `docs/01–04`, `lemma_dependency_ledger.md`, `README.md`
- Canvas: `canvases/hilbert-smith-audit.canvas.tsx`

Follow-ups (Step 5)

1. Expert check of WEAK lemmas (Pardon / Woolf / Balmer–Walter users).
2. Lean targets: `thm:dense-doubling`, `prop:equal-parts`, `thm:sphere-lattice`.
3. Homology-manifold scope note if literature finds $\mathbb{Z}_p$ actions there.

Cleanup (Step 6)

`/tmp/hs_paper` scratch copy may be deleted after confirming `paper/` is complete. No temporary diagnostic code was added to the proof.

Appendix: contents

Contents

Proof overview (Q1, Q2, Q4, Q6) **2**

Proof overview — OpenAI Hilbert–Smith preprint (2026-09-23) . . . 2
Statement . . . 2
Mechanism in one paragraph . . . 2
Q1 — Why no p-adic homeomorphisms near id? . . . 4
Q2 — Geometric or algebraic? . . . 4
Q4 — What the proof says about the structure of $\mathbb{R}^n$ . . . 4
Q6 — Is the contradiction "orbit space factoring + wonky local sphere maps"? . . . 5
Section map . . . 5
Verdict pointer . . . 5

Audit ledger (Targets A–C) **6**

Targeted audit ledger (Sherlock targets A–C) . . . 6
Target A — Homotopy invariance + bounded lattice . . . 6
`thm:homotopy` — Homotopy invariance . . . 6
`thm:bounded-comparison` — Bounded $\mathcal{P} \to \mathcal{T}$ comparison on spheres . . . 6
`prop:sphere-rank` — Rational sphere calculation . . . 6
`thm:sphere-lattice` — Fixed sphere lattice . . . 7
`prop:generic-signature` (supporting) . . . 7
Target B — Germs, localization, dense doubling . . . 7
`thm:germs` — Germ description of quotient Homs . . . 7
`thm:witt-localization` — Balmer–Walter LES . . . 7
`prop:open-cover` — Dense inclusion from open covers . . . 8
`thm:dense-doubling` — Explicit d with $\iota d = d\iota = 2$ . . . 8
`prop:contractible-quotient` — Integral iso after killing contractible open . . . 8
Target C — Character classes and equal parts . . . 8
`lem:character-coproduct` . . . 8
`lem:orthogonality` . . . 9
`def:character-classes` / `prop:sum-classes` . . . 9
`prop:equal-parts` . . . 9
`prop:move-test` . . . 9
Cross-cutting residual risks (not graded FAIL) . . . 10
Suggested expert / Lean priorities . . . 10
Sanity check (finite $\mathbb{Z}/p$) . . . 10

Sponge stress test (Q5) **11**

Sponge / non-manifold stress test . . . 11
Sanity: finite $\mathbb{Z}/p$ on $\mathbb{R}^2$ . . . 11
Break-point matrix . . . 11
Detailed sponge notes . . . 13
1. Dranishnikov Menger actions (including the sponge) . . . 13
2. Solenoid . . . 13
3. Raymond–Williams . . . 14
4. Homology manifolds (edge) . . . 14

Q5 answer (sponges as $\mathbb{R}^n$ replacements) . . . 14
"Proves too much" summary . . . 15

Public responses (Q3) **16**
Public responses to the OpenAI Hilbert–Smith preprint (Q3) . . . 16
Executive finding . . . 16
News / commentary (process, not proof audit) . . . 16
Expert / community technical venues . . . 16
Lean / formalization adjacency . . . 17
What to watch next . . . 17
Implication for this Sherlock session . . . 17

Lemma dependency ledger **18**
Lemma dependency ledger . . . 18
Core dependency chain (contradiction) . . . 18
Targeted audit nodes (by Sherlock target) . . . 18
External citations used as black boxes . . . 19
Full labeled environment inventory . . . 19

Sherlock session log **20**
Sherlock session — Hilbert–Smith OpenAI preprint . . . 20
Outcome . . . 20
Deliverables . . . 20
Follow-ups (Step 5) . . . 20
Cleanup (Step 6) . . . 20

Repository README **23**
Hilbert–Smith reverse verification + homology-manifold boundary . . . 23
Homology manifolds (HM-1..HM-6) . . . 23
Homology-manifold deliverables . . . 24
Checkpoints (git tags) . . . 24
HM-6 construction sketch . . . 24
2026 preprint audits . . . 25
OpenAI (2026-09-23) . . . 25
Dabrowski (2026-10-08) . . . 26
MathOverflow / posts . . . 26

Repository README

Hilbert–Smith reverse verification + homology-manifold boundary

Sherlock-mode audits of two 2026 preprints claiming the Hilbert–Smith conjecture in every finite dimension, plus a **relative** geometric-topology program that maps those arguments across homology / cohomology manifolds (HM-1..HM-6) and isolates an open construction target when $\dim_{\mathbb{Z}} X = n$ but $\dim X = \infty$.

Both 2026 arguments sit in the same family — **integrality versus p-divisibility of a signature / Witt invariant** after a Euclidean chart and Haar averaging — not Yang dim-raising and not Pardon's 3-manifold mapping-class theorem.

Author: Alan J. Coppola
AI disclaimer: This work is AI-assisted and AI-generated using Cursor, with Grok, Anthropic, and OpenAI models, under the author's direction. See `docs/AI_DISCLAIMER.md`.
Repo: https://github.com/ajjcoppola/hilberts-5th-oai

Homology manifolds (HM-1..HM-6)

Paper: Alan J. Coppola, *p-Adic Transfer, Signature Tails, and Homology Manifolds* (18 Oct 2026 draft) —
`papers/padic-homology-manifolds/main.pdf`

Class	What it is	Faithful continuous $\mathbb{Z}_p$?
HM-1	Non-*clc* / coefficient-mismatch Čech cohomology manifolds	Counterexample *mechanism* (outside 2026 scopes)
HM-2	Shrinking necklace / junction $\mathbb{Z}$-homology manifolds	**Explicit** counterexample (local homology fails)
HM-3	Connected polyhedral L-homology manifolds	**No** — reduces to manifold case (Thm C)
HM-4	Resolvable ENR / nonempty manifold stratum	**No** — chart or $X \times \mathbb{R}^2$ (Thm D)
HM-5	BFMW ENR, $X_{\text{man}} = \emptyset$, Quinn $I(X) \neq 1$	**No** (relative to OpenAI ENR rephrase or Dabrowski k-tower); bounded-dim inverse limits blocked
HM-6	Non-ENR: $\dim_{\mathbb{Z}} X = n < \infty$, $\dim X = \infty$	**Open candidate** — escapes E/F/G(2); Borel-join EW+Williams scaffolding; $clc_{\mathbb{Z}}$ glue still open

Honesty: Exclusion on manifolds/ENRs is **relative** to OpenAI/Dabrowski geometric engines. This repo does **not** claim a standalone classical Hilbert–Smith proof. HM-6 is a construction program, not a claimed existence theorem.

Homology-manifold deliverables

Deliverable	Path
Paper PDF / TeX	`papers/padic-homology-manifolds/`
Main slides	`presentations/padic-homology-2026/inde`
Invariants deck (Quinn / Witt / Dabrowski + geometric germ)	`presentations/invariants-quinn-witt-da`
Reviewer checklist / pack	`presentations/padic-homology-2026/`
Campaign + ledger + HM notes	`counterexamples/hilbert-smith/` (`HOMOLOGY_MANIFOLDS.md`, `CAMPAIGN.md`)
Model catalog (incl. E4 = HM-6)	`counterexamples/models/CATALOG.md`
CoppolaHS Lean	`lean/CoppolaHS/` (`CyclotomicTower`, `QuinnIndex`, `DranishnikovHM6`, …)
HM-6 harness / loop	`scripts/hm6_equivariant_ew_harness.py`, `HM6_LOOP_LOG.md`
Finish-line plan (Trunks A/B + glue P3)	`plans/HM6_FINISH_LINE.md`
Literature library (arXiv PDFs + construction notes)	`library/README.md`
Outreach / collaborators	`docs/OUTREACH_COLLABORATORS.md`
Post drafts (MO / X / LinkedIn)	`docs/POST_DRAFTS.md`

Checkpoints (git tags)

Tag	Contents
`checkpoint/hm6-2026-10-18`	Archive paper, CoppolaHS Lean, HM-6 harness/loop, counterexample campaign
`checkpoint/invariants-deck-2026-10-19`	Quinn / Witt / Dabrowski invariants slides (KaTeX + geometric germ narrative)

Notes: `docs/CHECKPOINT_HM6_2026-10-18.md`.

HM-6 construction sketch

Open frontier where OpenAI ENR rephrase, Dabrowski k-tower, and bounded-dimension Euler obstruction all fail:

- **Trunk A:** growing lens / solenoid / circle-bundle stages (`library/constructions/trunk_A_lens_solenoid`
- **Trunk B:** finite recursive C_{p^i} + Edwards–Walsh Borel-join (`library/constructions/trunk_B_cyclic_EW.`
- **Shared finish P3:** $clc_{\mathbb{Z}}$ Poincaré glue (`library/constructions/glue_P3_clc.md`)

Finite-stage algebraic gates G1–G4 **PASS** symbolically; continuous glue remains **EDGE / open**.

2026 preprint audits

Proof	Date	Packaging	Audit PDF
OpenAI	2026-09-23	Witt/signature lattice on S^d; equal parts $4/p^k$	`docs/Hilbert_Smith_Audit.pdf`
Dabrowski	2026-10-08	Integral tail in $L_4(A_G(T))$; $\mathbb{CP}^2$ realization vs $p\mathbb{Z}$	`dabrowski/docs/Dabrowski_HS_Audit.pdf`

Side-by-side: `dabrowski/docs/05_vs_openai.md`.
Geometric packaging of the invariants: `presentations/invariants-quinn-witt-dabrowski/`.

Counterexample protocol: `counterexamples/README.md`.

OpenAI (2026-09-23)

Upstream: https://github.com/openai/math (vendored at `adc7f1241b42e322a6451854ab7e4b4c146bf78a`)

Quick answers

Q	Answer
1. Near-id exclusion	Small open $\mathbb{Z}_p$ subgroups + Haar averaging give a degree-1 test through the orbit space; finite groups cannot.
2. Geometric vs algebraic	Mostly Witt/signature lattice arithmetic; thin geometric shell (Newman, chart, Serre).
3. Public responses	Process news only (2026-10-07); no technical rebuttal; no Lean for this proof.
4. Structure of $\mathbb{R}^n$	Signature-rigid under continuous $\mathbb{Z}_p$ orbit factorization after odd stabilization into S^d.
5. Menger sponges	Theorem false for sponges; proof breaks at chart/Newman — correctly scoped, does not overclaim.
6. Orbit factoring?	Necessary but not the whole story; contradiction is equal parts vs fixed denominator lattice.

Targeted audit: no `FAIL`; several `WEAK` (dense doubling, Balmer–Walter bridge, character coproducts, equal-parts isometry).

- `paper/` — vendored LaTeX + PDF + provenance
- `docs/01_proof_overview.md` — Q1, Q2, Q4, Q6
- `docs/02_audit_ledger.md` — lemma grades
- `docs/03_sponge_stress_test.md` — Q5 + sanity
- `docs/04_public_responses.md` — Q3
- `docs/lemma_dependency_ledger.md`
- Rebuild: `python3 scripts/build_audit_pdf.py`

Dabrowski (2026-10-08)

Upstream: https://github.com/adbrw/HS_proof (vendored at `abb0c44d58d711235f20b666ef422a4ea87b9add`)

Quick answers

Q	Answer
1. Near-id exclusion	Small open tails of $\mathbb{Z}_p$ Haar-average a degree-1 pinch in a chart; unbounded finite quotients feed an L_4 tail of signature 1, which divisibility forces into $p\mathbb{Z}$.
2. Geometric vs algebraic	Mostly ultimate quadratic L-theory (Karoubi / Ranicki); thin Euclidean shell (chart, PL dual cells, $\mathbb{CP}^2$).
3. Public responses	None technical as of 2026-10-08 morning; claimed Lean 4 certificate not rerun here.
4. Structure of $\mathbb{R}^n$	Signature-tail rigid: a Euclidean germ realizes $\mathbb{CP}^2$ with $\sigma = 1$, which cannot be eventually p-divisible.
5. Menger sponges	Breaks at chart / combinatorial dual cells — correctly scoped.
6. Orbit factoring?	Necessary but not sufficient; contradiction is $\sigma_{\text{tail}} = 1$ vs eventual p-divisibility.

Targeted audit: no `FAIL` on load-bearing PDF lemmas; `WEAK` on NegK/tail extraction, Mayer–Vietoris nilpotence, the graph corner, $\mathbb{CP}^2$ evaluation, and Lean `#print axioms`.

Local `#print axioms HSFormal.hilbertSmith` (optional palomar clone): `propext`, `Classical.choice`, `Quot.sound` — see `dabrowski/PROVENANCE.md`, `dabrowski/lean-extension/BUILD_STATUS.md`.

- `dabrowski/paper/` — vendored `HSproof.pdf` + Challenge/Solution + lake files
- `dabrowski/lean/` — sparse load-bearing modules (not the full 238)
- `dabrowski/docs/01_proof_overview.md`
- `dabrowski/docs/02_audit_ledger.md`
- `dabrowski/docs/03_sponge_stress_test.md`
- `dabrowski/docs/04_public_responses.md`
- `dabrowski/docs/05_vs_openai.md`
- Rebuild: `python3 scripts/build_dabrowski_audit_pdf.py`

MathOverflow / posts

Paste-ready Euclidean-structure rewrite and HM-6 research question (not work-checking): `docs/POST_DRAFTS.md`.