

**EMERGENCY
2013-2018**Now
Published
as**Archives of Academic
Emergency Medicine**[Arch Acad Emerg Med](#). 2020; 8(1): e29.

Published online 2020 Mar 21.

PMCID: PMC7085862

PMID: [32232214](#)

Potential Treatments for COVID-19; a Narrative Literature Review

[Ali Rismanbaf](#)^{1,*}

¹Department of Clinical Pharmacy and Pharmacy Practice, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran.

*Corresponding author: Ali Rismanbaf; Isfahan University of Medical Sciences, Hezar Jerib Street, Isfahan, Iran. Email: alirismanbaf74@gmail.com

Received 2020 Jan; Accepted 2020 Feb.

[Copyright notice](#)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/3.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

SARS-CoV-2 is a newly emerging human infectious coronavirus that causes COVID-19, which has been recognized as a pandemic by the World Health Organization (WHO) on March 11th. There is still no vaccine or definitive treatment for this virus because its pathogenesis and proliferation pathways are still unknown. Therefore, in this article, new potential COVID-19 therapies are briefly reviewed.

Key Words: Coronavirus, Drug therapy, Clinical trial, Case reports, Review, COVID-19

Introduction

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) is a newly emerging human infectious coronavirus, originated in Wuhan, China, and has been spreading rapidly in China and other countries since December 2019 ([1](#)). The World Health Organization (WHO) also declared a global emergency on January 31st due to increasing concerns over its fast spread, and on March 11th the disease was recognized as a pandemic. Since the bases for pathogenesis of this virus and its proliferation is unclear, there is still no vaccine or definitive treatment against it. Thus, medications used against SARS-CoV-2 are mainly based on their effectiveness on earlier strains of coronavirus, SARS-CoV and MERS-CoV. Therefore, the immediate introduction of potential COVID-19 treatments can be essential and salvaging. In this article, new potential COVID-19 therapies are briefly reviewed.

Methodology

Articles were extracted, irrespective of time, using PubMed, Embase, and Google Scholar search engines, searching terms "COVID-19", "SARS-CoV-2", and "2019-nCoV" in titles, abstracts and keywords. Afterwards, clinical trials, clinical reports, case reports, and suggestions for potential medications against COVID-19 were briefly reviewed.

Results

Clinical reports

Clinical reports on COVID-19 treatment mainly described empirical treatments and clinical experiences during its treatment. In 2020, Gao et al. studied the effect of chloroquine and hydroxychloroquine in treatment of COVID-19 in over 100 patients and 10 hospitals in Wuhan, Jingzhou, Guangzhou, Beijing, Shanghai, Chongqing, and Ningbo. The results of this study showed that chloroquine phosphate is effective in preventing the exacerbations of pneumonia, decreasing lung involvements in imaging findings, promoting a virus-negative conversion and shortening the disease course. In addition, there were no serious adverse effects observed at therapeutic doses (2).

Also, according to Jian-ya et al., treatment of 51 COVID-19 patients with traditional Chinese medicine, interferon, Lopinavir, Ritonavir and short-term (3 to 5 days) corticosteroids was successful and resulted in recovery and discharge of 50 patients (3). Qin et al. also reported that administration of moxifloxacin, lopinavir, and interferon to non-ICU patients and the addition of methylprednisolone to the above treatment for ICU patients resulted in 26 patients being discharged from intensive care unit (ICU) and 16 patients being discharged from hospital (4). Also, Zhou et al. reported that short-term moderate-dose corticosteroid (160 mg/day) plus immunoglobulin (20 g/day) significantly reduced lung injury, normalized lymphocyte counts, body temperature, C-reactive protein levels, and oxygenation index in 10 COVID-19 patients (5). On the other hand, while studying 416 COVID-19 patients, Shang et al. reported that corticosteroid therapy and gamma globulin administration increased mortality and appeared to be useful only in patients with lower lymphocyte counts (6). According to the mentioned clinical reports, the administration of corticosteroids for COVID-19 patients is still questionable.

Case reports

So far, there are three published case reports on the successful treatment of patients with COVID-19. In the first report Lim et al. described a 54-year-old man with COVID-19 who was treated with Lopinavir/Ritonavir from day 10 of illness, 2 tablets (Lopinavir 200mg / Ritonavir 50mg) every 12 hours. Since first day of administration, β -coronavirus viral load started to decrease, and little or no detectable coronavirus titers have been observed since then (7).

In another case report, Zhang et al. described a couple who were both 38 years old and were suffering from COVID-19. Their treatment included Methylprednisolone 40 mg daily intravenous (IV) injections for one and five days for the male and the female patient respectively, human gamma globulin 10g IV qd for five and seven successive days for the male and the female patient, respectively, and then the dose was changed to 5g for both of them, in addition to Moxifloxacin, Oseltamivir, Arbidol hydrochloride, and Tanreqing (Chinese patent medicine). After 11 days, the female patient and after 14 days the male patient recovered with regards to inflammatory factors and were discharged from the hospital (8).

In the third case report, Chen et al. reported a 45-year-old woman with COVID-19 and stated that after treatment with Thalidomide (100 mg orally once a day) and Methylprednisolone (40 mg intravenously bid for 3 days then reduced to once a day for 5 days) the overall patient status was improved, oxygen index was increased, symptoms of nausea and vomiting were alleviated, and cytokine levels were decreased (9).

Potential drugs

Several articles have suggested medicines, potentially effective for the treatment of COVID-19 (Table 1). Most of these suggestions are based on in vitro studies, virtual screenings and records of their effects on SARS and MERS.

Table 1

Potential drugs for COVID-19

Study	Method	Medicine	Mechanism of Action
Wang et al. (2020) (12)	In vitro study	Chloroquine Remdesivir	Reducing viral copy numbers in the cell supernatant and viral infection
Zhang et al. (2020) (13)	In vitro study	Teicoplanin	Preventing the entrance of SARS-CoV-2-Spike-pseudoviruses into the cytoplasm
Xu et al. (2020) (14)	Virtual screening	Nelfinavir	Binding to SARS-CoV-2 M ^{pro}
		Colistin Valrubicin Icatibant Bepotastine	
Liu et al. (2020) (15)	Virtual screening	Epirubicin Epoprostenol Vapreotide Aprepitant Caspofungin perphenazine	Binding to SARS-CoV-2 M ^{pro}
Shang et al. (2020) (16)	Virtual screening	Rupintrivir Lopinavir Remdesivir	Binding to SARS-CoV-2 M ^{pro}
Jin et al. (2020) (17)	Virtual screening	Ebselen	Binding to SARS-CoV-2 M ^{pro}
Sekhar et al. (2020) (18)	Virtual screening	Beclabuvir Saquinavir	Binding to SARS-CoV-2 M ^{pro}
		(Angiotensin II human acetate)	
Contini et al. (2020) (19)	Virtual screening	GHRP-2 Indinavir Cobicistat Caspofungin acetate Lopinavir Atazanavir	Binding to SARS-CoV-2 M ^{pro} : Angiotensin II human acetate, GHRP-2, Indinavir, and Cobicistat Binding to SARS-CoV-2 3C-like proteinase (3CL ^{pro}): Angiotensin II human acetate, GHRP-2, Indinavir, Caspofungin acetate, Lopinavir, and Atazanavir

In addition to these medications, Tocilizumab has recently been suggested as a COVID-19 treatment. Studies have shown that IL-6 levels significantly correlated with the severity of COVID-19, C-reactive protein (CRP), lactate dehydrogenase (LDH), and D-dimer levels and T cell counts, and it has been

suggested that Tocilizumab, with its inhibitory effect on IL-6, may be effective in treatment of COVID-19 (10, 11). However, no clinical study has demonstrated the effects of Tocilizumab on COVID-19 and further studies are indeed required.

Conclusion

Apparently, in addition to the drugs currently prescribed to treat COVID-19, Arbidol hydrochloride, interferon, and Thalidomide plus Methylprednisolone can also be used due to their effects reported in clinical studies. However, more studies are needed to confirm the use of corticosteroids, as there are conflicting reports regarding their efficacy. Also, potential drugs listed in Table 1, such as Remdesivir, Atazanavir, Saquinavir, and Formoterol, and Tocilizumab can be introduced as treatments for COVID-19 if they prove to be effective in animal and clinical studies.

Conflict of interest

None

Funding and support

None

References

1. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *BioRxiv*. 2020 [[Google Scholar](#)]
2. Gao J, Tian Z, Yang X. Breakthrough: Chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *BioScience Trends*. 2020 [[PubMed](#)] [[Google Scholar](#)]
3. Jian-ya G. Clinical characteristics of 51 patients discharged from hospital with COVID-19 in Chongqing, China. *medRxiv*. 2020 [[Google Scholar](#)]
4. Qin X, Qiu S, Yuan Y, Zong Y, Tuo Z, Li J, et al. Clinical Characteristics and Treatment of Patients Infected with COVID-19 in Shishou, China. *China*. 2020;18 [[Google Scholar](#)]
5. Zhou Z-G, Xie S-M, Zhang J, Zheng F, Jiang D-X, Li K-Y, et al. Short-Term Moderate-Dose Corticosteroid Plus Immunoglobulin Effectively Reverses COVID-19 Patients Who Have Failed Low-Dose Therapy. 2020 [[Google Scholar](#)]
6. Shang J, Du R, Lu Q, Wu J, Xu S, Ke Z, et al. The Treatment and Outcomes of Patients with COVID-19 in Hubei, China: A Multi-Centered, Retrospective, Observational Study. 2020 [[Google Scholar](#)]
7. Lim J, Jeon S, Shin H-Y, Kim MJ, Seong YM, Lee WJ, et al. Case of the Index Patient Who Caused Tertiary Transmission of COVID-19 Infection in Korea: the Application of Lopinavir/Ritonavir for the Treatment of COVID-19 Infected Pneumonia Monitored by Quantitative RT-PCR. *Journal of Korean medical science*. 2020;35:6. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
8. Zhang Z, Li X, Zhang W, Shi Z-L, Zheng Z, Wang T. Clinical Features and Treatment of 2019-nCov Pneumonia Patients in Wuhan: Report of A Couple Cases. *Virologica Sinica*. 2020:1–7. [[PubMed](#)] [[Google Scholar](#)]
9. Chen C, Qi F, Shi K, Li Y, Li J, Chen Y, et al. Thalidomide combined with low-dose glucocorticoid in the treatment of COVID-19 Pneumonia. 2020 [[Google Scholar](#)]

10. Liu T, Zhang J, Yang Y, Zhang L, Ma H, Li Z, et al. The potential role of IL-6 in monitoring coronavirus disease 2019. medRxiv. 2020 [[Google Scholar](#)]
11. Diao B, Wang C, Tan Y, Chen X, Liu Y, Ning L, et al. Reduction and Functional Exhaustion of T Cells in Patients with Coronavirus Disease 2019 (COVID-19) medRxiv. 2020 [[Google Scholar](#)]
12. Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. Cell research. 2020;30(3):269–71. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
13. Zhang J, Ma X, Yu F, Liu J, Zou F, Pan T, et al. Teicoplanin potently blocks the cell entry of 2019-nCoV. bioRxiv. 2020 [[Google Scholar](#)]
14. Xu Z, Peng C, Shi Y, Zhu Z, Mu K, Wang X, et al. Nelfinavir was predicted to be a potential inhibitor of 2019-nCoV main protease by an integrative approach combining homology modelling, molecular docking and binding free energy calculation. bioRxiv. 2020 [[Google Scholar](#)]
15. Liu X, Wang X-J. Potential inhibitors for 2019-nCoV coronavirus M protease from clinically approved medicines. bioRxiv. 2020 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
16. Ton A-T, Gentile F, Hsing M, Ban F, Cherkasov A. Rapid Identification of Potential Inhibitors of SARS-CoV-2 Main Protease by Deep Docking of 13 Billion Compounds. ChemRxiv. 2020 [[PubMed](#)] [[Google Scholar](#)]
17. Jin Z, Du X, Xu Y, Deng Y, Liu M, Zhao Y, et al. Structure-based drug design, virtual screening and high-throughput screening rapidly identify antiviral leads targeting COVID-19. bioRxiv. 2020 [[Google Scholar](#)]
18. Sekhar T. Virtual Screening based prediction of potential drugs for COVID-19 [[Google Scholar](#)]
19. Contini A. Virtual screening of an FDA approved drugs database on two COVID-19 coronavirus proteins. 2020 [[Google Scholar](#)]
20. Wang J. Fast Identification of Possible Drug Treatment of Coronavirus Disease-19 (COVID-19) Through Computational Drug Repurposing Study. 2020 [[Google Scholar](#)]
21. Wang Q, Zhao Y, Chen X, Hong A. Virtual Screening of Approved Clinic Drugs with Main Protease (3CLpro) Reveals Potential Inhibitory Effects on SARS-CoV-2. 2020 [[Google Scholar](#)]
22. Chen YW, Yiu C-P, Wong K-Y. Prediction of the 2019-nCoV 3C-like protease (3CLpro) structure: virtual screening reveals velpatasvir, ledipasvir, and other drug repurposing candidates. ChemRxiv. 2020 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
23. Beck BR, Shin B, Choi Y, Park S, Kang K. Predicting commercially available antiviral drugs that may act on the novel coronavirus (2019-nCoV), Wuhan, China through a drug-target interaction deep learning model. bioRxiv. 2020 [[Google Scholar](#)]
24. Elfiky AA, Ibrahim NS. Anti-SARS and Anti-HCV Drugs Repurposing Against the Papain-like Protease of the Newly Emerged Coronavirus (2019-nCoV) 2020 [[Google Scholar](#)]
25. Arya R, Das A, Prashar V, Kumar M. Potential inhibitors against papain-like protease of novel coronavirus (SARS-CoV-2) from FDA approved drugs. 2020 [[Google Scholar](#)]
26. Smith M, Smith JC. Repurposing Therapeutics for COVID-19: Supercomputer-Based Docking to the SARS-CoV-2 Viral Spike Protein and Viral Spike Protein-Human ACE2 Interface. 2020 [[Google Scholar](#)]

27. Li Z, Bai T, Yang L, Hou X. Discovery of potential drugs for COVID-19 based on the [\[Google Scholar\]](#)
28. Richardson P, Griffin I, Tucker C, Smith D, Oechsle O, Phelan A, et al. Baricitinib as potential treatment for 2019-nCoV acute respiratory disease. *The Lancet*. 2020;395(10223):e30–e1. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
29. Nowak JK, Walkowiak J. Is lithium a potential treatment for the novel Wuhan (2019-nCoV) coronavirus? A scoping review. *F1000Research*. 2020;9(93) [\[Google Scholar\]](#)
30. Sun M, Yang J, Sun Y, Su G. Inhibitors of RAS Might Be a Good Choice for the Therapy of COVID-19 Pneumonia. *Zhonghua jie he he hu xi za zhi= Zhonghua jiehe he huxi zazhi= Chinese journal of tuberculosis and respiratory diseases*. 2020;43:E014. [\[PubMed\]](#) [\[Google Scholar\]](#)

Articles from Archives of Academic Emergency Medicine are provided here courtesy of **Shahid Beheshti University of Medical Sciences**