

Improved package of practice for Dhaincha



Dr. Girija Prasad Patnaik

National Consultant

Sustainable Impact for Rice based Systems

International Rice Research Institute



DHAINCHA

Seed production



Land requirement

- Free of volunteer plants.
- Previous crop **not be the same variety or other varieties** of the same crop.
- It can be the same variety if it is certified as per the procedures of certification agency.




Season

- *Rabi* and summer are best seasons
- October-November sowing will give better seed yield
- Can be cultivated in *kharif* with good soil moisture content


Isolation distance

- Foundation seed production - 10 m
- Certified seed production - 5 m
- From all sides




Seed treatment

- Rhizobium 10 g/kg of seed
- Carbendazim @ 2 g /kg of seeds



Seed rate and spacing

- 20 - 30 kg / ha
- Spacing: 60 x 10-20 cm



Nutrient management

Dose: 20-30:40:20 kg/ha			
Fertilizer sources	Per acre	Per hectare	Time of application
Lime application	200 kg	500 kg	Basal
Urea	5 kg	12.5 kg	
DAP	35 kg	87.5 kg	
MOP	16 kg	40 kg	
Foliar spray of DAP (2%)			35 DAS 45 DAS




Irrigation management

- Three irrigation one each at 3 DAS, at flowering and at pod development stages.
- Controlled flooding irrigation

Weed management

- Imazethapyr @ 0.075 kg a.i. ha⁻¹ at 5 DAS + hand weeding on 15 DAS
- Spray Neem oil 2% to control pod borer



Rain-gun system in Dhanicha



Seed grading

- Grade the seeds with BSS 7 x 7 wire mesh sieve

Storage

- Store the seeds in gunny or cloth bags for short term storage (8 - 9 months) with a seed moisture content of 9 %
- Store the seeds in 700-gauge polythene bag for long term storage (more than 15 months) with a seed moisture content of 8 %.





Flower bud



Pod initiation



Pod development



Root nodules

DHAINCHA

In-situ Green manuring



Dhaincha (Green manuring *In-situ*)

Time of sowing

- Best time for sowing is just after pre-monsoon showers or with good soil moisture it can be grown in all seasons

Soil

- Grown in all soil conditions, suitable for medium and lowland soils

Seed rate

- 40 kg/ha, seed is broadcasted in soil

Seed treatment

- Rhizobium @ 10g/kg seed



Nutrient management

- Lime application @ 500 kg/ha broadcasted just after incorporation of green manure and before final puddling
- The entire quantity of phosphatic fertilizer (DAP) of the succeeding rice crop can be applied to dhaincha. So that succeeding rice crop gets the N and P from the decomposing green manure.

Irrigation

- Irrigation is given once in 15 – 20 days



Incorporation of Dhaincha

Time of incorporation: **45-50 DAS** or **at flower bud stage**. It is incorporated with the help of plough and water is impounded for 2-3 days followed by final puddling.

Yield

Green biomass – 20-25 t/ha

Flower bud stage





Dhaincha incorporation

