

MTU 1010 (Cottondora Sannalu)* (Krishnaveni x IR64)

Product Profile

A high-yielding, early-maturing, medium-tall variety with medium-slender heavy grains, this mega variety is cultivated in India under irrigated lowland conditions. Developed and released in 1999 by ANGRAU University (Maruteru), MTU-1010 resulted from the hybridization of a local variety 'Krishnaveni' with IR64, a popular rice variety from IRRI. Known for its excellent tillering capacity and sturdy stem, it demonstrates susceptibility to leaf and sheath blight, as well as BPH. However, it shows tolerance to leaf blast, which enhances its resilience against this particular disease.

Priority Market Segment - TEMS-R

Planting Method		Ecosystem		
PTR	DSR	Irrigated Lowland	Rainfed Lowland	Upland
Recommended	Not Recommended	Recommended	Not Recommended	Not Recommended

Positioning

Position	Irrigated Lowland	Rainfed Lowland	Upland
Replace	IR-64	None	None
Companion	MTU-1001	None	None

Key Strengths

- Good yield
- Medium tall; Sturdy stem
- Medium slender and heavy grain
- Early maturity
- Tolerant to leaf blast
- Excellent productive tillers

Considerations

- Susceptible to leaf and sheath blight and BPH
- Average performance in low phosphorus soil
- Prone to shattering

Relative Variety Performance - Grain Yield, Agronomic Traits and Grain Shape

Variety	Grain yield		Maturity (days)	Plant height (cm)	Lodging	HRR %	1000 grain weight (g)	Grain Shape
	(Kg/ha)	% Adv						
MTU 1010 (Cottondora Sannalu)	4000-4500	5	120-125	105-108	Tolerant	58	25	MS
IR-64	4000-4300		115-120	100-105	Tolerant	49	23	LS

Relative Variety Performance - Biotic and Abiotic Stress

Variety	Leaf blast	Leaf blight	Sheath blight	BLB	Drought
MTU 1010 (Cottondora Sannalu)	Tolerant	Susceptible	Susceptible	Susceptible	Susceptible
IR-64	Tolerant	Resistant	Susceptible	Tolerant	Susceptible

Producibility: MTU 1010 (Cottondora Sannalu)= 1600-1800Kgs. / acre

Recommendation: Released for cultivation in Andhra Pradesh, Telangana, Odisha&west Bengal.Zonal check in advance varietal trial (mid-early)in central and southern India.



*Parental line (s) from IRRI germplasm