

**Homework 1**

STA4322/STA5328

Introduction to Statistics Theory

Spring 2020, MWF 3:00pm

Due date: Wednesday, January 22nd, 2020

All work must be shown for complete credit. W.M.S. denotes the course textbook, *Mathematical Statistics with Applications, 7th Ed.*

1. Let  $X_1, X_2, \dots, X_n$  be independent random variables following the uniform distribution on the interval  $[0, \theta]$  for some  $\theta > 0$ .
  - (a) Find the probability distribution function of  $X_{(n)} = \max(X_1, X_2, \dots, X_n)$ .
  - (b) Find the probability density function of  $X_{(n)}$ .
  - (c) Compute the mean and variance of  $X_{(n)}$ .
2. As in Problem 1, let  $X_1, X_2, \dots, X_n$  be independent random variables following the uniform distribution on the interval  $[0, \theta]$  for some  $\theta > 0$ . Find the density of  $X_{(k)}$ , the  $k$ th order statistic, where  $k$  is an integer between 1 and  $n$ .
3. (WMS 8.12) Let  $Y_1, \dots, Y_n$  be a random sample where for  $i = 1, \dots, n$ , we know  $Y_i \sim \text{Uniform}(\theta, \theta + 1)$  where  $\theta > 0$  is an unknown parameter.
  - (a) Show that  $\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i$  is a *biased* estimator of  $\theta$ .
  - (b) Propose an estimator of  $\theta$  which is unbiased. Verify that this estimator is unbiased.
  - (c) Compute the MSE of  $\bar{Y}$  with respect to  $\theta$ , i.e., compute  $\text{MSE}_\theta(\bar{Y})$
4. Suppose that the random variables  $X_1$  and  $X_2$  are independent and that both have the normal distribution with mean  $\mu$  and variance  $\sigma^2$ . Prove that  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$  and  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$  are independent.

**Hint:** Let  $Y_1 = X_1 + X_2$  and  $Y_2 = X_1 - X_2$ . Show that  $\bar{X}$  and  $s^2$ , respectively, can be written in terms of  $Y_1$  and  $Y_2$  alone. Then, verify that  $Y_1$  and  $Y_2$  are independent using that if two normally distributed random variables have zero covariance, they are independent.

5. Suppose that a random sample  $X_1, X_2, \dots, X_n$  is to be taken from the uniform distribution on the interval  $[0, \phi]$  with  $\phi > 0$  unknown. What  $n$  is required so that

$$P(|\max(X_1, X_2, \dots, X_n) - \phi| < 0.1\phi) \geq 0.95$$

for all possible  $\phi$ ?