References for Project Proposal

[1] Hutchinson, G.E. (1959) Homage to santa rosalia or why are there so many kinds of animals? *Am. Nat.* 93, 145–159

[2] Prates, I. *et al.* (2016) Inferring responses to climate dynamics from historical demography in neotropical forest lizards. *Proc. Natl. Acad. Sci. U. S. A.* 113, 7978–7985

[3] Carnaval, A.C. *et al.* (2014) Prediction of phylogeographic endemism in an environmentally complex biome. *Proc. Biol. Sci.* 281

[4] Barnosky, A.D. *et al.* (2011) Has the earth's sixth mass extinction already arrived? *Nature* 471, 51–57

[5] Erwin, D.H. (2015) *Extinction: How Life on Earth Nearly Ended 250 Million Years Ago*. Princeton University Press

[6] MacArthur, R.H. and Wilson, E.O. (1967) *The theory of island biogeography*. Princeton University Press

[7] Hubbell, S.P. (2001) *The unified neutral theory of biodiversity and biogeography*, vol. 32. Princeton University Press

[8] Gillespie, R. (2004) Community assembly through adaptive radiation in hawaiian spiders. *Science* 303, 356–359

[9] Owens, H.L. *et al.* (2017) The latitudinal diversity gradient in new world swallowtail butterflies is caused by contrasting patterns of out-of- and into-the-tropics dispersal. *Glob. Ecol. Biogeogr.* 26, 1447–1458

[10] Janzen, D.H. (1967) Why mountain passes are higher in the tropics. *Am. Nat.* 101, 233–249

[11] Mittelbach, G.G. *et al.* (2007) Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecol. Lett.* 10, 315–331

[12] Pianka, E.R. (1966) Latitudinal gradients in species diversity: A review of concepts. *Am. Nat.* 100, 33–46

[13] Ricklefs, R.E. (2006) The unified neutral theory of biodiversity: do the numbers add up? *Ecology* 87, 1424–1431

[14] Chisholm, R.A. and O'Dwyer, J.P. (2014) Species ages in neutral biodiversity models. *Theor. Popul. Biol.* 93, 85–94

[15] Rosindell, J. *et al.* (2015) Unifying ecology and macroevolution with individual-based theory. *Ecol. Lett.* 18, 472–482

[16] Rosindell, J. *et al.* (2012) The case for ecological neutral theory. *Trends Ecol. Evol.* 27, 203–208

[17] Chisholm, R.A. and Pacala, S.W. (2010) Niche and neutral models predict asymptotically equivalent species abundance distributions in high-diversity ecological communities. *Proc. Natl. Acad. Sci. U. S. A.* 107, 15821–15825

[18] McGill, B. (2003) Strong and weak tests of macroecological theory. *Oikos* 102, 679–685

[19] McGill, B.J. *et al.* (2007) Species abundance distributions: moving beyond single prediction theories to integration within an ecological framework. *Ecol. Lett.* 10, 995–1015

[20] Leibold, M.A. and Chase, J.M. (2017) *Metacommunity Ecology*. Princeton University Press

[21] Rominger, A.J. *et al.* Linking evolutionary and ecological theory illuminates non-equilibrium biodiversity. *Trends Ecol. Evol.*

[22] Etienne, R.S. (2007) A neutral sampling formula for multiple samples and an 'exact' test of neutrality. *Ecology Letters* 10, 608–618

[23] Chesson, P. (2000) Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* 31, 343–366

[24] Ricklefs, R.E. and Bermingham, E. (2001) Nonequilibrium diversity dynamics of the lesser antillean avifauna. *Science* 294, 1522–1524

[25] Qian, H. *et al.* (2005) Beta diversity of angiosperms in temperate floras of eastern asia and eastern north america. *Ecol. Lett.* 8, 15–22

[26] Yu, X.Q. *et al.* (2017) Insights into the historical assembly of east asian subtropical evergreen broadleaved forests revealed by the temporal history of the tea family. *New Phytol.* 215, 1235–1248

[27] Carnaval, A.C. and Moritz, C. (2008) Historical climate modelling predicts patterns of current biodiversity in the brazilian atlantic forest. *J. Biogeogr.* 35, 1187–1201

[28] Carnaval, A.C. *et al.* (2009) Stability predicts genetic diversity in the brazilian atlantic forest hotspot. *Science* 323, 785–789

[29] Burbrink, F.T. *et al.* (2016) Asynchronous demographic responses to pleistocene climate change in eastern nearctic vertebrates. *Ecol. Lett.* 19, 1457–1467

[30] Barnosky, A.D. *et al.* (2012) Approaching a state shift in earth/'s biosphere. *Nature* 486, 52–58

[31] Guralnick, R. *et al.* (2017) Humboldt core – toward a standardized capture of biological inventories for biodiversity monitoring, modeling and assessment. *Ecography*

[32] Overcast, I. *et al.* An integrated model of population genetics and community ecology. *Mol. Ecol.*

[33] Beaumont, M.A. (2010) Approximate bayesian computation in evolution and ecology. *Annu. Rev. Ecol. Evol. Syst.*

[34] Rosindell, J. *et al.* (2010) Protracted speciation revitalizes the neutral theory of biodiversity. *Ecol. Lett.* 13, 716–727

[35] Rosindell, J. and Phillimore, A.B. (2011) A unified model of island biogeography sheds light on the zone of radiation. *Ecol. Lett.* 14, 552–560

[36] Haegeman, B. and Etienne, R.S. (2017) A general sampling formula for community structure data. *Methods Ecol. Evol.*

[37] Grilli, J. *et al.* (2017) Higher-order interactions stabilize dynamics in competitive network models. *Nature* 548, 210–213

[38] HilleRisLambers, J. *et al.* (2012) Rethinking community assembly through the lens of coexistence theory. *Annual Review of*

[39] Adler, P.B. *et al.* (2010) Coexistence of perennial plants: an embarrassment of niches. *Ecol. Lett.* , no–no

[40] Jabot, F. *et al.* (2008) Reconciling neutral community models and environmental filtering: theory and an empirical test. *Oikos* 117, 1308–1320

[41] Tuomisto, H. *et al.* (2003) Dispersal, environment, and floristic variation of western amazonian forests. *Science* 299, 241–244

[42] Schrider, D.R. and Kern, A.D. (2018) Supervised machine learning for population genetics: A new paradigm. *Trends Genet.*

[43] R Development Core, T. (2013) R: A language and environment for statistical computing. *Vienna, Austria*

[44] Losos, J.B. and Schluter, D. (2000) Analysis of an evolutionary species–area relationship. *Nature* 408, 847

[45] Wagner, C.E. *et al.* (2014) Cichlid species-area relationships are shaped by adaptive radiations that scale with area. *Ecol. Lett.* 17, 583–592

[46] Ogden, R. and Thorpe, R.S. (2002) Molecular evidence for ecological speciation in tropical habitats. *Proc. Natl. Acad. Sci. U. S. A.* 99, 13612–13615

[47] Rundle, H.D. and Nosil, P. (2005) Ecological speciation. *Ecol. Lett.* 8, 336–352

[48] May, R.M. (1973) Qualitative stability in model ecosystems. *Ecology* 54, 638–641

[49] Nuismer, S.L. *et al.* (2013) Coevolution and the architecture of mutualistic networks. *Evolution* 67, 338–354

[50] Quental, T.B. and Marshall, C.R. (2013) How the red queen drives terrestrial mammals to extinction. *Science*

[51] O'Dwyer, J.P. and Chisholm, R. (2014) A mean field model for competition: From neutral ecology to the red queen. *Ecol. Lett.* 17, 961–969

[52] Rabosky, D.L. (2009) Ecological limits and diversification rate: alternative paradigms to explain the variation in species richness among clades and regions. *Ecol. Lett.* 12, 735–743

[53] Rominger, A.J. *et al.* (2015) Community assembly on isolated islands: macroecology meets evolution. *Glob. Ecol. Biogeogr.*

[54] Gillespie, R.G. and Baldwin, B.G. (2010) Island biogeography of remote archipelagoes. *The theory of island biogeography revisited* , 358–387

[55] Rominger, A.J. and Huang, E.G. (2016). hdimDB: an R package for biodiversity database management with google sheets. `https://github.com/hawaiiDimensions/db/tree/master/hdimDB`

[56] RStudio, Inc (2013) *Easy web applications in R*

[57] Wickham, H. and Francois, R. (2015) dplyr: A grammar of data manipulation. *R package version 0. 4* 3

[58] Flemons, P. *et al.* (2007) A web-based GIS tool for exploring the world's biodiversity: The global biodiversity information facility mapping and analysis portal application (GBIF-MAPA). *Ecol. Inform.* 2, 49–60

[59] Sherry, S.T. *et al.* (2001) dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.* 29, 308–311

[60] Benson, D. *et al.* (1993) GenBank. *Nucleic Acids Res.* 21, 2963–2965

[61] Hinchliff, C.E. *et al.* (2015) Synthesis of phylogeny and taxonomy into a comprehensive tree of life. *Proc. Natl. Acad. Sci. U. S. A.* 112, 12764–12769

[62] Jetz, W. *et al.* (2012) Integrating biodiversity distribution knowledge: toward a global map of life. *Trends Ecol. Evol.* 27, 151–159

[63] Bartomeus, I. and Molina, P.J. (2018). TraitBase. `https://traitbase.info/`

[64] Poelen, J.H. *et al.* (2014) Global biotic interactions: An open infrastructure to share and analyze species-interaction datasets. *Ecol. Inform.* 24, 148–159

[65] Chamberlain, S. *et al.* (2014) rgbif: Interface to the global biodiversity information facility API. *R package version 0. 7* 7

[66] Winter, D.J. (2017) *rentrez: An R package for the NCBI eUtils API. Tech. Rep. e3179v2*, PeerJ Preprints

[67] Michonneau, F. *et al.* (2016) rotl: an R package to interact with the open tree of life data. *Methods Ecol. Evol.* 7, 1476–1481

[68] Molina, P.J. and Bartomeus, I. (2017). traitbaser: R package providing API access to traitbase https://www.traitbase.info. `https://github.com/metadevpro/traitbaser`

[69] Poelen, J. *et al.* (2018). rglobi: R interface to global biotic interactions

[70] Douglas, K. and Douglas, S. (2003) *PostgreSQL: A Comprehensive Guide to Building, Programming, and Administering PostgresSQL Databases*. Sams Publishing

[71] Brown, J.H. *et al.* (2004) Toward a metabolic theory of ecology. *Ecology* 85, 1771–1789

[72] Price, C.A. *et al.* (2012) Testing the metabolic theory of ecology. *Ecology letters* 15, 1465–1474

[73] Elith, J. and Leathwick, J.R. (2009) Species distribution models: ecological explanation and prediction across space and time. *Annual review of ecology, evolution, and systematics* 40, 677–697

[74] Krehenwinkel, H. *et al.* (2017) Estimating and mitigating amplification bias in qualitative and quantitative arthropod metabarcoding. *Sci. Rep.* 7, 17668

[75] Che, J. *et al.* (2012) Universal coi primers for dna barcoding amphibians. *Molecular Ecology Resources* 12, 247–258

[76] Bi, Y. *et al.* (2018) Chloroplast genomic resources for phylogeny and dna barcoding: a case study on fritillaria. *Scientific reports* 8, 1184

[77] Lopez-Gonzalez, G. *et al.* (2011) ForestPlots.net: a web application and research tool to manage and analyse tropical forest plot data. *J. Veg. Sci.* 22, 610–613

[78] ter Steege, H. and et al. (2011). Amazon tree diversity network. `http://atdn.myspecies.info/`

[79] RAINFOR (2018). Amazon forest inventory network. `http://www.rainfor.org/`

[80] Webb, C.O. *et al.* (2008) Phylocom: software for the analysis of phylogenetic community structure and trait evolution. *Bioinformatics* 24

[81] Harmon, L.J. *et al.* (2013) Arbor: Comparative analysis workflows for the tree of LifeCall for SubmissionsAVAToL CollectionBlog PostTwitter. *PLoS Curr.*

[82] Rosindell, J. and Harmon, L.J. (2013) A unified model of species immigration, extinction and abundance on islands. *J. Biogeogr.* 40, 1107–1118

[83] Felsenstein, J. (2012) A comparative method for both discrete and continuous characters using the threshold model. *Am. Nat.* 179, 145–156

[84] Wilson, W.G. *et al.* (2003) Biodiversity and species interactions: extending Lotka–Volterra community theory. *Ecol. Lett.* 6, 944–952

[85] Kendall, D.G. (1948) On some modes of population growth leading to r. a. fisher's logarithmic series distribution. *Biometrika* 35, 6–15

[86] Rosenberg, N.A. and Nordborg, M. (2002) Genealogical trees, coalescent theory and the analysis of genetic polymorphisms. *Nat. Rev. Genet.* 3, 380–390

[87] Waples, R.S. and Yokota, M. (2007) Temporal estimates of effective population size in species with overlapping generations. *Genetics* 175, 219–233

[88] Kelleher, J. *et al.* (2016) Efficient coalescent simulation and genealogical analysis for large sample sizes. *PLoS Comput. Biol.* 12, e1004842

[89] Anderson, J. *et al.* (2014) The more, the merrier: the blessing of dimensionality for learning large gaussian mixtures. In *Conference on Learning Theory.* jmlr.org, pp. 1135–1164

[90] Breiman, L. (2001) Statistical modeling: The two cultures (with comments and a rejoinder by the author). *Stat. Sci.* 16, 199–231

[91] Chan, J. *et al.* (2018) A Likelihood-Free inference framework for population genetic data using exchangeable neural networks

[92] Tingley, M.W. and Beissinger, S.R. (2009) Detecting range shifts from historical species occurrences: new perspectives on old data. *Trends Ecol. Evol.* 24, 625–633

[93] May, F. *et al.* (2017) mobsim: An R package for the simulation and measurement of biodiversity across spatial scales

[94] Chao, A. *et al.* (2010) Phylogenetic diversity measures based on hill numbers. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 365, 3599–3609

[95] Karp, D.S. *et al.* (2012) Intensive agriculture erodes ?-diversity at large scales. *Ecol. Lett.* 15

[96] Barnosky, A.D. *et al.* (2001) Temperate terrestrial vertebrate faunas in north and south america: Interplay of ecology, evolution, and geography with biodiversity. *Conserv. Biol.* 15, 658–674

[97] do Amaral, F.R. *et al.* (2016) The "atlantis forest hypothesis" does not explain atlantic forest phylogeography. *Proc. Natl. Acad. Sci. U. S. A.* 113, E2097–E2098

[98] Prates, I. *et al.* (2016) A mid-pleistocene rainforest corridor enabled synchronous invasions of the atlantic forest by amazonian anole lizards. *Mol. Ecol.* 25, 5174–5186

[99] Maestri, R. *et al.* (2016) Predictors of intraspecific morphological variability in a tropical hotspot: comparing the influence of random and non-random factors. *J. Biogeogr.* 43, 2160–2172

[100] Zamborlini Saiter, F. *et al.* (2016) Environmental correlates of floristic regions and plant turnover in the atlantic forest hotspot. *J. Biogeogr.* 43, 2322–2331

[101] Montade, V. *et al.* (2016) Pollen-based characterization of montane forest types in northeastern brazil. *Rev. Palaeobot. Palynol.* 234, 147–158

[102] Bernal, J.P. *et al.* (2016) High-resolution holocene south american monsoon history recorded by a speleothem from botuverá cave, brazil. *Earth Planet. Sci. Lett.* 450, 186–196

[103] Bustamante, M.G. *et al.* (2016) Holocene changes in monsoon precipitation in the andes of NE peru based on $\delta18O$ speleothem records. *Quat. Sci. Rev.* 146, 274–287

[104] Gu, F. *et al.* (2017) Long-term vegetation, climate and ocean dynamics inferred from a 73,500 years old marine sediment core (GeoB2107-3) off southern brazil. *Quat. Sci. Rev.* 172, 55–71

[105] Dawson, M.N. (2016) Island and island-like marine environments. *Glob. Ecol. Biogeogr.* 25, 831–846

[106] Dawson, M.N. *et al.* (2016) Biogeography of islands, lakes, and mountaintops; evolutionary

[107] Meyerhof, M.S. *et al.* (2016) Microbial community diversity, structure and assembly across oxygen gradients in meromictic marine lakes, palau. *Environ. Microbiol.* 18, 4907–4919

[108] Schiebelhut, L.M. *et al.* (2017) A comparison of DNA extraction methods for high-throughput DNA analyses. *Mol. Ecol. Resour.* 17, 721–729

[109] Swift, H.F. *et al.* (2016) Three routes to crypsis: Stasis, convergence, and parallelism in the mastigias species complex (scyphozoa, rhizostomeae). *Mol. Phylogenet. Evol.* 99, 103–115

[110] Wilson, J. *et al.* (2017) Primary production, community respiration, and net community production along oxygen and nutrient gradients: Environmental controls and biogeochemical feedbacks within and across "marine lakes". *Frontiers in Marine Science* 4, 12

[111] Brewer, M.S. *et al.* (2015) Shifting habitats, morphology, and selective pressures: developmental polyphenism in an adaptive radiation of hawaiian spiders. *Evolution* 69, 162–178

[112] Brewer, M.S. *et al.* (2014) New sequencing technologies, the development of genomics tools, and their applications in evolutionary arachnology. *J. Arachnol.* 42, 1–15

[113] Gillespie, R.G. (2013) Adaptive radiation: Convergence and non-equilibrium. *Curr. Biol.* 23, R71–R74

[114] Shaw, K.L. and Gillespie, R.G. (2016) Comparative phylogeography of oceanic archipelagos: Hotspots for inferences of evolutionary process. *Proc. Natl. Acad. Sci. U. S. A.* 113, 7986–7993

[115] Gillespie, R.G. (2016) Island time and the interplay between ecology and evolution in species diversification. *Evol. Appl.* 9, 53–73

[116] Gillespie, R.G. and Roderick, G.K. (2014) Evolution: Geology and climate drive diversification. *Nature* 509, 297–298

[117] Krehenwinkel, H. *et al.* (2017) A cost-efficient and simple protocol to enrich prey DNA from extractions of predatory arthropods for large-scale gut content analysis by illumina sequencing. *Methods Ecol. Evol.* 8, 126–134

[118] Warren, B.H. *et al.* (2015) Islands as model systems in ecology and evolution: prospects fifty years after MacArthur-Wilson. *Ecol. Lett.* 18, 200–217

[119] Yim, K.M. *et al.* (2014) Comparative transcriptomics of maturity-associated color change in hawaiian spiders. *J. Hered.* 105 Suppl 1, 771–781

[120] Krehenwinkel, H. *et al.* (2018) The effect of DNA degradation bias in passive sampling devices on metabarcoding studies of arthropod communities and their associated microbiota. *PLoS One* 13, e0189188

[121] Graham, N.R. *et al.* (2017) Island ecology and evolution: challenges in the anthropocene. *Environ. Conserv.* 44, 323–335

[122] Guralnick, R.P. *et al.* (2007) Towards a collaborative, global infrastructure for biodiversity assessment. *Ecol. Lett.* 10, 663–672

[123] Guralnick, R. *et al.* (2009) Sharing: lessons from natural history's success story. *Nature* 462, 34

[124] Guralnick, R. and Constable, H. (2010) VertNet: Creating a data-sharing community. *Bioscience* 60, 258–259

[125] Parr, C.S. *et al.* (2012) Evolutionary informatics: unifying knowledge about the diversity of life. *Trends Ecol. Evol.* 27, 94–103

[126] Stucky, B.J. *et al.* (2014) The BiSciCol triplifier: bringing biodiversity data to the semantic web. *BMC Bioinformatics* 15, 257

[127] Yilmaz, P. *et al.* (2011) Minimum information about a marker gene sequence (MIMARKS) and minimum information about any (x) sequence (MIxS) specifications. *Nat. Biotechnol.* 29, 415

[128] Overcast, I. *et al.* (2017) Strategies for improving approximate bayesian computation tests for synchronous diversification. *BMC Evol. Biol.* 17, 203

[129] Xue, A.T. and Hickerson, M.J. (2017) MultiDICE: R package for comparative population genomic inference under hierarchical codemographic models of independent singlepopulation size changes. *Mol. Ecol. Resour.*

[130] Emerson, B.C. *et al.* (2015) Model misspecification confounds the estimation of rates and exaggerates their time dependency. *Mol. Ecol.* 24, 6013–6020

[131] Myers, E.A. *et al.* (2016) Asynchronous diversification of snakes in the north american warm deserts. *J. Biogeogr.*

[132] Boehm, J.T. *et al.* (2016) Urbanization shapes the demographic history of a native rodent (the white-footed mouse, peromyscus leucopus) in new york city. *Biology*

[133] Joseph, T.A. *et al.* (2016) Demographic inference under a spatially continuous coalescent model. *Heredity* 117, 94–99

[134] Burbrink, F.T. *et al.* (2016) Asynchronous demographic responses to pleistocene climate change in eastern nearctic vertebrates. *Ecol. Lett.*

[135] Brown, J.L. *et al.* (2016) Predicting the genetic consequences of future climate change: The power of coupling spatial demography, the coalescent, and historical landscape changes. *Am. J. Bot.* 103, 153–163

[136] Prates, I. *et al.* (2016) Inferring responses to climate dynamics from historical demography in neotropical forest lizards. *Proceedings of the National Academy of Sciences* 113, 7978–7985

[137] Demos, T.C. *et al.* (2015) Comparative population genomics of african montane forest mammals support population persistence across a climatic gradient and quaternary climatic cycles. *PLoS One* 10, e0131800

[138] Lipshutz, S.E. *et al.* (2017) Behavioural response to song and genetic divergence in two subspecies of white-crowned sparrows (zonotrichia leucophrys). *Mol. Ecol.* 26, 3011–3027

[139] Harris, S.E. *et al.* (2016) Urbanization shapes the demographic history of a native rodent (the white-footed mouse, peromyscus leucopus) in new york city. *Biol. Lett.* 12

[140] Alvarado-Serrano, D.F. and Hickerson, M.J. (2015) Spatially explicit summary statistics for historical population genetic inference. *Methods Ecol. Evol.* 7, 418–427

[141] Xue, A.T. and Hickerson, M.J. (2015) The aggregate site frequency spectrum (aSFS) for comparative population genomic inference. *Mol. Ecol.*

[142] Emerson, B.C. and Hickerson, M.J. (2015) Lack of support for the time-dependent molecular evolution hypothesis. *Mol. Ecol.* 24, 702–709

[143] Boehm, J.T. *et al.* (2015) Population genomics reveals seahorses (hippocampus erectus) of the western Mid-Atlantic coast to be residents rather than vagrants. *PLoS One* 10, e0116219

[144] Robinson, J.D. *et al.* (2014) Sampling strategies for frequency spectrum-based population genomic inference. *BMC Evol. Biol.* 14, 254

[145] Robinson, J.D. *et al.* (2014) ABC inference of multipopulation divergence with admixture from unphased population genomic data. *Mol. Ecol.* 23, 4458–4471

[146] Smith, B.T. *et al.* (2014) The drivers of tropical speciation. *Nature*

[147] Chan, Y.L. *et al.* (2014) Detecting concerted demographic response across community assemblages using hierarchical approximate bayesian computation. *Mol. Biol. Evol.* , msu187

[148] Demos, T.C. *et al.* (2014) Uncovering cryptic diversity and refugial persistence among small mammal lineages across the eastern afromontane biodiversity hotspot. *Mol. Phylogenet. Evol.* 71, 41–54

[149] Hickerson, M.J. *et al.* (2014) Recommendations for using msbayes to incorporate uncertainty in selecting an abc model prior: a response to oaks et al. *Evolution* 68, 284–294

References for Student Training Plan

[1] Woodin, T. *et al.* (2010) Vision and change in biology undergraduate education, a call for action–initial responses. *CBE Life Sci. Educ.* 9, 71–73

[2] Cook, J.A. *et al.* (2014) Natural history collections as emerging resources for innovative education. *BioScience* 64, 725–734

[3] Lacey, E.A. *et al.* (2017) Climate change, collections and the classroom: using big data to tackle big problems. *Evolution: Education and Outreach* 10, 2

[4] Goring, S.J. *et al.* (2014) Improving the culture of interdisciplinary collaboration in ecology by expanding measures of success. *Frontiers in Ecology and the Environment* 12, 39–47

[5] Wuchty, S. *et al.* (2007) The increasing dominance of teams in production of knowledge. *Science* 316, 1036–1039

[6] Cheruvelil, K.S. *et al.* (2014) Creating and maintaining high-performing collaborative research teams: the importance of diversity and interpersonal skills. *Frontiers in Ecology and the Environment* 12, 31–38

[7] Jones, B.F. *et al.* (2008) Multi-university research teams: Shifting impact, geography, and stratification in science. *Science* 322, 1259–1262