

Mathematical Correspondence Between the Free Fermion and Magnetized Partition Functions

Andrew Steinmetz,^{*} Cheng-Tao Yang, and Johann Rafelski
Department of Physics, The University of Arizona, Tucson, AZ 85721-0081, USA
(Dated: February 17, 2023)

Abstract. To be written.

I. INTRODUCTION

When exposed to external magnetic fields, matter can become magnetized enhancing or reducing the overall net field within the material. The origin of this magnetization is well known to be quantum mechanical in origin. The magnetic susceptibility χ is a measure of the material response given by

$$\mathbf{M} = \chi \mathbf{H}, \quad (1)$$

$$\mathbf{B} = \mu_{vac.} (\mathbf{H} + \mathbf{M}) = \mu_{vac.} (1 + \chi) \mathbf{H}, \quad (2)$$

where \mathbf{H} is the external magnetic field strength and \mathbf{B} is the total magnetic flux density. The magnetization \mathbf{M} is the magnetic moment density of the medium. The coefficient $\mu_{vac.}$ is the vacuum permeability, which with χ can define a material permeability $\mu_{mat.} = \mu_{vac.} (1 + \chi)$. Paramagnetism occurs when $\chi > 0$ which increases the magnetic flux as the magnetization \mathbf{M} is aligned with the external field. In contrast, diamagnetism is when $\chi < 0$ reducing the overall magnetic flux as the magnetization is anti-aligned with the external field. In the most general case, the susceptibility can be described by a tensor which is neither paramagnetic or diamagnetic. In this work, we will be considering only the simple case where the susceptibility is single-valued.

Before we handle an ensemble system, we will look at the magnetic moment $\boldsymbol{\mu}$ of a single-particle quantum system. To avoid confusion, the permeability will always be denoted either by a subscript for the vacuum or medium to always differentiate from magnetic moment. If we apply an external field with the local constant value of B which is primarily responsible for the particle's response, the magnetic moment can be evaluated from the matrix element

$$|\boldsymbol{\mu}_n| = - \left\langle n \left| \partial \hat{\mathcal{H}} / \partial B \right| n \right\rangle = - \frac{\partial E_n}{\partial B}, \quad (3)$$

where $\hat{\mathcal{H}}$ is the Hamiltonian of the system. It is valuable to point out that the individual dipole's response within a medium is only uniquely dependent on the external field \mathbf{H} in the case of weak medium magnetization. However, if the bulk magnetization is may easily be large and thus influential to each individual dipole. Physically each dipole is sensitive to the total magnetic flux \mathbf{B} which includes a mixture of external field and bulk magnetization \mathbf{M} from its neighbors. The magnetization density of the quantum system is then

$$M_n(B) = \frac{1}{V} |\boldsymbol{\mu}_n|. \quad (4)$$

If we couple the system to a thermal reservoir of temperature T , the averaged magnetization at thermal and chemical equilibrium is

$$M(B, T, \eta) = \frac{\sum_n M_n e^{-\beta(E_n - \eta N)}}{\sum_n e^{-\beta(E_n - \eta N)}}, \quad (5)$$

where $\beta = 1/k_B T$, k_B is the Boltzmann constant and η is the chemical potential. We then introduce the grand

^{*} ajsteinmetz@arizona.edu

potential Φ defined by

$$\Phi = -\frac{1}{\beta} \ln \left(\sum_n e^{-\beta(E_n - \eta N)} \right) = -\frac{1}{\beta} \ln(\mathcal{Z}) , \quad (6)$$

where \mathcal{Z} is the grand partition function. This allows us to rewrite eq. (5) as

$$M(B, T, \eta) = -\frac{1}{V} \frac{\partial \Phi}{\partial B} . \quad (7)$$

Combining eq. (6) and eq. (7) in the grand ensemble yields

$$M = \frac{1}{\beta V} \frac{\partial}{\partial B} \ln(\mathcal{Z}) . \quad (8)$$

The magnetic susceptibility is related to the magnetization via

$$\chi = \mu_{vac.} \frac{\partial M}{\partial B} . \quad (9)$$

If a given thermodynamic system is well described by a partition function, we can evaluate the susceptibility using eq. (9).

II. ENERGY EIGENVALUES

To evaluate the susceptibility of the charged Fermi gas submerged in a magnetic field, we need the energy eigenvalues of the system. We consider an external homogeneous magnetic field in the z-direction

$$\mathbf{B} = B\hat{z} , \quad \mathbf{A}_L = -By\hat{x} , \quad (10)$$

where we have specifically chosen the Landau gauge \mathbf{A}_L . Charged particles in homogeneous magnetic fields have their motion perpendicular to the direction of the magnetic field quantized into Landau orbits. Our next task is to choose our description of the quantum system. The second order KGP equation is given by

$$\left((i\hbar\partial_\mu - eA_\mu)^2 - m^2 - e\hbar\frac{g}{2}\frac{\sigma_{\mu\nu}F^{\mu\nu}}{2} \right) \Psi = 0 , \quad (11)$$

where $\sigma_{\mu\nu}$ is the spin tensor proportional to the commutator of the gamma matrices and $F^{\mu\nu}$ is the EM field tensor. The KGP equation is an alternate to the Dirac equation in describing spin one-half particles, though the two formulations are equivalent for the natural Dirac value of the g -factor of $g = 2$. The primary difference between the two is that the KGP is second order in derivatives and the magnetic moment of the particle is explicitly given by the Pauli term in the wave equation rather than implicitly present in the spinor structure of the Dirac equation.

The detailed solution to the Landau problem for the KGP particle is located in our prior work as well as a more detailed description of the KGP equation. The resulting energy levels are

$$E_{n,s}(p_z, B) = \sqrt{m^2 + p_z^2 + 2qB \left(n + \frac{1}{2} \right) - qBgs} , \quad (12)$$

where $n \in \mathbb{Z}^+$ for the Landau levels and $s \in \pm 1/2$ for the spin states. The magnitude of the electric charge of the particle is q , its momentum in the z-direction is p_z , and its mass is m . These energy levels differ from the related Dirac energies only by the presence of an arbitrary g -factor. The g -factor of the particle is defined in terms of the anomalous magnetic moment via

$$\frac{g}{2} = 1 + a . \quad (13)$$

We have put the above equation into natural units by setting $\hbar = c = 1$. As we are in the Landau gauge, these




FIG. 1. Qualitative value of the primordial magnetic field over the evolutionary lifespan of the universe. As magnetic flux is conserved over co-moving surfaces, the primordial relic field is expected to dilute as $B \propto 1/a(t)^2$ meaning the contemporary small bounded values of $5 \times 10^{-12} \text{ T} > B_{\text{relic}} > 10^{-20} \text{ T}$ may have once represented large magnetic fields in the early universe. This relic magnetic field would then be generated by the last phase of significant magnetization in the early universe. This figure is meant to be illustrative and it is unlikely the magnetization of the universe would proceed unhindered and unaltered into the ultra-dense plasma phases of the early universe. The values of the Schwinger critical field and the upper bound of surface magnetar field strength are included for scale.

states are also degenerate in p_x as it commutes with eq. (11). The degeneracy of the spin one-half Landau levels is broken for $g \neq 2$ except for values

$$g_k/2 = 1 + k, \quad k \in \mathbb{Z}, \quad (14)$$

where the degeneracy of the levels is restored. Using eq. (3) we can evaluate the KGP magnetic moment yielding

$$|\mu|_{p_z, n, s} = -\frac{1}{2E} (e\hbar c^2(2n + 1 - gs)). \quad (15)$$

From the numerator in eq. (15) we see that the magnetic moment is broken into two distinct parts: (a) an orbital part related to the Landau quantization and (b) a spin part containing the natural moment of the particle. This is the expected result as the overall magnetic moment of the system arises from the Amperian current of the particle completing the Landau orbits and the intrinsic moment present by virtue of its spin. As the moment is weighted in the denominator by the energy, the two kinds of magnetic moment cannot truly be separated out except under certain limits.

To better connect eq. (15) to our traditional understanding of magnetic moment, we introduce the following

substitutions (which also put the KGP equation into a Schoedinger-like form)

$$E = m'c^2, \quad \mu' = \frac{e\hbar}{2m'}, \quad (16)$$

yielding

$$|\boldsymbol{\mu}|_{p_z, n, s} = -\mu'(2n + 1 - gs). \quad (17)$$

Here we see μ' take on the roll of the effective magneton. In the non-relativistic limit, the coefficient becomes the Bohr magneton $\mu' \rightarrow \mu_B$. This limit also uniquely separates out the two contributions to the overall moment. We introduce an effective mass term which also incorporates the ground state component of the Landau energy

$$\tilde{m}_s^2(B) = m^2 + qB(1 - gs) > 0, \quad (18)$$

where we require that the effective mass remain positive definite. This effective mass is also spin dependent. This assumption breaks down for very strong magnetic fields

$$B_{\text{crit}} = \frac{m^2}{qa} = \frac{\mathcal{B}_S}{a}, \quad (19)$$

where \mathcal{B}_S is the Schwinger critical field for the particle.

III. LANDAU EIGEN-ENERGIES IN COSMOLOGY

The standard model of cosmology with a cosmological constant plus cold dark matter (Λ CDM) is described by the Friedman-Lemaître-Robertson-Walker (FLRW) metric. The spatially flat (Gaussian curvature $k = 0$) FLRW metric with metric signature $(+1, -1, -1, -1)$ in Cartesian coordinates is

$$ds^2 = dt^2 - a^2(t) [dx^2 + dy^2 + dz^2]. \quad (20)$$

The scale factor $a(t)$ denotes the change of proper distances in an expanding (or contracting) universe which is both homogeneous and isotropic. The evolutionary expansion of the universe is then traditionally defined in terms of the Hubble parameter $H(t)$ as follows

$$H(t)^2 \equiv \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G_N}{3} \rho_{\text{tot}}, \quad \frac{\ddot{a}}{a} = -qH^2, \quad q \equiv -\frac{a\ddot{a}}{\dot{a}^2}, \quad \dot{H} = -H^2(1 + q). \quad (21)$$

where G_N is the Newtonian constant of gravitation, ρ_{tot} is the total energy density of the universe and q is the cosmic deceleration parameter. Eq. (21) is also known as the Friedmann equations.

There is another natural scale for the magnetic field besides Eq. (??) when considering the consequences of FLRW expansion on the e^\pm fluid. As the universe expands, different terms in the energies and thus partition function evolve as a function of the scale factor $a(t)$ which arises in the FLRW metric. We can consider the expansion to be an adiabatic process which results in a smooth shifting of the relevant dynamical quantities. From the conservation of magnetic flux through a co-moving surface, the magnetic field under expansion starting at some initial time t_0 is given by

$$B(t) = B(t_0) \frac{a(t_0)^2}{a(t)^2}. \quad (22)$$

As the universe expands, the temperature also cools as the cosmological redshift reduces the momenta of particles in the universe lowering their contribution to the energy content of the universe. This cosmological redshift is written as

$$p_i(t) = p_i(t_0) \frac{a(t_0)}{a(t)}, \quad T(t) = T(t_0) \frac{a(t_0)}{a(t)}. \quad (23)$$

The momenta scale with the same factor as temperature as it is the origin of cosmological redshift. The energy of massive free particles in the universe scales differently based on their momentum (and thus temperature). When hot and relativistic, particle energy scales with inverse scale factors like radiation. However as particles transition to non-relativistic momenta, their energies scale with the inverse square of the scale factor like magnetic flux.

$$E(t) = E(t_0) \frac{a(t_0)}{a(t)} \xrightarrow{\text{NR}} E(t_0) \frac{a(t_0)^2}{a(t)^2}. \quad (24)$$

This occurs because of the functional dependence of energy on momentum in the relativistic versus non-relativistic cases. The argument in the Boltzmann statistical factor is given by

$$X_n^s \equiv \frac{E_n^s}{T}. \quad (25)$$

We can explore this relationship for the magnetized system explicitly by writing out Eq. (25) using the KGP eigen-energies as

$$X_n^s = \sqrt{\frac{m_e^2}{T^2} + \frac{p_z^2}{T^2} + \frac{2eB}{T^2} \left(n + \frac{1}{2} - \frac{gs}{2} \right)}, \quad (26)$$

where we now introduce the expansion scale factor via Eq. (22) - Eq. (??). The Boltzmann factor can then be written as

$$X_n^s[a(t)] = \sqrt{\frac{m_e^2}{T^2(t_0)} \frac{a(t)^2}{a(t_0)^2} + \frac{p_z^2(t_0)}{T^2(t_0)} + \frac{2eB(t_0)}{T^2(t_0)} \left(n + \frac{1}{2} - \frac{gs}{2} \right)}. \quad (27)$$

This reveals that only the mass contribution is dynamic over cosmological time. For any given eigen-state, the mass term increases driving the state into the non-relativistic limit while the momenta and magnetic contributions are frozen by initial conditions. As a point of comparison, the Boltzmann factor for the DP eigen-energies becomes

$$X_n^s|_{DP} = \sqrt{\left(\sqrt{\frac{m_e^2}{T^2} + \frac{2eB}{T^2} \left(n + \frac{1}{2} - s \right)} - \frac{eB}{2mT} (g-2)s \right)^2 + \frac{p_z^2}{T^2}}, \quad (28)$$

which over cosmological time under expansion scales as

$$X_n^s[a(t)]|_{DP} = \sqrt{\left(\sqrt{\frac{m_e^2}{T^2(t_0)} \frac{a(t)^2}{a(t_0)^2} + \frac{2eB(t_0)}{T^2(t_0)} \left(n + \frac{1}{2} - s \right)} - \frac{eB(t_0)}{2mT(t_0)} \frac{a(t_0)}{a(t)} (g-2)s \right)^2 + \frac{p_z^2(t_0)}{T^2(t_0)}}. \quad (29)$$

We note here one important difference between KGP and DP eigen-energies in the context of cosmology: The anomalous magnetic moment portion of the DP statistics is suppressed by $1/a(t)$ over cosmological time while the AMM contribution is preserved in the KGP model. That the universe's expansion makes a distinction between $g = 2$ magnetic moment and AMM for DP fermions appears as a rather artificial and nonphysical trait. While the suppression of AMM may often be small for particles such as electrons, this suppression is non-trivial for particles with large AMM values such as the proton. That cosmological redshift would push DP protons to be described by $g = 2$ eigen-energies in the non-relativistic limit counts as a malaise for the model and further strengthens our thinking that the KGP model is more appropriate for cosmological studies. Motivated by Eq. (27), we can introduce a dimensionless cosmic magnetic scale which is frozen in the homogeneous case as

$$b_0 \equiv \frac{eB}{T^2} = \frac{eB\hbar c^2}{(k_B T)^2} \text{ (S.I.)}, \quad (30)$$

where we've included the expression explicitly in full SI units. We can estimate the value of b_0 from the bounds on the extra-galactic magnetic field strength and the temperature of the universe today. If the origin of deep space extra-galactic magnetic fields are relic fields from the early universe, which today are expected to exist between

$5 \times 10^{-12} \text{ T} > B_{relic} > 10^{-20} \text{ T}$, then at temperature $T = 2.7 \text{ K}$, the value of the cosmic magnetic scale is between

$$5.5 \times 10^{-5} > b_0 > 1.1 \times 10^{-11}. \quad (31)$$

This should remain constant in the universe at-large up to the last epoch the universe was sufficiently magnetized to disturb this value. As the electron-proton plasma which generated the CMB was relatively dilute over its duration, it was unlikely sufficiently magnetized to significantly alter this value over extra-galactic scales. Rather, the best candidate plasma to have been sufficiently magnetized and dense to have set the relic field magnetic scale would have been the electron-positron plasma which existed during the duration of Big Bang Nucleosynthesis (BBN) and beforehand.

Higher order non-minimal magnetic contributions which can be introduced via Eq. (??) to the eigen-energies like $\approx \mu_B^2 B^2 / T^2$ are even more suppressed over cosmological time which drives the system into minimal electromagnetic coupling with the exception of the anomalous magnetic moment in the KGP case. It is interesting to note that cosmological expansion serves to “smooth out” the characteristics of more complex BSM electrodynmaics erasing them from a statistical perspective in favor of the minimal or minimal-like dynamics. As b_0 is a constant of expansion, assuming the electron-proton plasma between the CMB and electron-positron annihilation did not greatly disturbed it, we can calculate the remnant values at the temperature $T = 50 \text{ keV}$ with the expression

$$B(T) = \frac{b_0}{e} T^2, \quad (32)$$

yielding a range of field strengths

$$2.3 \times 10^5 \text{ T} > B(T = 50 \text{ keV}) > 4.6 \times 10^{-4} \text{ T}, \quad (33)$$

during which the electron-positron plasma in the universe had a number density comparable to that of the Solar core with $n_e = 4.49 \times 10^{24} \text{ cm}^{-3}$ at $r = 0.01 R_\odot$.

IV. PARTITION FUNCTION

Now that we’ve obtained the energy eigenvalues of the KGP particle, we can work towards evaluating the grand partition function of the system and ultimately the magnetic properties of the magnetized KGP Fermi gas. In addition to considering polarization, there are also the positive and negative energy states. The grand partition function for the relativistic Fermi-Dirac ensemble is given by the standard definition

$$\ln(\mathcal{Z}_L) = \sum_{\alpha} \ln(1 + z_{\sigma} \exp(-X_s)) , \quad (34)$$

where we are summing over all possible quantum numbers $\alpha = \{p_z, n, s, \sigma, \tilde{g}\}$. The summation over \tilde{g} represents the occupancy of Landau states which are matched to the available phase space within $\Delta p_x \Delta p_y$. The fugacity z_{σ} is defined as

$$z_{\sigma} = \exp(\sigma \eta) , \quad (35)$$

where η is the chemical potential of the species and $\sigma \in \pm 1$ denotes between particles and antiparticles. The chemical potential η is an increasing function of the number of particles present within a given volume and fixed by the particle number

$$N = \sum_{\alpha} \langle n_{\alpha} \rangle = \sum_{\alpha} \frac{1}{\exp(X_s) z_{\sigma}^{-1} + 1} . \quad (36)$$

If we consider the Landau energies to represent the transverse momentum $p_T^2 = p_x^2 + p_y^2$ of the system, then the relationship that defines \tilde{g} is given by

$$\frac{L^2}{(2\pi)^2} \Delta p_x \Delta p_y = \frac{qBL^2}{2\pi} \Delta n, \quad \tilde{g} = \frac{eBL^2}{2\pi} . \quad (37)$$

The summation over the continuous p_z can be replaced with an integration

$$\sum_{p_z} \rightarrow \frac{L}{2\pi} \int_{-\infty}^{+\infty} dp_z, \quad (38)$$

where L defines the boundary length of our considered volume. The partition function Eq. (34) can be then rewritten as

$$\ln(\mathcal{Z}_L) = \sum_{\sigma} \sum_s^{\pm 1} \frac{2eBL^3}{(2\pi)^2} \int_0^{+\infty} dp_z \sum_{n=0}^{\infty} \ln(1 + z_{\sigma} \exp(-X_s)). \quad (39)$$

We note that the partition function can be broken into four quantum gasses: Particles and antiparticles, and spin aligned and antialigned. This can be represented by separate partition functions $\ln(\mathcal{Z}_s^{\sigma})$ where

$$\ln(\mathcal{Z}_L) = \sum_{\sigma, s} \ln(\mathcal{Z}_s^{\sigma}), \quad (40)$$

For KGP particles, the anomalous moment (a) breaks the degeneracy between Landau levels of opposite alignment and (b) depending on the size of the anomaly, flips the sign of the magnetic contribution to the energy for certain eccentric states. For $0 < a < 2$ the aligned ground state is uniquely separated from the rest of the states while for $2 < a < 4$ the aligned ground state and first excited state are uniquely separated. The pattern of unique eccentric states continues for further extreme values of anomalous moment. While we consider the g -factor to be the immutable description of magnetic moment, it is useful in the case of Landau levels to consider the anomaly a as from the energies given in eq. (11), g always is linearly added or subtracted from the Landau quantum number which are integers.

Degeneracy is restored for values of the anomalous parameter given in eq. (14). While generating a large number of eccentric states from large anomalous moment may be of theoretical interest, it is of practical interest to consider particles like the proton or electron where only the ground state is uniquely disturbed. Due to the properties of logs, the overall partition function will be sum of the partition function of each species, which allows us to consider each species separately. The next step is to replace the sum over n Landau levels and replace it with an integration using the Euler-Maclaurin integration formula. The Euler-Maclaurin formula is given by

$$\sum_{n=a}^b f(n) - \int_a^b f(x) dx = \frac{1}{2} (f(b) + f(a)) + \sum_{i=1}^k \frac{b_{2i}}{(2i)!} \left(f^{(2i-1)}(b) - f^{(2i-1)}(a) \right) + R(k), \quad (41)$$

where b_n are the Bernoulli numbers and $R(k)$ is the error remainder defined by integrals over Bernoulli polynomials. The integer k is chosen for the level of approximation that is desired. Using Eq. (41) allows us to convert the sum over n quantum numbers in Eq. (39) into an integral. We define

$$f(p_z, n, s, \sigma) = \ln(1 + z_{\sigma} \exp(-X_s)), \quad (42)$$

The result for $k = 1$ is

$$\ln(\mathcal{Z}_s^{\sigma}) = \frac{2eBL^3}{(2\pi)^2} \int_0^{+\infty} dp_z \left(\int_0^{+\infty} dn f(n) + \frac{1}{2} f(0) + \frac{1}{12} \frac{\partial f(n)}{\partial n} \Big|_{n=0} + R(1) \right) \quad (43)$$

It is to be recognized that the first term in expression Eq. (43) is merely the free particle fermion partition function with a rescaled mass m_s . This then allows us to achieve our goal of a “separated” partition function which explicitly reveals the particle portion of the phase-phase and the spin portions as different contributions. This separation is however not fully independant as the rescaled mass m_s is intrinsically a function of the spin alignment. It will be demonstrated further below that the free-like portion is the relativistic equivalent of the Hamiltonian modified by $E_M = -\boldsymbol{\mu} \cdot \mathbf{B}$ dipole energies. We introduce the dimensionless variables

$$a_s = \frac{m_s}{T}, \quad y_s = \frac{p_z}{m_s}, \quad u_s = \frac{2qB}{m_s^2} n, \quad (44)$$

which allows us to express Eq. (25) as

$$X_s(y_s, u_s) = a_s \sqrt{1 + y_s^2 + u_s}. \quad (45)$$

Via substitution of dimensionless variables, and some rearrangement, we can write the magnetized partition function in terms of free fermion partition functions of differing masses, yielding,

$$\ln(\mathcal{Z}_s^\sigma) = \ln(\mathcal{Z}_F^\sigma)|_{m_s} + \frac{2eBL^3}{(2\pi)^2} m_s \int_0^{+\infty} dy \left(\frac{1}{2} f(0) + \frac{1}{12} \frac{\partial f(n)}{\partial n} \Big|_{n=0} + R(1) \right), \quad (46)$$

where we have dropped the spin subscript from the dimensionless z-momentum y as it does not impact the limits of integration. The free-like portion of the partition function is given by

$$\ln(\mathcal{Z}_F^\sigma)|_{m_s} = \frac{L^3}{(2\pi)^3} \int_{-\infty}^{+\infty} d\mathbf{p}^3 \ln(1 + z_\sigma \exp(-X_F^s)), \quad X_F^s = X^s|_{u_s=0} = a_s \sqrt{1 + y^2}, \quad (47)$$

In dimensionless units we can represent Eq. (47) as

$$\ln(\mathcal{Z}_F^\sigma)|_{m_s} = \frac{L^3}{(2\pi)^2} m_s^3 \int_{-\infty}^{+\infty} dy [y^2 \ln(1 + z_\sigma \exp(-X_F^s))], \quad (48)$$

Integrating by parts, we arrive at

$$\ln(\mathcal{Z}_F^\sigma)|_{m_s} = \frac{1}{3} \frac{L^3}{(2\pi)^2} m_s^4 \left(\frac{m_s}{T} \right) \int_{-\infty}^{+\infty} dy \left[\frac{y^4}{\epsilon} F[X_s, \sigma] \right], \quad \epsilon = m_s \sqrt{1 + y^2}, \quad (49)$$

$$F[X_s, \sigma] = \frac{1}{\exp(X_s) z_\sigma^{-1} + 1}, \quad (50)$$

where ϵ is the free eigen-energy and F is the Fermi particle(antiparticle) distribution. By introducing a change of variables of $y = \sinh(t)$ this simplifies to

$$\ln(\mathcal{Z}_F^\sigma)|_{m_s} = \frac{1}{3} \frac{L^3}{(2\pi)^2} m_s^4 \left(\frac{m_s}{T} \right) \int_{-\infty}^{+\infty} dt \left[\frac{\sinh(t)^4}{\exp(a_s \cosh(t)) z_\sigma^{-1} + 1} \right], \quad (51)$$

This integrand is well behaved and has a finite integration value within the restriction that m_s remains physical. The magnetization of this term, defined by Eq. (8), is found to be

$$M_F^{s,\sigma} = \frac{T}{V} \frac{\partial m_s}{\partial B} \frac{\partial}{\partial m_s} \ln(\mathcal{Z}_F^\sigma), \quad (52)$$

with the total magnetization given by the sum over the four species

$$M_F = \sum_{s,\sigma} M_F^{s,\sigma}. \quad (53)$$

The term by term magnetization evalutes as

$$M_F^{s,\sigma} = \frac{e(1-gs)}{2m_s} \left(\frac{5}{m_s} \ln(\mathcal{Z}_F^\sigma)|_{m_s} - \frac{1}{3} \frac{L^3}{(2\pi)^2} m_s^3 \left(\frac{m_s}{T} \right)^2 \int_{-\infty}^{+\infty} dt \left[z_\sigma^{-1} e^{a_s \cosh(t)} \cosh(t) \sinh^4(t) F^2[X_s, \sigma] \right] \right) \quad (54)$$