

```
> fitMed <- mediate(fitM, fitY, treat = "X", mediator = "M")
> summary(fitMed)
```

```
> summary(fitMed)
```

Causal Mediation Analysis

Quasi-Bayesian Confidence Intervals

	Estimate	95% CI Lower	95% CI Upper	p-value
ACME	0.276803	0.144987	0.43	<2e-16 ***
ADE	-0.115043	-0.316462	0.07	0.268
Total Effect	0.161760	0.000729	0.31	0.048 *
Prop. Mediated	1.653327	0.507091	9.66	0.048 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Sample Size Used: 100

Simulations: 1000

- By running this we get the Average Causal Mediation Effects (ACME), our Average Direct Effects (ADE), combined indirect and direct effects (Total Effect), and the ratio of these estimates (Prop. Mediated).
- The ACME is the indirect effect of M (Total Effects - ADE) and the associated p -value value tells us if our mediation effect is significant.
- We can bootstrap our data and fit a model based on our estimated population parameters (which is recommend over the default CI estimation method above)...

```
> fitMedBoot <- mediate(fitM, fitY, boot = TRUE, sims =
10000, treat = "X", mediator = "M")
> summary(fitMedBoot)
```

```
> summary(fitMedBoot)
```

Causal Mediation Analysis

Nonparametric Bootstrap Confidence Intervals with the Percentile Method

	Estimate	95% CI Lower	95% CI Upper	p-value
ACME	0.28078	0.14112	0.43	<2e-16 ***
ADE	-0.11179	-0.29548	0.09	0.273
Total Effect	0.16899	-0.00862	0.35	0.066 .
Prop. Mediated	1.66151	-3.92801	10.91	0.066 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Sample Size Used: 100

Simulations: 10000

- We now see that the ACME is the only one that is significant - this tells us we have a significant moderator - with no direct effect of our predictor or combined effect of predictor and moderator when the moderator is taken into consideration.