

```
> mixed_model <- lmer(rt ~ condition + (1 | subject) + (1 | item), data = fulldata)
> summary(mixed_model)
Linear mixed model fit by REML. t-tests use Satterthwaite's method ['lmerModLmerTest']
Formula: rt ~ condition + (1 | subject) + (1 | item)
Data: fulldata
```

REML criterion at convergence: 1276.5

Scaled residuals:

	Min	1Q	Median	3Q	Max
	-2.59882	-0.62360	0.07231	0.57203	2.91523

Random effects:

Groups	Name	Variance	Std.Dev.
subject	(Intercept)	7952.1	89.17
item	(Intercept)	436.3	20.89
Residual		20938.7	144.70

Number of obs: 100, groups: subject, 10; item, 5

Fixed effects:

	Estimate	Std. Error	df	t value	Pr(> t)
(Intercept)	1067.99	36.07	12.62	29.61	4.82e-13 ***
conditionsmall	187.83	28.94	85.00	6.49	5.46e-09 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Correlation of Fixed Effects:

	(Intr)
conditnsml1	-0.401

More
variability in
subjects than
in scenarios.

The intercept corresponds to the RT to the Large Condition - going from Large to Small contexts increases RT by around 188 ms.

- To determine whether our mixed effects model is significant, we need to know whether it differs from what we'd expect if Condition didn't influence Reaction Times.

```
mixed_model_null <- lmer(rt ~ (1 | subject) + (1 | item), data = fulldata)
```

- This model which we call `mixed_model_null` removes Condition as a predictor - in other words, it simply contains our random effects.