

```
> fitMedBoot <- mediate(fitM, fitY, boot = TRUE, sims =
10000, treat = "X", mediator = "M")
> summary(fitMedBoot)
```

```
> summary(fitMedBoot)
```

Causal Mediation Analysis

Nonparametric Bootstrap Confidence Intervals with the Percentile Method

	Estimate	95% CI Lower	95% CI Upper	p-value
ACME	0.28078	0.14112	0.43	<2e-16 ***
ADE	-0.11179	-0.29548	0.09	0.273
Total Effect	0.16899	-0.00862	0.35	0.066 .
Prop. Mediated	1.66151	-3.92801	10.91	0.066 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Sample Size Used: 100

Simulations: 10000

- We now see that the ACME is the only one that is significant - this tells us we have a significant moderator - with no direct effect of our predictor or combined effect of predictor and moderator when the moderator is taken into consideration.

Moderation Effects

- Moderation tests whether a variable (Z) affects the direction and/or strength of the relation between an IV (X) and a DV (Y).
- In other words, moderation tests for interactions that affect when relationships between variables occur.
- Imagine we are interested in whether the relationship between the number of hours of sleep (X) a student receives and the attention that they pay to this lecture (Y) is influenced by their consumption of coffee (Z).