

Border rank two does not force a tensor-square rank saving

Matthew J. Colbrook*

11 September 2026

Authorship is recorded at the submitter's request. AI assistance is disclosed. An independent agent reviewed this complete mathematical source against the original repository targets; a separate agent checked the accompanying computations. This is not external human peer review or formal certification, and no priority claim is made. The original proposed manuscripts have now passed independent review; reports and live eligibility checks accompany this attributed edition.

Abstract

An explicit smooth, irreducible, nondegenerate curve $X \subset \mathbb{P}^{11}$ and a point p satisfy $\underline{R}_X(p) = 2$, $R_X(p) = 3$, and $R_{X \times X}(p \otimes p) = 9$. Thus the implication in repository problem TR-27 is false for its stated class of projective varieties. More generally, for every $r \geq 3$ and every $m \geq 2$, a smooth projected rational normal curve has a point of border rank two and rank r whose first m tensor powers have ranks $r, r^2, \dots, r^m$. The proof is elementary: a projection preserves independence of small sets of distinct curve points while identifying a high-rank tangent vector with a sum of r curve points. All tensor products retain their separate factors.

1 The claim being answered

For a reduced irreducible nondegenerate projective variety $X \subset \mathbb{P}(V)$ over $\mathbb{C}$, let $R_X(p)$ be the smallest number of points of X whose span contains p . Border rank $\underline{R}_X(p)$ is defined by Zariski closure of these rank loci. Scalars in linear combinations are unrestricted. For $k \geq 1$, write $X^{\times k}$ for its Segre-embedded product in $\mathbb{P}(V^{\otimes k})$.

Problem TR-27 asks whether

$$\underline{R}_X(p) < R_X(p) \implies R_{X \times X}(p \otimes p) < R_X(p)^2 \quad (1)$$

always holds. The same implication is Conjecture 1.1 of Ballico–Bernardi–Gesmundo–Oneto–Ventura [2], and is restated as Conjecture 4.9 in [3]. The construction below disproves precisely this general-variety statement. It does not assert a counterexample when X is required to be a Segre or a Veronese variety.

Theorem 1. *Let $r \geq 3$ and $m \geq 2$ be integers, and put $D = r^m + r$. There are a smooth rational nondegenerate curve $X \subset \mathbb{P}^{D-1}$ and a point p such that*

$$\underline{R}_X(p) = 2, \quad R_X(p) = r, \quad R_{X^{\times k}}(p^{\otimes k}) = r^k \quad (1 \leq k \leq m).$$

In particular, $r = 3$, $m = 2$, and $D = 12$ give a counterexample to (1).

*Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge, United Kingdom. Email: m.colbrook@damtp.cam.ac.uk.

2 Two elementary independence lemmas

Let $E_D = \mathbb{C}^{D+1}$ with basis $e_0, \dots, e_D$. Write

$$c_D(t) = \sum_{j=0}^D t^j e_j \quad (t \in \mathbb{C}), \quad c_D(\infty) = e_D.$$

The projective points $[c_D(t)]$ form the rational normal curve C_D . Distinct points of C_D are linearly independent whenever their number is at most $D + 1$. This follows from the Vandermonde determinant, including the usual last-coordinate version when one point is at infinity.

Lemma 2. *For $D \geq 2$, $R_{C_D}([e_1]) = D$.*

Proof. Suppose e_1 were a linear combination of $\ell \leq D - 1$ distinct curve vectors, with parameter set $S \subset \mathbb{P}^1$. Define

$$f(T) = T \prod_{a \in S \setminus \{0, \infty\}} (T - a).$$

Then $f'(0) \neq 0$. Every finite parameter in S is a root of f . If $\infty \notin S$, then $\deg f \leq \ell + 1 \leq D$; if $\infty \in S$, then $\deg f \leq \ell \leq D - 1$. Write $f(T) = \sum_{j=0}^D f_j T^j$. The linear functional $L_f(e_j) = f_j$ annihilates every curve vector in the putative decomposition: for finite a , $L_f(c_D(a)) = f(a)$, while $L_f(c_D(\infty)) = f_D = 0$ when infinity occurs. But $L_f(e_1) = f_1 = f'(0) \neq 0$, a contradiction.

For the upper bound, summation over the D th roots of unity gives

$$e_1 = \frac{1}{D} \sum_{\zeta^D=1} \zeta^{-1} c_D(\zeta).$$

Indeed, for $0 \leq j \leq D$, the sum of ζ^{j-1} is nonzero only at $j = 1$. □

Lemma 3. *Suppose that every set of at most K distinct points of $X \subset \mathbb{P}(V)$ is linearly independent. If $R_X(p) = r$ and $r^k - 1 \leq K$, then*

$$R_{X^{\times k}}(p^{\otimes k}) = r^k.$$

Proof. An r -term expression for p gives the upper bound by expansion. Suppose, for a contradiction, that a representative v of p satisfies

$$v^{\otimes k} = \sum_{\ell=1}^L \lambda_\ell x_{1\ell} \otimes \cdots \otimes x_{k\ell}, \quad L \leq r^k - 1, \tag{2}$$

where $[x_{j\ell}] \in X$. For each factor j , collect the distinct projective points that occur in that factor and choose one representative for each. Denote this set of representatives by A_j . It has at most $L \leq K$ elements, so is linearly independent.

Contract (2) in all factors except j with any functional $\varphi \in V^*$ satisfying $\varphi(v) = 1$. It follows that $v \in \text{span } A_j$. In the unique expansion of v in the basis A_j , at least r coefficients are nonzero, since otherwise $R_X(p) < r$.

The products of the bases $A_1, \dots, A_k$ form a basis of their tensor-product span. In this basis, $v^{\otimes k}$ has at least r^k nonzero coordinates: its support is the Cartesian product of the nonzero supports of the k expansions of v . The right-hand side of (2) uses at most L such basis vectors, even after repeated products are combined. This is impossible for $L < r^k$. □

3 The projected curve

Choose pairwise distinct nonzero complex numbers $a_1, \dots, a_r$; the choices $a_i = i$ suffice. Set

$$b = \sum_{i=1}^r c_D(a_i), \quad z = e_1 - b, \quad V = E_D/\mathbb{C}z,$$

and let $\pi : E_D \rightarrow V$ be the quotient map. Lemma 2 and subadditivity of rank give

$$D = R_{C_D}([e_1]) \leq R_{C_D}([z]) + R_{C_D}([b]) \leq R_{C_D}([z]) + r. \quad (3)$$

Consequently $R_{C_D}([z]) \geq D - r = r^m$; in particular, $z \neq 0$ and $[z] \notin C_D$. The restriction of projective projection to C_D is therefore a morphism. Let X be its image, with its reduced structure. It is closed and irreducible because C_D is projective and irreducible. It is nondegenerate because C_D spans $\mathbb{P}(E_D)$ and π is surjective.

Proposition 4. *Every set of at most $K = r^m - 1$ distinct points of X is linearly independent.*

Proof. Choose lifts $u_1, \dots, u_s$ on C_D of $s \leq K$ distinct image points. The lifts are distinct. A linear dependence among their images would give

$$\sum_{i=1}^s \alpha_i u_i = \lambda z.$$

If $\lambda \neq 0$, this expresses z using at most $s \leq r^m - 1$ curve points, contrary to (3). If $\lambda = 0$, independence of at most $D + 1$ distinct rational-normal-curve points forces every $\alpha_i = 0$. Thus no nontrivial dependence exists. $\square$

Now let $p = [\pi(e_1)]$. Since $\pi(z) = 0$,

$$\pi(e_1) = \pi(b) = \sum_{i=1}^r \pi(c_D(a_i)). \quad (4)$$

The r image points in this sum are distinct and independent. Distinctness also follows directly from (3): identifying two curve points would put z in their span and give rank at most two. In particular, the sum in (4) is nonzero.

This expression has minimal length r . Otherwise a shorter expression and (4) would give two different expressions in the span of at most $2r - 1$ distinct points of X . Because $2r - 1 \leq r^m - 1 = K$, Proposition 4 makes that union independent. Uniqueness of basis coordinates would force all r original points to appear in the shorter expression, a contradiction. Thus $R_X(p) = r$.

For border rank, as $t \rightarrow 0$ we have

$$\frac{\pi(c_D(t)) - \pi(c_D(0))}{t} = \pi(e_1) + \sum_{j=2}^D t^{j-1} \pi(e_j) \longrightarrow \pi(e_1).$$

For all sufficiently small nonzero t this is a nonzero vector of X -rank at most two. Its projective limit is p , so $\underline{R}_X(p) \leq 2$. Since $p \notin X$ and X is closed, the border rank is not one. Therefore $\underline{R}_X(p) = 2$.

Finally, for $1 \leq k \leq m$ we have $r^k - 1 \leq K$. Applying Lemma 3 proves all rank equalities in Theorem 1.

3.1 The curve is smooth

For completeness, the construction is not relying on singularities of X . In fact,

$$\underline{R}_{C_D}([z]) = r + 2. \quad (5)$$

The upper bound is the sum of a tangent vector, of border rank two, and r curve vectors. To prove the lower bound, note that $D \geq 2r + 2$ and form the $(r + 2) \times (r + 2)$ Hankel matrix

$$H = (z_{i+j})_{0 \leq i, j \leq r+1}, \quad z_j = \delta_{j1} - \sum_{\nu=1}^r a_\nu^j.$$

Every curve vector maps to a matrix of rank at most one under this linear map, including the point at infinity. Hence a nonzero $(r + 2)$ -minor is a border-rank lower bound.

Put $u(a) = (1, a, \dots, a^{r+1})^\top$ and

$$B = [u(0) \ u'(0) \ u(a_1) \ \cdots \ u(a_r)], \quad M = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \oplus (-I_r).$$

Directly, $H = BMB^\top$. The square matrix B is invertible: a polynomial of degree at most $r + 1$ in the kernel of $B^\top$ would have a double zero at zero and zeros at all a_i , and hence would be divisible by a polynomial of degree $r + 2$. It must be zero. Thus H is invertible, proving (5).

In particular, the projection center lies on neither a secant nor a tangent line of C_D . The projection is injective on C_D and its differential is injective at every point: a failure of either assertion would place the center on the corresponding secant or tangent line. A proper injective immersion of this smooth complex projective curve is an embedding. Thus X is a smooth rational curve, as claimed.

4 An explicit counterexample in twelve coordinates

Take $r = 3$, $m = 2$, $D = 12$, and $a_1 = 1, a_2 = 2, a_3 = 3$. For $0 \leq j \leq 12$, set $S_j = 1 + 2^j + 3^j$, and use the coordinate index set

$$J = \{0, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}.$$

Here $z_1 = 1 - S_1 = -5$. A quotient map with kernel $\mathbb{C}z$ is

$$P : E_{12} \longrightarrow \mathbb{C}^{12}, \quad (P(v))_j = 5v_j - S_j v_1 \quad (j \in J).$$

It has rank twelve since its columns with indices in J form $5I_{12}$, and $P(z) = 0$. The curve is the image of the following explicit homogeneous parametrization:

$$[s : t] \longmapsto \left[\left(5s^{12-j}t^j - S_j s^{11}t \right)_{j \in J} \right]. \quad (6)$$

The common-factor-free, base-point-free property also follows from the quotient construction. Directly, at $s = 0$ the last coordinate is $5t^{12} \neq 0$; on the chart $s = 1$, vanishing of the first coordinate forces $t = 5/3$, at which the $j = 2$ coordinate is $-85/9 \neq 0$.

On this affine chart write $g(t) = (5t^j - S_j t)_{j \in J}$. A representative of the required point is

$$\begin{aligned} v = P(e_1) &= (-S_j)_{j \in J} \\ &= (-3, -14, -36, -98, -276, -794, -2316, -6818, \\ &\quad -20196, -60074, -179196, -535538). \end{aligned}$$

The identities

$$g'(0) = v, \quad g(1) + g(2) + g(3) = v$$

exhibit, respectively, its border-two degeneration and its three-term expression. The preceding proofs give

$$\boxed{\mathbf{R}_X([v]) = 2, \quad \mathbf{R}_X([v]) = 3, \quad \mathbf{R}_{X \times X}([v \otimes v]) = 9.}$$

An explicit nine-term decomposition is

$$v \otimes v = \sum_{a=1}^3 \sum_{b=1}^3 g(a) \otimes g(b).$$

The lower bound applies to arbitrary complex decompositions on the whole curve, not only to decompositions using these three parameter values.

The 5×5 center Hankel determinant, using coordinates $z_0, \dots, z_8$, equals 5184. This provides an additional exact check that the center has border rank at least five and that the example is a smooth projection.

5 Why there is no conflict with eventual strict submultiplicativity

The known eventual-power statement in [2, Section 1] concerns sufficiently large powers, not the square. There is an elementary eventual-saving bound for the present examples. Put

$$q(t) = \pi(c_D(t) - c_D(0)) = tv + t^2\pi(e_2) + \dots + t^D\pi(e_D).$$

For an integer $k \geq 1$, set $L = k(D - 1) + 1$. Extracting the coefficient of t^k by a roots-of-unity sum gives

$$v^{\otimes k} = \frac{1}{L} \sum_{\zeta^L=1} \zeta^{-k} q(\zeta)^{\otimes k}.$$

The degrees of $q(t)^{\otimes k}$ lie between k and kD , so no other exponent is congruent to k modulo L . Each $q(\zeta)$ is a difference of two curve vectors. Therefore

$$\mathbf{R}_{X \times k}(p^{\otimes k}) \leq (k(D - 1) + 1)2^k.$$

For fixed D and $r > 2$, this is less than r^k for all sufficiently large k . For the explicit $D = 12$ example, it already proves a saving at $k = 13$, since $144 \cdot 2^{13} = 1,179,648 < 3^{13} = 1,594,323$. Thus the examples delay a saving; they do not prevent one at every power.

6 Verification and scope

The exact-arithmetic script `verify.py` checks the coordinates, quotient rank and kernel, Hankel determinant, nine-term expression, and interpolation identities. It does not replace the universal independence and rank lower-bound proofs.

This counterexample answers the universal implication in TR-27. Arbitrary finite delay strengthens the same result; it is not a second problem claimed solved. The Segre and Veronese cases are not settled here.

References

- [1] OpenProblemsInNLA, “TR-27—Border-rank deficiency forcing strict submultiplicativity at the tensor square,” problem statement, checked September 11, 2026. <https://github.com/ajt60gaibb/OpenProblemsInNLA/tree/main/tensor-computations/TR-27>.
- [2] E. Ballico, A. Bernardi, F. Gesmundo, A. Oneto, and E. Ventura, “Geometric conditions for strict submultiplicativity of rank and border rank,” *Annali di Matematica Pura ed Applicata* **200** (2021), 187–210. Conjecture 1.1. <https://arxiv.org/abs/1909.03811>.
- [3] A. Oneto and E. Ventura, “Ranks of tensors: geometry and applications,” *Bollettino dell’Unione Matematica Italiana* **18** (2025), 751–778. Conjecture 4.9. <https://doi.org/10.1007/s40574-025-00472-9>.