

Nonnegative and positive factorizations

Exact nonnegative rank of regular polygon slack matrices

Difficulty: challenging **Importance:** interesting to the community

Status: open; checked 2026-09-08

Area: structured nonnegative matrix factorization

Context and notation

All factorizations are over the real numbers. For $X \in \mathbb{R}_{\geq 0}^{m \times n}$, define

$$\text{rank}_+(X) = \min\{r \geq 0 : X = WH, \quad W \in \mathbb{R}_{\geq 0}^{m \times r}, H \in \mathbb{R}_{\geq 0}^{r \times n}\}.$$

Problem statement

For each integer $n \geq 3$, define the $n \times n$ nonnegative matrix

$$S_n(i, j) = \cos(\pi/n) - \cos((2i + 1 - 2j)\pi/n), \quad 0 \leq i, j < n.$$

It is the slack matrix obtained from the vertices of the regular n -gon on the unit circle and its supporting facet inequalities.

Question

With $k = \lceil \log_2 n \rceil$, is it true for every $n \geq 3$ that

$$\text{rank}_+(S_n) = \begin{cases} 2k - 1, & 2^{k-1} < n \leq 2^{k-1} + 2^{k-2}, \\ 2k, & 2^{k-1} + 2^{k-2} < n \leq 2^k? \end{cases}$$

References

Arnaud Vandaele, Nicolas Gillis, François Glineur, and Daniel Tuytens, *Heuristics for Exact Nonnegative Matrix Factorization*, Journal of Global Optimization **65** (2016), 369–400, §6.2, Conjecture 2. Vandaele, Gillis, and Glineur, *On the Linear Extension Complexity of Regular n -gons*, Linear Algebra and its Applications **521** (2017), 217–239, proves the corresponding upper bound. Nicolas Gillis, *Nonnegative Matrix Factorization*, SIAM, 2020, §3.6.3.4, p. 90.

Status evidence

Baeckelant, Vandaele, and Gillis, *Computing Lower Bounds on the Nonnegative Rank via Non-Convex Optimization Solvers*, revised July 6, 2026, Appendix A.2, still identifies sharpness as conjectural. Searches for the conjecture and regular-polygon nonnegative ranks in 2025–2026 found no subsequent resolution. The family is counted once.