

Linear systems and elimination

The exact four-step amplification of restarted Anderson acceleration

Difficulty: challenging **Importance:** interesting to the community

Status: Solved

Independently reviewed resolution - 2026-09-11

Negative resolution. Section 2 refutes the exact four-step identity using $M = \text{diag}(1/10, 1/2, 3/5)$ and $v = (1, 1, 1)^T$: the squared norm ratio is $1920682/21289638243 > 1/14641$, the square of the proposed factor. Both M and $I - M$ are positive definite. Section 3 proves unbounded underestimation over a parameter family. The separate asymptotic convergence question is not resolved.

Author: Matthew J. Colbrook, Department of Applied Mathematics and Theoretical Physics, University of Cambridge. [Complete manuscript](#), [independent proof review](#), and [submission record](#). The supplied notes were reconstructed with substantial AI assistance. This is independent agent verification, not external human peer review or formal proof-assistant certification; no novelty or priority claim is made.

The difficulty, importance and rating rationale below are historical assessments of the original open target. Original statements, references and dated audits are preserved.

Rating rationale: Challenging reflects a global maximization of a nonlinear homogeneous residual map in arbitrary dimension; community impact is an exact convergence factor for restarted Anderson acceleration.

Let $n \geq 2$ and let $M \in \mathbb{R}^{n \times n}$ be nonzero and symmetric, with $1 \notin \sigma(M)$. Put $A = I - M$. Define the positively homogeneous map

$$R(v) = M \left(v - \frac{v^T A v}{\|A v\|_2^2} A v \right) \quad (v \neq 0), \quad R(0) = 0.$$

Two steps of restarted Anderson acceleration with memory one applied to $x = Mx + b$ propagate the residual by R . If $m_1, \dots, m_n$ are the eigenvalues of M , is

$$\max_{v \neq 0} \frac{\|R(R(v))\|_2}{\|v\|_2} = \max_{i \neq j} \left(\frac{m_i m_j (m_j - m_i)}{|m_i(m_i - 1)| + |m_j(m_j - 1)|} \right)^2 ?$$

A term with zero denominator is defined as zero: under the assumptions this can occur only when $m_i = m_j = 0$.

The conjecture says that a largest four-step residual amplification is attained using only two orthogonal eigenvectors, irrespective of the dimension. It supplies an exact worst-case factor for this restart scheme, relevant to accelerated stationary solvers and multigrid.

References

O. A. Krzysik, H. De Sterck, and A. Smith, *Asymptotic convergence of restarted Anderson acceleration for certain normal linear systems*, SISC 47(2025), Conjecture 10, (9),(16),(24),(27) ([journal](#); [arXiv v4](#)). Conjecture 10 states the maximum through two-eigenvector nonlinear eigenvalues; equation(24) evaluates their scalar maximum. The map formulation above also covers exact termination without an undefined $\alpha(0)$.

Status check — 2026-09-08

The arXiv history lists v4,12 May 2025, as latest. Its Conjecture 10 and journal abstract retain the conditional result. Searches for the paper title, “restarted Anderson Conjecture 10”, and later proof/counterexample results found no resolution. Windowed AA, GMRES(1), and restarted CG use different residual maps. The September 2026 Forsythe paper resolves the CG restart question and is not a solution claim for the map above.

Audit update — 2026-09-10

Rechecked [version 4](#), Conjecture 10 and its surrounding discussion, against the [2025 journal record](#). The general-dimensional identity is still conjectural; the order-two case alone does not justify a substantive partial-status label. Targeted searches for this four-step amplification conjecture found no later resolution.