

Eigenvalues and inverse problems

The smallest-multiplier rule for nearest unit-absolute-determinant matrices

Difficulty: challenging **Importance:** interesting to the community

Status: explicit open root-selection question; no resolution located

For $n \geq 2$, define

$$\mathcal{G}_n = \{X \in \mathbb{R}^{n \times n} : \det X \in \{-1, 1\}\}, \quad \|Y\|_F^2 = \sum_{i,j} y_{ij}^2.$$

For a real data matrix U , consider the real stationary pairs

$$\mathcal{C}(U) = \{(X, c) \in \mathcal{G}_n \times \mathbb{R} : X^T(U - X) = cI_n\}.$$

For generic U , this set is finite. Choose a pair (X_*, c_*) whose multiplier has minimum absolute value. Does

$$\|U - X_*\|_F = \min_{X \in \mathcal{G}_n} \|U - X\|_F$$

hold for every $n \geq 2$ and generic real U ?

Here generic means outside an exceptional proper real algebraic set; in particular U is invertible and its squared singular values are distinct. If minimum absolute multipliers tie, the question can be restricted to the generic locus where the choice is unique. The source phrases the same selection question using the real roots of its elimination polynomial $R_1(c)$, which correspond one-to-one to stationary matrices for generic data.

Relevance

This is a concrete question about selecting the global nearest matrix from the stationary solutions of a structured matrix nearness problem. A positive answer would supply a scalar criterion for identifying the global solution. Both determinant signs are allowed; this must not be silently replaced by projection onto $\mathrm{SL}_n(\mathbb{R})$ alone.

References

- J. A. Baaijens and J. Draisma, *Euclidean distance degrees of real algebraic groups*, Linear Algebra and its Applications 467 (2015), 174–187, §4.1, equations (4)–(5), elimination algorithm, and Problem 4.3 on p.186 ([version of record](#); [journal](#)).
- P. Jaap and O. Sander, *How to project onto $SL(n)$* , arXiv:2501.19310 (2025), introduction pp.1–2, discussing the Baaijens–Draisma conjectural global selection rule ([primary manuscript](#)); Journal of Optimization Theory and Applications 209 (2026), article 40 ([journal](#)).

Status check — 2026-09-08

The January 2025 manuscript explicitly describes the earlier proposed global-minimum selection as conjectural. Its journal version appeared in April 2026; the accessible journal abstract and arXiv version history were checked, but the subscription-only journal body was not inspected. Searches for “nearest determinant-one matrix”, “Baaijens”, “smallest absolute value”, “projection $\mathrm{SL}(n)$ ”, and 2025–2026 found no resolution of this exact rule. The problem retains the source’s generic-data scope.