

Matrix inequalities and norms

The triangle conjecture for the maximal symmetric modulus

Difficulty: challenging **Importance:** interesting to specialist

Problem statement

For a complex square matrix X , let $|X| = (X^*X)^{1/2}$ and define its maximal symmetric modulus by the finite-dimensional limit

$$M(X) = \lim_{r \rightarrow \infty} (|X|^r + |X^*|^r)^{1/r}, \quad r \in \mathbb{N}.$$

This is the supremum of $|X|$ and $|X^*|$ in Olson's spectral order. For every $n \geq 1$ and every $A, B \in \mathbb{C}^{n \times n}$, must there exist unitary U, V of order n satisfying

$$M(A + B) \leq UM(A)U^* + VM(B)V^*?$$

The order in this displayed inequality is ordinary positive-semidefinite order.

Why it matters

The modulus combines both positive polar factors through spectral order. A sharp triangle inequality would give a new way to control positive representatives of nonnormal matrix sums.

References

1. J.-C. Bourin and E.-Y. Lee, *Averages over matrix unitary orbits and spectral order*, arXiv:2606.15624v2 (18 June 2026), §§3.1–3.3, Question 3.11 and Conjecture 3.12. [Primary text](#).
2. J.-C. Bourin and E.-Y. Lee, *Some hybrid matrix triangle inequalities*, arXiv:2606.29188v1 (28 June 2026), §3, Lemma 3.3 and Theorem 3.4. [Primary text](#).

Status check — 2026-09-08

Latest versions v2 and v1 were checked. The follow-up proves inequalities with the ordinary modulus of the total sum on the left, while the present conjecture has its maximal symmetric modulus there. Its conclusion therefore does not establish the displayed assertion. Searches included *maximal symmetric modulus conjecture*, *Bourin Lee 2606.15624 conjecture*, and *Some hybrid matrix triangle inequalities*. No later resolution was located.