

Matrix functions and stability

The uniform expansion threshold for Toeplitz symbols with higher-order zeros

Difficulty: challenging **Importance:** interesting to specialist**Status:** Open

Rating rationale: Matching bulk asymptotics with extreme eigenvalues at the exact breakdown order is challenging; the detailed expansion threshold is mainly important to structured spectral specialists.

Statement

For each integer $m \geq 3$, let

$$g_m(\theta) = (2 \sin(\theta/2))^{2m}, \quad \hat{g}_{m,k} = \frac{1}{2\pi} \int_{-\pi}^{\pi} g_m(\theta) e^{-ik\theta} d\theta.$$

Let $T_n(g_m) = (\hat{g}_{m,j-k})_{j,k=1}^n$, with eigenvalues $\lambda_{n,1} \leq \dots \leq \lambda_{n,n}$ counted with multiplicity. For real continuous functions $d_0, \dots, d_{2m}$ on $[0, \pi]$, independent of n, j , write

$$R_{p,n,j} = \lambda_{n,j} - \sum_{k=0}^p \frac{d_k(j\pi/(n+2))}{(n+2)^k}.$$

Conjecture 8.4 of Barrera–Böttcher–Grudsky–Maximenko. Such coefficient functions exist, with $d_0 = g_m$, for which all three assertions hold:

1. For every integer $0 \leq p \leq 2m - 1$, there are constants $D_p > 0$ and $N_p \in \mathbb{N}$ such that

$$|R_{p,n,j}| \leq D_p (n+2)^{-p-1} \quad (n \geq N_p, 1 \leq j \leq n).$$

2. There are constants $D_{2m} > 0$ and $N_{2m} \in \mathbb{N}$ such that

$$|R_{2m,n,j}| \leq D_{2m} (n+2)^{-2m-1}$$

whenever $n \geq N_{2m}$ and $\lceil (\log(n+2))^2 \rceil \leq j \leq n$.

3. No constants $D > 0, N \in \mathbb{N}$ make the bound $|R_{2m,n,j}| \leq D(n+2)^{-2m-1}$ valid for every $n \geq N$ and every $1 \leq j \leq n$.

All constants may depend on m and the expansion order; $\log$ denotes the natural logarithm. The same coefficient functions are used throughout. The continuity and existence quantifiers make explicit the regular-expansion convention of the source's Theorem 1.2 and §8; these are not matrix-size-dependent fitted coefficients.

Numerical significance

Regular eigenvalue expansions support accurate computation without forming large Toeplitz matrices. This conjecture identifies exactly where extreme eigenvalues obstruct a uniform expansion for these polynomial symbols. It retains all three parts as one problem.

References and status check

- M. Barrera, A. Böttcher, S. M. Grudsky and E. A. Maximenko, *Eigenvalues of even very nice Toeplitz matrices can be unexpectedly erratic*, Oper. Theory Adv. Appl. **268** (2018), 51–77, DOI; [author preprint](#), Conjecture 8.4, p. 26, equation (8.4); Theorem 1.2 supplies the proved $m = 2$ analogue.
- M. Barrera, S. Grudsky, V. Stukopin and I. Voronin, *Asymptotics of the eigenvalues of seven-diagonal Toeplitz matrices of a special form*, Adv. Oper. Theory **9** (2024), 79, DOI; [preprint](#), Theorems 2.3–2.6. This studies the sixth-order-zero case with more elaborate formulas.
- A. Böttcher, *Ten years with Sergei Grudsky in the eigenvalue bulk of Toeplitz matrices*, J. Math. Sci. **298** (2026), 363–376, DOI, section “Beyond the simple-loop class,” Theorems 2–5 and discussion of higher-order zeros.

On 2026-09-10, checked the original full preprint, the seven-diagonal follow-up’s full preprint and publication record, the 2026 survey, and targeted title/conjecture/2025–2026 searches. No resolution of all parts for every $m \geq 3$ was located. Later local second-order expansions and results confined to $m = 3$ do not establish this full statement. This bounded search is not a proof of openness.

Audit — 2026-09-10

Independently rechecked [Conjecture 8.4](#), [the seven-diagonal paper’s Theorems 2.3–2.6](#), and [Böttcher’s survey](#). Corrected the status to Open: the proved $m = 2$ analogue lies outside the displayed range, and the $m = 3$ local and second-order formulas do not establish the three-part threshold assertion. The [March 2026 eigenvalue-superposition paper](#) assumes simple-loop symbols, excluding these higher-order zeros. Title and conjecture searches found no complete resolution. Challenging difficulty and specialist importance are retained.