

Tensor computations

Degrees of the two Rayleigh–Ritz discriminant parts for rational normal curves

Difficulty: challenging **Importance:** interesting to specialist**Status:** Partially resolved

Rating rationale: Separating the isotropic and nonisotropic degree contributions requires substantial algebraic analysis beyond the known total. The result mainly interests specialists in conditioning of constrained eigenvalue problems.

Problem statement

For every integer $d \geq 2$, use the specific rational normal curve

$$C_d = \{[a^d : a^{d-1}b : \dots : b^d] : [a : b] \in \mathbb{P}^1\} \subset \mathbb{P}^d.$$

For a complex symmetric matrix H of order $d+1$, consider $\rho_H(z) = z^\top H z / (z^\top z)$ on the part of C_d where $z^\top z \neq 0$. Transposes here do not conjugate.

Let $\mathcal{R}$ be the Zariski closure in $C_d \times \mathbb{P}(\text{Sym}_{d+1}(\mathbb{C}))$ of all pairs $([z], [H])$ for which $[z]$ is a critical point of $\rho_H|_{C_d}$. Choose homogeneous generators g_i of its ideal and let

$$\mathcal{B} = \{([z], [H]) \in \mathcal{R} : \text{rank}(\partial g_i / \partial z_j) \leq d-1\}.$$

Put $Q = \{([z], [H]) : z^\top z = 0\}$ and let π project to the matrix factor. Define

$$\Sigma_{\text{iso}} = \pi(\mathcal{B} \cap Q), \quad \Sigma_{\text{noniso}} = \overline{\pi(\mathcal{B} \setminus Q)}^{\text{Zar}}.$$

Are these projective hypersurfaces with

$$\deg \Sigma_{\text{noniso}} = 6(d-1), \quad \deg \Sigma_{\text{iso}} = 2d?$$

Degree refers to each reduced algebraic set, including all its components. The question is Conjecture 3.14 in the source. Both parts belong to one problem; the standard unweighted embedding and the bilinear quadratic form are fixed.

Why it matters

These discriminants locate parameter values at which constrained Rayleigh-quotient critical points collide or cease to form a reduced finite set. They describe degeneracy in a basic nonlinear analogue of matrix eigenvalue computations.

References and status check

1. V. Borovik, H. Friedman, S. Hoşten, and M. Pfeffer, *Numerical Algebraic Geometry for Energy Computations on Tensor Train Varieties*, [arXiv:2512.06939v2](https://arxiv.org/abs/2512.06939v2), §3.2, Conjecture 3.14, pp.13–14; the correspondence and ramification definitions are on pp.7–13.
2. The same paper, Proposition 3.10, proves the total discriminant degree $2(4d - 3)$; Examples 3.12–3.13 compute small cases.

Status check — 2026-09-10

Checked the June 2026 revision, Proposition 3.10, Examples 3.12–3.13 and Conjecture 3.14, and searched the title and rational-normal-curve Rayleigh–Ritz discriminants. Example 3.12 gives an explicit symbolic factorization for $d = 2$, with isotropic degree four and nonisotropic degree six; this is a proved small-degree case supporting Partially resolved. Example 3.13 also reports symbolic degree computation for $d = 3$. The separate numerical checks through $d = 10$ are evidence, not a general proof. The total-degree theorem alone does not determine the component split. No later general proof or counterexample was located.