
Original source distribution among included 02D records
n = 599 records
cattle feces
bovine manure
cattle manure
cattle fecal samples
beef
bovine feces
bovine fecal
feces (Bos taurus)
animal−cattle−steer
Rectal fecal grab samples from a commercial feedlot
cattle manure with soil
cattle
fecal beef calf
Cattle feces
calf feces
Rectal fecal grab samples from finishing cattle in a commercial feedlot
animal−cattle−heifer
Animal−Calf−Formula−fed Veal
Animal−Calf−Bob Veal
n=140 (23.37%)
n=116 (19.37%)
n=108 (18.03%)
n=87 (14.52%)
n=70 (11.69%)
n=28 (4.67%)
n=17 (2.84%)
n=8 (1.34%)
n=5 (0.83%)
n=4 (0.67%)
n=3 (0.5%)
n=3 (0.5%)
n=2 (0.33%)
n=2 (0.33%)
n=2 (0.33%)
n=1 (0.17%)
n=1 (0.17%)
n=1 (0.17%)
n=1 (0.17%)
Original source
0
40
80
120
160
Number of records


Original sources within each included 02D cohort/matrix g
n = 599 records
bovine manure [Unspecified_Manure_Environment_Preharvest | Manure/environment]
cattle feces [Unspecified_Feces_Rectal_Preharvest | Feces/rectal]
cattle manure [Unspecified_Manure_Environment_Preharvest | Manure/environment]
cattle fecal samples [Unspecified_Feces_Rectal_Preharvest | Feces/rectal]
beef [Beef_Preharvest | Other/unknown]
cattle feces [Beef_Preharvest | Feces/rectal]
bovine feces [Unspecified_Feces_Rectal_Preharvest | Feces/rectal]
bovine fecal [Unspecified_Feces_Rectal_Preharvest | Feces/rectal]
feces (Bos taurus) [Unspecified_Feces_Rectal_Preharvest | Feces/rectal]
animal−cattle−steer [Beef_Preharvest | Bovine/cattle non−fecal]
Rectal fecal grab samples from a commercial feedlot [Beef_Preharvest | Feces/rectal]
cattle manure with soil [Unspecified_Manure_Environment_Preharvest | Manure/environment]
cattle [Beef_Preharvest | Bovine/cattle non−fecal]
fecal beef calf [Beef_Preharvest | Feces/rectal]
Cattle feces [Beef_Preharvest | Feces/rectal]
calf feces [Unspecified_Feces_Rectal_Preharvest | Feces/rectal]
Rectal fecal grab samples from finishing cattle in a commercial feedlot [Beef_Preharvest | Feces/rectal]
animal−cattle−heifer [Beef_Preharvest | Bovine/cattle non−fecal]
Animal−Calf−Formula−fed Veal [Beef_Preharvest | Calf−associated non−fecal]
Animal−Calf−Bob Veal [Beef_Preharvest | Calf−associated non−fecal]
n=116 (51.1%)
n=110 (43.65%)
n=108 (47.58%)
n=87 (34.52%)
n=70 (100%)
n=30 (76.92%)
n=28 (11.11%)
n=17 (6.75%)
n=8 (3.17%)
n=5 (55.56%)
n=4 (10.26%)
n=3 (1.32%)
n=3 (33.33%)
n=2 (5.13%)
n=2 (5.13%)
n=2 (0.79%)
n=1 (2.56%)
n=1 (11.11%)
n=1 (50%)
n=1 (50%)
Original source [cohort | matrix class]
0
50
100
Number of records


Original source distribution by included 02D cohort/matrix group
n = 599 records
Beef_Preharvest | Bovine/cattle non−fecal
animal−cattle−steer
n=5
cattle
n=3
animal−cattle−heifer
n=1
Beef_Preharvest | Calf−associated non−fecal
Animal−Calf−Formula−fed Veal
n=1
Animal−Calf−Bob Veal
n=1
Beef_Preharvest | Feces/rectal
cattle feces
n=30
Rectal fecal grab samples from a commercial feedlot
n=4
fecal beef calf
n=2
Cattle feces
n=2
Rectal fecal grab samples from finishing cattle in a commercial feedlot
n=1
Beef_Preharvest | Other/unknown
beef
n=70
Unspecified_Feces_Rectal_Preharvest | Feces/rectal
cattle fecal samples
n=87
cattle feces
n=110
bovine feces
n=28
bovine fecal
n=17
feces (Bos taurus)
n=8
calf feces
n=2
Unspecified_Manure_Environment_Preharvest | Manure/environment
bovine manure
n=116
cattle manure
n=108
cattle manure with soil
n=3
Original source
0
50
100
Number of records

O157:H7 Focus: Beef_Preharvest

O157:H7 vs Non−O157:H7 vs No Call



n = 120 genomes

O157:H7 Focus: Unspecified_Feces_Rectal_Preharvest

O157:H7 vs Non−O157:H7 vs No Call



n = 252 genomes

O157:H7 Focus: Unspecified_Manure_Environment_Preharvest

O157:H7 vs Non−O157:H7 vs No Call



n = 227 genomes