Original sources within each included 02D cohort/matrix group

n = 599 records

Original source [cohort | matrix class]

bovine manure [Unspecified_Manure_Environment_Preharvest | Manure/environment] — n=116 (51.1%)
cattle feces [Unspecified_Feces_Rectal_Preharvest | Feces/rectal] — n=110 (43.65%)
cattle manure [Unspecified_Manure_Environment_Preharvest | Manure/environment] — n=108 (47.58%)
cattle fecal samples [Unspecified_Feces_Rectal_Preharvest | Feces/rectal] — n=87 (34.52%)
beef [Beef_Preharvest | Other/unknown] — n=70 (100%)
cattle feces [Beef_Preharvest | Feces/rectal] — n=30 (76.92%)
bovine feces [Unspecified_Feces_Rectal_Preharvest | Feces/rectal] — n=28 (11.11%)
bovine fecal [Unspecified_Feces_Rectal_Preharvest | Feces/rectal] — n=17 (6.75%)
feces (Bos taurus) [Unspecified_Feces_Rectal_Preharvest | Feces/rectal] — n=8 (3.17%)
animal−cattle−steer [Beef_Preharvest | Bovine/cattle non−fecal] — n=5 (55.56%)
Rectal fecal grab samples from a commercial feedlot [Beef_Preharvest | Feces/rectal] — n=4 (10.26%)
cattle manure with soil [Unspecified_Manure_Environment_Preharvest | Manure/environment] — n=3 (1.32%)
cattle [Beef_Preharvest | Bovine/cattle non−fecal] — n=3 (33.33%)
fecal beef calf [Beef_Preharvest | Feces/rectal] — n=2 (5.13%)
Cattle feces [Beef_Preharvest | Feces/rectal] — n=2 (5.13%)
calf feces [Unspecified_Feces_Rectal_Preharvest | Feces/rectal] — n=2 (0.79%)
Rectal fecal grab samples from finishing cattle in a commercial feedlot [Beef_Preharvest | Feces/rectal] — n=1 (2.56%)
animal−cattle−heifer [Beef_Preharvest | Bovine/cattle non−fecal] — n=1 (11.11%)
Animal−Calf−Formula−fed Veal [Beef_Preharvest | Calf−associated non−fecal] — n=1 (50%)
Animal−Calf−Bob Veal [Beef_Preharvest | Calf−associated non−fecal] — n=1 (50%)

0
50
100

Number of records