



A Computational Model for Prediction of Cardiac Arrest Outcome Using a Large Multi-Center Database

Blue Team Members:

Han Kim
Qingchu Jin
Tatiana Gelaf Romer
Hieu “Hugh” Nguyen
Sharmila Tamby

Biomedical PIs:

Dr. Robert Stevens
Dr. Jose Suarez
Dr. Christian Storm


Engineering PIs:

Dr. Raimond Winslow
Dr. Joseph Greenstein


Teaching Assistant:

Ran Liu


Problem Introduction



Importance and Significance




eICU Database




Aims


Aim 1: Data preprocessing and data exploration



Aim 2: Prediction of cardiac arrest recovery outcome



Aim 3: Early prediction of favorable recovery outcome if TTM is administered



Feature Space & Label Space

Feature Space

Time-Series (first 48 hours)

Blood
Pressure

SPO2

Breathing
Rate

Heart Rate

Temperature

Numerical

pH

Age

GCS at
admission

Categorical

Cardiac Arrest
Location


Intubation

Initial Rhythm

Sex

Label Space



GCS Score at Discharge



Label 1

Label 2

Aim 1: Data preprocessing




Aim 1: Learning Methods

Unsupervised learning


- K-means clustering
- Spectral clustering
- Principle component analysis
- ...

Supervised learning

- Generalized linear model
- Random forest
- Lasso regression
- ...



Aim 2: Prediction of Cardiac Arrest Recovery Outcome



Aim 2: Approaches

Supervised learning techniques

GLM

K-nearest neighbors


Random Forest

Boosting Algorithms

Support Vector
Machines (SVMs)

Possible improvements:

1. Try multiclass models



2. Try neural networks


ie. Recurrent Neural network

Gated Recurrent Unit, Long short-term memory

Aim 3: Targeted Temperature Management Efficacy Prediction

Predict which patient will have increased recovery if TTM is administered

- 6-8 hours after admission
- Temperature below 36°C for at least 12 hours.




Aim 3 Approaches

1. Locate TTM patients
2. Isolate features from first 6-8 hours
3. Use similar framework from Aims 1 & 2

Identify Patients who will either:

1. Respond favorably to TTM
2. Will not benefit from TTM




Acknowledgements



Dr. Robert Stevens



Dr. Jose Suarez



Dr. Christian Storm



Dr. Raimond Winslow



Dr. Sridevi Sarma



Dr. Joseph Greenstein



Ran Liu

Thank you for listening!

Questions? Comments? Suggestions?

APPENDIX

Targeted Temperature Management

- Targeted Temperature Management (formerly Therapeutic Hypothermia) is the intentional cooling of a patient to 33-36°C for 18-24 hours post-cardiac arrest.
- Multiple studies have shown induced therapeutic hypothermia improves neurologic function in post-cardiac arrest patients when initiated within six hours of the arrest; starting with two landmark studies in 2002.
- The American Heart Association in 2005 listed induced hypothermia as one of the guidelines for post-cardiac arrest care.



How Does It Work?



- Initial neurologic injury occurs when circulatory collapse impairs oxygen flow to the brain. Without oxygen, the brain switches to anaerobic metabolism, resulting in excessive calcium and glutamate excretion. This makes brain cells more excitable, leading to further hypoxemia, in turn causing mitochondrial and cellular death.
- Cellular death results in cerebral edema, producing further damage.
- The initial injury also disrupts the blood-brain barrier, which increases fluid in the brain and worsens cerebral edema.
- Hypothermia counteracts neuro-excitation in brain cells by stabilizing calcium and glutamate release, reducing the degree of cell death. It also stabilizes the blood-brain barrier and suppresses the inflammatory process, reducing cerebral edema. Cerebral metabolism decreases 6% to 10% for every degree Celsius that body temperature drops. As cerebral metabolism declines, the brain needs less oxygen.
- Hypothermia counteracts many of the destructive mechanisms of cardiac arrest. Similarly, hypothermia halts destructive brain mechanisms and lets the brain reset itself to normal functioning.
- Decard & Ebright (2011).

<https://cdn.ymaws.com/npofoklahoma.com/resource/collection/C3C77701-15EF-4C38-86ED-993BCB850168/Hypothermia%20in%20Cardiac%20Arrest.pdf>

How Does It Work?




- Initial neurologic injury occurs when circulatory collapse impairs oxygen flow to the brain. Without oxygen, the brain switches to anaerobic metabolism, resulting in excessive calcium and glutamate excretion. This makes brain cells more excitable, leading to further hypoxemia, in turn causing mitochondrial and cellular death.
- Cellular death results in cerebral edema, producing further damage.
- The initial injury also disrupts the blood-brain barrier, which increases fluid in the brain and worsens cerebral edema.
- Hypothermia counteracts neuro-excitation in brain cells by stabilizing calcium and glutamate release, reducing the degree of cell death. It also stabilizes the blood-brain barrier and suppresses the inflammatory process, reducing cerebral edema. Cerebral metabolism decreases 6% to 10% for every degree Celsius that body temperature drops. As cerebral metabolism declines, the brain needs less oxygen.
- Hypothermia counteracts many of the destructive mechanisms of cardiac arrest. Similarly, hypothermia halts destructive brain mechanisms and lets the brain reset itself to normal functioning.
- Decard & Ebright (2011).

<https://cdn.ymaws.com/npofoklahoma.com/resource/collection/C3C77701-15EF-4C38-86ED-993BCB850168/Hypothermia%20in%20Cardiac%20Arrest.pdf>


Targeted Temperature Management Review

- After cardiac arrest and successful resuscitation, patient can be determined if he/she is a candidate for TTM
 - Several Inclusion & Exclusion Criteria
- Multiple Cooling Methods
 - Ice Packs
 - Cooled Saline
 - Gaymar Unit
 - Zoll Catheter
- Cooled for 24 hours
- Slowly rewarmed
- Neurological Function assessed



Targeted Temperature Management Review

- After cardiac arrest and successful resuscitation, patient can be determined if he/she is a candidate for TTM
 - Several Inclusion & Exclusion Criteria
- Multiple Cooling Methods
 - Ice Packs
 - Cooled Saline
 - Gaymar Unit
 - Zoll Catheter
- Cooled for 24 hours
- Slowly rewarmed
- Neurological Function assessed



Targeted Temperature Management

Inclusion Criteria

- Cardiac Arrest/ROSC within 6 Hours
- GCS <8
- SBP ~90
- Mechanical Ventilation
- >16 Year Old Patient

Exclusion Criteria

- ROSC >6hrs Prior
- GCS >8; Purposeful Movements
- Initial Temp <30
- MAP <60 with Interventions
- Terminal Illness
- DNR Standing
- Coagulopathy/Active Bleeding
- Primary Intra-Cranial Event
- Arrest Secondary to Severe Sepsis
- Major Surgery in last 72 Hours
- Trauma Patient at Risk for Bleeding

Cardiac Arrest

- Full Advanced Cardiac Life Support Measures
 - Intubation
 - CPR
 - Defibrillation
 - Medications
 - Epinephrine
 - Bicarbonate
 - Amiodarone
- Correct Abnormalities
 - ABG; pH
 - Acidosis
 - Electrolytes
- Hemodynamic Support
 - Pressors; Inotropes, etc.



<https://cdn.ymaws.com/npofoklahoma.com/resource/collection/C3C77701-15EF-4C38-86ED-993BCB850168/Hypothermia%20in%20Cardiac%20Arrest.pdf>

Vegetative state vs. Brain death

For brain death:

1. ***Absence of Brainstem Reflexes:*** Patient should be unresponsive to stimuli that otherwise would trigger an involuntary response (such as dilation of the pupils in the presence of a bright light)
2. ***Apnea Test:*** The patient, when disconnected from a respirator, should not have respiratory movements and will show other measurable signs supporting the diagnosis of brain death.