Supplemental Material: Dimensional Uniqueness of $\mu_0 = H_0/c$ in Khronon Ghost Condensation

Sheng-Kai Huang[1,*]

[1] *Independent Researcher*

(Dated: April 28, 2026)

We show that the cosmological scale appearing in the Blanchet–Skordis kinetic function $K(Q) = \mu^2(Q-1)^2$ is uniquely fixed, up to an $O(1)$ prefactor, to be $\mu_0 = H_0/c$. Two independent physical constraints—(i) the requirement that the condensate energy density reproduce $\rho_{\rm crit} \sim H_0^2/G$ without a hierarchy, and (ii) the requirement that μ_0 is the IR scale where Petz retrodictability fails—both single out H_0/c as the unique dimensionally consistent choice. Other cosmological scales ($\sqrt{\Lambda}$, $G\rho_{\rm m}/c^4$, particle masses) either reduce to H_0/c at $z = 0$ or fail one of the two constraints. The numerical prefactor 1 versus $1/(2\pi)$ versus 2 is not fixed by dimensional analysis; in the main text the combination $a_0 = c^2\mu_0/(2\pi)$ that appears in the galactic sector singles out the $1/(2\pi)$ factor through the de Broglie–like spatial Fourier normalization.

I. SETUP

[KNOWN] The Blanchet–Skordis (BS) Khronon action [? ?] contains a kinetic function of the timelike scalar

$$K(Q) = \mu^2(Q-1)^2 + \cdots, \tag{1}$$

where $Q = -g^{\mu\nu}\nabla_\mu\tau\nabla_\nu\tau$ is dimensionless and μ has dimension $[\text{length}]^{-1}$ (in units with $\hbar = 1$, equivalently [mass] in natural units). Ghost condensation sits at the minimum $Q = 1$. The cosmological background of Paper 3 is at $Q_0 = 1 + \delta_0$ with $\delta_0 \approx 0.34$.

The dark matter density follows from

$$\rho_K = c^2\,\frac{\mu^2}{8\pi G}\,\delta_0(2+\delta_0), \tag{2}$$

so that the dimensionless density parameter is

$$\Omega_{\rm DM} = \frac{\rho_K}{\rho_{\rm crit}} = \frac{c^2\mu^2}{H_0^2}\,\frac{\delta_0(2+\delta_0)}{3}, \tag{3}$$

using $\rho_{\rm crit} = 3H_0^2/(8\pi G)$.

The question is: *what fixes the scale μ?* The BS papers leave μ as a free parameter set by matching to observations. We argue here that one *specific* choice is selected by dimensional analysis combined with the ghost-condensation IR structure—and that this choice happens to reproduce $\Omega_{\rm DM}$ within 0.5% once δ_0 is fitted, with no further hierarchy.

II. AVAILABLE SCALES

[DERIVED] At cosmological scales today, the available fundamental constants and parameters with dimension are

- the speed of light c ($[c] = \text{m/s}$);
- Newton's constant G ($[G] = \text{m}^3\text{kg}^{-1}\text{s}^{-2}$);
- the Hubble rate H_0 ($[H_0] = \text{s}^{-1}$);
- particle masses m_i ($[m_i] = \text{kg}$, with the lightest neutrino $m_\nu \gtrsim 10^{-2}\,\text{eV}/c^2$);
- the cosmological constant $\Lambda \sim H_0^2/c^2$ (not independent of H_0);
- the Planck mass $M_{\rm Pl} = (\hbar c/G)^{1/2}$ (with $\hbar$ included only for quantum gravity).

We restrict $\hbar$ to enter only via $M_{\rm Pl}$ (the classical limit). Then any inverse length must be expressible as

$$\mu \;\propto\; H_0^n\, c^m\, G^k\, m_i^\ell, \tag{4}$$

for real exponents n, m, k, ℓ satisfying

$$[\text{m}^{-1}] = [\text{s}^{-n}]\cdot[\text{m/s}]^m\cdot[\text{m}^3/(\text{kg}\,\text{s}^2)]^k\cdot[\text{kg}]^\ell, \tag{5}$$

giving three constraints: $3k+m = -1$ (length), $-n-m-2k = 0$ (time), $-k+\ell = 0$ (mass). This is a one-parameter family $\{n, m, k, \ell\} = \{n,\, n-1,\, 0,\, 0\} + (G\text{-axis})$.

Removing irrelevant axes. Ghost condensation is an IR mechanism: the kinetic function $K(Q)$ describes classical, long-wavelength dynamics of the Khronon, so $\hbar$ should not appear, removing the $M_{\rm Pl}$ axis. At cosmological scales the only relevant matter density is $\rho_{\rm m}$; combinations such as $G\rho_{\rm m}/c^2$ have dimension of H_0^2/c^2 on the FRW background by Friedmann's equation, so any G-dependence collapses onto H_0^2. Particle masses m_i are at energies $\gtrsim 10^{15}H_0$ above the cosmological IR scale and would introduce a hierarchy of $\sim 10^{60}$ in $\Omega_{\rm DM}$ if they entered μ directly.

After these reductions, the only dimensionally consistent scale linear in H_0 is

$$\boxed{\mu_0 \;=\; \xi\,\frac{H_0}{c}\,,} \tag{6}$$

with ξ a dimensionless $O(1)$ constant.

* akai@fawstudio.com

III. FIXING THE PREFACTOR

A. Constraint 1: matching $\rho_{\rm crit}$

[DERIVED] Substituting $\mu_0 = \xi H_0/c$ into (**??**),

$$\rho_K = \frac{c^2}{8\pi G}\,\frac{\xi^2 H_0^2}{c^2}\,\delta_0(2+\delta_0) = \frac{\xi^2 H_0^2}{8\pi G}\,\delta_0(2+\delta_0)\,, \quad (7)$$

so that

$$\Omega_{\rm DM} = \frac{\xi^2\,\delta_0(2+\delta_0)}{3}\,. \quad (8)$$

With $\xi = 1$ and $\delta_0 = 0.340$, $\Omega_{\rm DM} = 0.2652$, agreeing with Planck 2018 $\Omega_{\rm DM} = 0.2665$ [**?**] to 0.49%. Any $\xi \neq 1$ requires a compensating shift in δ_0 (or vice versa); $\xi = 1$ is the *minimal* choice with no fine-tuning of the prefactor against δ_0.

B. Constraint 2: IR retrodictability scale

[NEW SYNTHESIS] The Petz recovery framework of Paper 1 [**?**] identifies H_0^{-1} as the *retrodictability horizon*: signals beyond this scale leak from the observable Hilbert space, and the recovery fidelity $F = e^{-\Sigma/2}$ degrades. For Khronon condensation to encode the cosmological arrow of time, the IR scale of the kinetic function $K(Q)$ must coincide with this horizon:

$$\mu^{-1} \sim cH_0^{-1} = d_H, \quad (9)$$

the Hubble length. This is satisfied by $\mu_0 = H_0/c$ and only by that choice. Multiplying or dividing by 2π shifts μ^{-1} by 2π, which corresponds to going from the wavelength to the wavenumber—this is the source of the $1/(2\pi)$ in the galactic sector $a_0 = c^2\mu_0/(2\pi)$.

IV. NUMERICAL EVALUATION

[DERIVED] With $H_0 = 67.4$ km/s/Mpc $= 2.184 \times 10^{-18}$ s^{-1} [**?**]:

$$\mu_0 = \frac{H_0}{c} = \frac{2.184\times10^{-18}\ \mathrm{s}^{-1}}{2.998\times10^{8}\ \mathrm{m/s}} = 7.286\times10^{-27}\ \mathrm{m}^{-1}. \quad (10)$$

The corresponding inverse length is $\mu_0^{-1} = 1.372 \times 10^{26}$ m $\approx d_H$, the Hubble radius.

Cross-check via $\rho_{\rm crit}$.

$$\frac{c^2\mu_0^2}{8\pi G} = \frac{c^2(H_0/c)^2}{8\pi G} = \frac{H_0^2}{8\pi G} = \frac{\rho_{\rm crit}}{3} = 2.844\times10^{-27}\ \mathrm{kg/m^3}, \quad (11)$$

exactly $\rho_{\rm crit}/3$. The factor of 3 is absorbed into $\delta_0(2+\delta_0)$, recovering $\Omega_{\rm DM} = 0.265$.

Cross-check via a_0. The galactic sector predicts the MOND scale

$$a_0 = \frac{c^2\mu_0}{2\pi} = \frac{cH_0}{2\pi} = \frac{(2.998\times10^8)(2.184\times10^{-18})}{2\pi} = 1.042\times10^{-10}\ \mathrm{m/s^2}, \quad (12)$$

within 13% of the empirical $a_0 \approx 1.2\times10^{-10}$ m/s^2 [**?**]. We emphasize this is suggestive, not a derivation of the empirical a_0: the BS galactic sector $J(Y)$ is a free function and a_0 enters as its small-Y asymptotic parameter.

V. WHY OTHER SCALES FAIL

We list candidate inverse-length scales and the constraint they violate.

$2H_0/c$ or $H_0/(2c)$: simple $O(1)$ rescalings change $\Omega_{\rm DM}$ by a factor of ξ^2, which would require shifting δ_0 to 0.158 or 0.812 respectively to restore Planck consistency. The preferred value $\xi = 1$ keeps δ_0 in its natural range $\delta_0 \sim O(0.1)$, with no tuning between prefactor and amplitude.

$\sqrt{\Lambda}$: numerically $\sqrt{\Lambda} \approx 1.27H_0/c$ today, but Λ is constant and H_0 runs. At recombination, $\sqrt{\Lambda}$ is unchanged while $H(z=1100)/c$ is $\sim 10^5\ H_0/c$; the two scales diverge in the past. $\sqrt{\Lambda}$ would freeze the IR scale of $K(Q)$ at the present-day horizon, giving the wrong scaling of $\rho_K(z)$. The running coupling $\mu(a) = H(a)/c$ is discussed in the companion supplement *paper3_running_coupling*.

$H_0^2/(cM_\nu)$ or $1/M_{\rm Pl}$: introduce a 10^{60}–10^{120} hierarchy between μ and the observed $\Omega_{\rm DM}$, requiring fine-tuning at the cosmological constant level.

Particle masses: any $\mu \propto m_i c/\hbar$ sets the IR scale to a Compton wavelength, far below the Hubble horizon, and is incompatible with the ghost condensation IR structure.

VI. CONNECTION TO PETZ RECOVERY (PAPER 1)

[NEW SYNTHESIS] Paper 1 establishes that H_0 is the natural retrodictability scale of the universe: the entropy production rate per Hubble time satisfies $\dot{\Sigma}/H_0 = O(1)$

TABLE I. Candidate cosmological scales for μ and the constraint each fails. Only $\mu_0 = H_0/c$ passes both.

Scale	Numerical	Failure mode
H_0/c	7.3×10^{-27} m^{-1}	— (passes both)
$2H_0/c$	1.5×10^{-26} m^{-1}	$\Omega_{\rm DM} = 1.06$ for fixed δ_0
$H_0/(2c)$	3.6×10^{-27} m^{-1}	$\Omega_{\rm DM} = 0.066$, 4× low
$\sqrt{\Lambda}$	$\sim H_0/c$	degenerate with H_0/c
$H_0^2/(cM_\nu)$	$\sim 10^{-65}$ m^{-1}	$\Omega_{\rm DM} \sim 10^{-77}$
$1/M_{\rm Pl}$	1.6×10^{35} m^{-1}	involves $\hbar$, IR/UV swap
$1/d_{\rm LSS}$	$\sim H_0/c$	last-scattering, degenerate

on cosmological backgrounds. The Khronon Q is the order parameter of this retrodictability, with $Q - 1$ measuring departure from the time-symmetric vacuum. In this picture $\mu_0 = H_0/c$ is not an independent parameter but the IR scale at which recovery fidelity $F = e^{-\Sigma/2}$ is order unity over the observable universe.

The factor of 2π between μ_0 and the galactic a_0 scale arises because a_0 enters the galactic sector through a spatial Fourier kernel (de Broglie–like normalization), while μ_0 enters the cosmological sector through the temporal mass scale of $K(Q)$. Both encode the same horizon, viewed from the kinetic and gradient sides of the Khronon action.

VII. SUMMARY

We have shown that $\mu_0 = H_0/c$ is the unique inverse-length scale built from cosmological constants that (i) does not invoke $\hbar$ or particle masses, (ii) reproduces $\rho_{\rm crit}$ to within the $\delta_0(2 + \delta_0)/3$ factor without a hierarchy, and (iii) coincides with the Petz retrodictability horizon. The $O(1)$ prefactor $\xi = 1$ is selected by the absence of a tuning between the kinetic scale and δ_0; the $1/(2\pi)$ factor in a_0 is the spatial Fourier dual of the same scale. No free parameter remains in the cosmological sector beyond the boundary condition δ_0, and even that is fixed within 2σ by Planck once $\xi = 1$ is imposed.

[1] L. Blanchet and C. Skordis, "Dark matter as a Khronon condensate," JCAP **11**, 040 (2024); arXiv:2404.06584.

[2] L. Blanchet and C. Skordis, "Khronon-Tensor theory reproducing MOND and the cosmological model," (2025); arXiv:2507.00912.

[3] Planck Collaboration, "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. **641**, A6 (2020); arXiv:1807.06209.

[4] M. Milgrom, "A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis," Astrophys. J. **270**, 365 (1983).

[5] S.-K. Huang, "The arrow of time as Petz recovery failure: a unifying framework," (2025); DOI: 10.5281/zenodo.14775757.

[6] N. Arkani-Hamed, H.-C. Cheng, M. A. Luty, and S. Mukohyama, "Ghost condensation and a consistent infrared modification of gravity," JHEP **05**, 074 (2004); arXiv:hep-th/0312099.