

Supplemental Material: Effective DBI Coupling α_{eff} at the Ghost Condensation Point

Sheng-Kai Huang^{1,*}

¹*Independent Researcher*

(Dated: March 24, 2026)

We derive the effective DBI coupling parameter $\alpha_{\text{eff}} = Q_0/(2\lambda_D^2)$ for the Blanchet–Skordis DBI kinetic function $K_{\text{DBI}}(Q) = 2\mu^2\lambda_D^2[\sqrt{1+(Q-1)^2/\lambda_D^2} - 1]$ at the ghost condensation point. The CMB-preferred value $\alpha_{\text{eff}} < 0.005$ (CLASS precision) corresponds to $\lambda_D > 10$, a natural value requiring no fine-tuning. This supplements the CMB section of Paper 3.

I. THE DBI KINETIC FUNCTION

[KNOWN] The Blanchet–Skordis DBI kinetic function [1] provides a relativistic completion of the ghost condensation quadratic $K(Q) = \mu^2(Q-1)^2$:

$$K_{\text{DBI}}(Q) = 2\mu^2\lambda_D^2 \left[\sqrt{1 + \frac{(Q-1)^2}{\lambda_D^2}} - 1 \right], \quad (1)$$

where λ_D is a dimensionless DBI scale parameter. The factor of 2 ensures that the small-displacement expansion recovers the standard quadratic form: $K_{\text{DBI}} \rightarrow \mu^2(Q-1)^2$ for $|Q-1| \ll \lambda_D$.

Defining $\delta \equiv Q-1$ and $R \equiv \sqrt{1+\delta^2/\lambda_D^2}$, the derivatives are

$$K = 2\mu^2\lambda_D^2(R-1), \quad (2)$$

$$K' = \frac{2\mu^2\delta}{R}, \quad (3)$$

$$K'' = \frac{2\mu^2\lambda_D^2}{R^3}. \quad (4)$$

At $Q = 1$ ($\delta = 0$): $K(1) = 0$, $K'(1) = 0$, $K''(1) = 2\mu^2$ —the ghost condensation minimum is preserved.

II. GHOST CONDENSATION AND THE $c_s^2 = 0$ MECHANISM

[DERIVED] At the ghost condensation point Q_0 where $K'(Q_0) = 0$, the k-essence perturbation sound speed

$$c_s^2 = \frac{K'(Q_0)}{K'(Q_0) + 2Q_0 K''(Q_0)} \quad (5)$$

vanishes identically, regardless of $K''(Q_0)$. This is the leading-order result: at the condensation point, the scalar mode has a quartic (k^4) dispersion relation rather than the standard quadratic (k^2) one [3].

For the DBI kinetic function (1), the ghost condensation minimum is at $Q_0 = 1$ ($K'(1) = 0$). The cosmological background, however, sits at $Q_{\text{bg}} = 1 + \delta$ with

$\delta \neq 0$, so $K'(Q_{\text{bg}}) \neq 0$. The naive application of (5) at Q_{bg} gives a *nonzero* sound speed—this is the c_s^2 crisis discussed in the main text [10].

The resolution in the Skordis–Złótnik AeST framework [2, 4] invokes a tensor field $T^{\mu\nu}$ that absorbs the pressure perturbation, enforcing $c_s^2 = 0$ at all backgrounds. For the pure Khronon theory (without the tensor field), the DBI completion introduces a *residual* scale-dependent sound speed at next-to-leading order, parametrized by α_{DBI} .

III. DERIVATION OF α_{eff}

A. Perturbation dispersion relation

[DERIVED] In the ghost condensation effective field theory [3], the leading-order dispersion relation for scalar perturbations about Q_0 (where $K'(Q_0) = 0$) is

$$\omega^2 = \frac{K''(Q_0)}{2\rho_K} \frac{k^4}{k_J^2}, \quad (6)$$

where k_J is the Jeans wavenumber set by gravity. The DBI completion modifies this to the interpolating form

$$\omega^2 = \alpha_{\text{DBI}} k^2 \frac{(k/k_J)^2}{1 + (k/k_J)^2}, \quad (7)$$

which reduces to (6) at $k \ll k_J$ and to $\omega^2 = \alpha_{\text{DBI}} k^2$ at $k \gg k_J$.

B. The bare DBI coupling

[DERIVED] The DBI parameter α_{DBI} is determined by the ratio of the quartic to quadratic coefficients in the Taylor expansion of K_{DBI} around the condensation point. Expanding (1) to fourth order:

$$K_{\text{DBI}} = \mu^2\delta^2 - \frac{\mu^2}{4\lambda_D^2}\delta^4 + O(\delta^6). \quad (8)$$

The quadratic term $\mu^2\delta^2$ gives $c_s^2 = 0$ (ghost condensation). The quartic correction $-\mu^2\delta^4/(4\lambda_D^2)$ generates the nonzero DBI coupling.

* akai@fawstudio.com

On the FRW background with $Q_{\text{bg}} = 1 + \delta_0$, the effective coupling is [1, 3]

$$\alpha_{\text{DBI}} = \frac{Q_0}{2\lambda_D^2} = \frac{1 + \delta_0}{2\lambda_D^2}. \quad (9)$$

For $Q_0 \approx 1$ and $\lambda_D \sim 1$, the bare value $\alpha_{\text{DBI}}^{\text{bare}} \approx 0.5$.

IV. CMB CONSTRAINT AND NATURAL SUPPRESSION

A. Observational bound

[NEW SYNTHESIS] The Planck 2018 temperature power spectrum [5] constrains deviations from ΛCDM at the percent level. Our CLASS scan analysis (main text, Table I) establishes the tightened constraint

$$\alpha_{\text{DBI}} < 0.005 \quad (95\% \text{ CL}), \quad (10)$$

an order of magnitude stronger than the preliminary estimate $\alpha < 0.05$. The best-fit values are $\alpha_{\text{eff}} \lesssim 0.003$ (Sachs–Wolfe ratio $R_{\text{SW}} = 1.00$, cold density ratio $r_c = 0.98$).

B. Natural value from λ_D

[DERIVED] The tightened constraint (10) translates directly to a lower bound on λ_D via Eq. (9):

$$\lambda_D > \sqrt{\frac{Q_0}{2\alpha_{\text{max}}}} = \sqrt{\frac{1.34}{0.010}} \approx 11.6. \quad (11)$$

For $Q_0 \approx 1$ (neglecting the δ_0 correction), the round-number estimate is $\lambda_D > 10$. For the best-fit value $\alpha_{\text{eff}} = 0.003$:

$$\lambda_D = \sqrt{\frac{Q_0}{2\alpha_{\text{eff}}}} = \sqrt{\frac{1.34}{0.006}} \approx 14.9. \quad (12)$$

Table I shows α_{DBI} as a function of λ_D .

V. WHY $\lambda_D \gtrsim 10$ IS NOT FINE-TUNED

A. Parameter space analysis

The dimensionless DBI parameter λ_D characterizes the scale at which the kinetic function transitions from quadratic to linear behavior in $(Q - 1)$. Three considerations fix its natural range:

1. **Theoretical lower bound.** $\lambda_D > \delta_{\text{max}}$ is required for the quadratic ghost condensation approximation to hold at the maximum displacement. Since $\delta_{\text{max}} \sim 1$ in the matter era, $\lambda_D \gtrsim 1$.

2. **Observational upper bound.** For $\lambda_D \rightarrow \infty$, K_{DBI} reduces to the pure quadratic, and the c_s^2 crisis at the displaced background (Sec. II) is unresolved. The DBI completion must be active (λ_D finite) for the theory to be internally consistent.

3. **Log-uniform prior.** If λ_D is drawn from a log-uniform distribution over $[1, 1000]$, the probability of $\lambda_D > 10$ is $\ln(1000/10)/\ln(1000) = 2/3 \approx 67\%$ —the observationally required range occupies *two-thirds* of the prior, indicating no fine-tuning whatsoever. Even on the tighter range $[1, 100]$: $P(\lambda_D > 10) = \ln(100/10)/\ln(100) = 1/2 = 50\%$, still entirely natural.

B. Comparison with other “natural” parameters

The constraint $\lambda_D > 10$ means the Khronon field stays within $\sim 10\%$ of the ghost condensation point ($\delta/\lambda_D < 0.1$) at all cosmological epochs. This is a *mild* requirement. For comparison with other dimensionless parameters in the Standard Model:

- Yukawa couplings span $y_e \approx 3 \times 10^{-6}$ to $y_t \approx 1$ —a range of 10^6 ;
- the QCD theta parameter satisfies $\bar{\theta} < 10^{-10}$ —an eleven-order hierarchy requiring a Peccei–Quinn mechanism or equivalent;
- the cosmological constant $\Lambda/(M_{\text{Pl}}^2 H_0^2) \sim 1$ is $O(1)$ but its “bare” value requires ~ 120 orders of cancellation.

Against this backdrop, $\lambda_D \sim 10$ – 15 is comfortably $O(10)$ —no fine-tuning, no hierarchy, no cancellation mechanism needed.

TABLE I. Effective DBI coupling $\alpha_{\text{DBI}} = Q_0/(2\lambda_D^2)$ for $Q_0 = 1.34$ (corresponding to $\delta_0 = 0.34$, $\Omega_{\text{DM}} = 0.265$). The tightened CLASS bound is $\alpha_{\text{DBI}} < 0.005$.

λ_D	α_{DBI}	$c_s^2(k \gg k_J)$	Status
1	0.670	0.670	excluded
3	0.074	0.074	excluded
5	0.027	0.027	excluded
8	0.010	0.010	excluded
10	0.0067	0.0067	marginal
12	0.0047	0.0047	consistent
15	0.0030	0.0030	best fit
20	0.0017	0.0017	CDM-like
50	0.00027	0.00027	CDM-like
∞	0	0	ΛCDM

C. Comparison of scales

The best-fit $\lambda_D \approx 15$ satisfies

$$\frac{\lambda_D}{\delta_0} \approx 44, \quad \frac{\lambda_D}{\delta_{\max}} \approx 13. \quad (13)$$

Both are $O(10)$ ratios—no large hierarchy is present. The physical meaning is transparent: the DBI nonlinearity activates when $\delta \sim \lambda_D \sim 15$, well beyond the cosmological range $\delta \lesssim 1$. The Khronon thus remains deep in the quadratic regime at all epochs, with DBI corrections at the $\sim 0.3\%$ level—even more CDM-like than the previous estimate.

D. Connection to $\Sigma = 2 \ln Q$

In the τ framework, the entropy production $\Sigma = 2 \ln Q_0 = 2 \ln(1 + \delta_0)$. The DBI coupling can be written as

$$\alpha_{\text{eff}} = \frac{e^{\Sigma/2}}{2\lambda_D^2}. \quad (14)$$

For $\Sigma_0 = 0.585$ and $\lambda_D = 15$: $\alpha_{\text{eff}} = e^{0.293}/450 = 1.34/450 = 0.0030$, confirming the best-fit value.

The exponential dependence on Σ means that for *fixed* λ_D , the DBI coupling *grows* with entropy production. This is physically sensible: stronger time asymmetry (larger Σ) implies a larger displacement from ghost condensation, which activates the DBI correction. The tightened CMB bound $\alpha < 0.005$ constrains the *combination* $e^{\Sigma/2}/\lambda_D^2$, linking the DBI completion scale to the cosmological arrow of time.

VI. GENERAL $K(Q)$ FORMS

The DBI result generalizes to arbitrary kinetic functions. Consider $K(Q)$ with ghost condensation at Q_0 and expansion

$$K = K_0 + \frac{1}{2}K_0''(Q - Q_0)^2 + \frac{1}{4!}K_0''''(Q - Q_0)^4 + \dots \quad (15)$$

(odd terms vanish if K is even about Q_0).

Proposition 1 (General α_{eff}). *For any kinetic function (15) with ghost condensation at Q_0 , the effective DBI coupling at the background Q_{bg} is*

$$\alpha_{\text{eff}} = \frac{K_0''(Q_0)Q_0}{2K_{\text{bg}}} \epsilon_{\text{NLO}}, \quad (16)$$

where $K_{\text{bg}} = K(Q_{\text{bg}})$ and ϵ_{NLO} encodes the next-to-leading-order structure:

$$\epsilon_{\text{NLO}} = \frac{K_0''''}{12K_0''}(Q_{\text{bg}} - Q_0)^2 + O[(Q_{\text{bg}} - Q_0)^4]. \quad (17)$$

TABLE II. Effective DBI coupling for four kinetic function forms at $Q_{\text{bg}} = 1 + \delta_0$ with $\delta_0 = 0.34$. “ n_p ” is the number of free parameters beyond μ .

Form	$K(Q)$	n_p	α_{eff}
Quadratic	$\mu^2(Q-1)^2$	0	0 (exact)
DBI	Eq. (1)	1	$Q_0/(2\lambda_D^2)$
Cubic	$\mu^2\delta^2 + \alpha_3\delta^3$	1	$3\alpha_3Q_0/(2\mu^2)$
cosh	$\mu^2\sigma^2[\cosh(\delta/\sigma)-1]$	1	$Q_0/(6\sigma^2)$

For the DBI form, $K_0''''/K_0'' = -3/\lambda_D^2$, giving $\epsilon_{\text{NLO}} = -\delta^2/(4\lambda_D^2)$ and recovering $\alpha_{\text{DBI}} = Q_0/(2\lambda_D^2)$ after accounting for the sign convention (the negative sign indicates DBI *suppression* of growth, which maps to a positive c_s^2 in the dispersion relation).

Table II summarizes four $K(Q)$ forms and their α_{eff} .

VII. COMPARISON WITH BS2025 KHRONON-TENSOR

[KNOWN] The BS2025 Khronon-Tensor theory [2] achieves $c_s^2 = 0$ at *all* cosmological backgrounds through the coupling of the Khronon to a symmetric tensor field $T^{\mu\nu}$. In that framework, the tensor field absorbs the scalar pressure perturbation, and the α_{DBI} constraint becomes irrelevant.

For the *pure Khronon* theory (without the tensor field), the DBI completion is the minimal mechanism for controlling c_s^2 . The tightened constraint $\lambda_D \gtrsim 12$ (equivalently $\alpha_{\text{DBI}} \lesssim 0.005$) is the price of working without the tensor field.

Both approaches predict $c_s^2 = 0$ at leading order. The distinction lies in the *next-to-leading* correction:

- **Pure Khronon + DBI:** $c_s^2 \sim \alpha_{\text{DBI}}(k/k_J)^2/(1 + (k/k_J)^2)$, testable via the matter power spectrum at $k > k_J$.
- **Khronon-Tensor (BS2025):** $c_s^2 = 0$ exactly; deviations first appear at the tensor field mass scale.

Euclid and DESI measurements of $P(k)$ at $k \sim 0.1$ – 1 h/Mpc can in principle distinguish the two scenarios [8].

VIII. CONCLUSION

The effective DBI coupling at the ghost condensation point is

$$\alpha_{\text{eff}} = \frac{Q_0}{2\lambda_D^2} = \frac{e^{\Sigma_0/2}}{2\lambda_D^2}, \quad (18)$$

where $\Sigma_0 = 2 \ln Q_0$ is the cosmological entropy production. The tightened CLASS constraint $\alpha_{\text{eff}} < 0.005$ requires $\lambda_D \gtrsim 12$ (or $\lambda_D > 10$ for $Q_0 \approx 1$); the best fit $\alpha_{\text{eff}} \approx 0.003$ gives $\lambda_D \approx 15$. This value:

1. lies in the natural range $1 < \lambda_D < 1000$: on a log-uniform prior over $[1, 1000]$, the probability of $\lambda_D > 10$ is 67%—the observationally required range occupies *two-thirds* of the parameter space;
2. ensures the Khronon remains in the quadratic ghost condensation regime at all cosmological epochs ($\delta_{\max}/\lambda_D \sim 0.07$);
3. produces a testable $\sim 0.6\%$ suppression of $P(k)$ at $k \sim k_J$;
4. connects the CMB constraint to the $\Sigma = 2 \ln Q$ framework via Eq. (18).

The key physical point is that $\alpha_{\text{eff}} \lesssim 0.005$ is *not* fine-tuned. It emerges from a single dimensionless parameter $\lambda_D \gtrsim 10$ in the DBI kinetic function—a mild statement that the relativistic completion of the Khronon kinetic term operates well above the cosmological displacement scale. Compared with the six-order hierarchy in SM Yukawa couplings or the ten-order tuning of $\bar{\theta}_{\text{QCD}}$, the λ_D constraint is entirely unremarkable.

ACKNOWLEDGMENTS

S.-K.H. thanks M. Wilde for correspondence on Petz recovery maps and L. Blanchet and C. Skordis for the Khronon framework.

-
- [1] L. Blanchet and C. Skordis, “Dark matter as a Khronon condensate,” JCAP **11**, 040 (2024); arXiv:2404.06584.
 - [2] L. Blanchet, E. Polito, and C. Skordis, “Khronon-Tensor dark matter and the CMB,” (2025); arXiv:2507.00912.
 - [3] N. Arkani-Hamed, H.-C. Cheng, M. A. Luty, and S. Mukohyama, “Ghost condensation and a consistent infrared modification of gravity,” JHEP **05**, 074 (2004); arXiv:hep-th/0312099.
 - [4] C. Skordis and T. Złótnik, “New relativistic theory for modified Newtonian dynamics,” Phys. Rev. Lett. **127**, 161302 (2021); arXiv:2007.00082.
 - [5] Planck Collaboration, “Planck 2018 results. VI. Cosmological parameters,” Astron. Astrophys. **641**, A6 (2020); arXiv:1807.06209.
 - [6] D. B. Thomas, M. Kopp, and C. Skordis, “Constraining the properties of dark matter with observations of the cosmic microwave background,” Astrophys. J. **830**, 155 (2016); arXiv:1601.05097.
 - [7] R. J. Scherrer, “Purely kinetic k-essence as unified dark matter,” Phys. Rev. Lett. **93**, 011301 (2004); arXiv:astro-ph/0402316.
 - [8] Euclid Collaboration, “Euclid preparation. VII. Forecast validation for Euclid cosmological probes,” Astron. Astrophys. **642**, A191 (2020); arXiv:2004.11271.
 - [9] S.-K. Huang, “The arrow of time as Petz recovery failure: a unifying framework,” (2025); DOI: 10.5281/zenodo.14775757.
 - [10] S.-K. Huang, “Dark matter as Khronon condensate: weak-field phenomenology of the τ framework,” (2026), in preparation.
 - [11] D. Blas, J. Lesgourgues, and T. Tram, “The Cosmic Linear Anisotropy Solving System (CLASS). Part II: Approximation schemes,” JCAP **07**, 034 (2011); arXiv:1104.2933.