One Equation, Three Scales: $\Sigma = D(\rho_{\text{spacetime}} \| \rho_{\text{matter}})$ as the Universal Action

Sheng-Kai Huang[1,*]

[1] *Independent Researcher*

(Dated: March 10, 2026)

[**TO WRITE:** Final abstract after Papers 1–3 are complete.]

Papers 1–3 of this series demonstrated that the entropy production $\Sigma = D(\rho\|\sigma) - D(\mathcal{N}(\rho)\|\mathcal{N}(\sigma))$ governs three seemingly unrelated domains: quantum information recovery (Paper 1), strong-field gravity near black holes (Paper 2), and galactic rotation curves (Paper 3). Here we propose that these are not three separate applications but three solutions of *one equation*:

$$\Sigma = D(\rho_{\text{spacetime}} \| \rho_{\text{matter}}),$$

where $\rho_{\text{spacetime}}$ and ρ_{matter} are density matrices encoding the geometry and matter content, respectively. We argue that three mathematical realizations—modular flow, gravitational Landauer principle, and quantum channel analysis—all converge on the universal identity $\Sigma_{\text{grav}} = -\ln(-g_{00})$, independent of any specific action principle. Bianconi's gravity-from-entropy framework (PRD 2025) provides one formal setting for $\Sigma = D(\rho_{\text{spacetime}} \| \rho_{\text{matter}})$, though its vacuum solutions give Schwarzschild $+O(r^{-4})$ corrections [35], not the exponential metric. The exponential metric instead follows from promoting the identity to an exact equation. Different boundary conditions yield different solutions: strong-field ($r \sim r_s$) $\to$ exponential metric, weak-field ($r \sim$ kpc) $\to$ running G and flat rotation curves, no-gravity limit $\to$ standard quantum information theory. We identify the matching conditions between regimes, derive the emergent arrow of time $\tau = 1 - \exp(-\Sigma/2)$ as a universal consequence, and discuss open problems including the Born rule, Planck-scale physics, and cosmological initial conditions.

THE THREE SCALES

Content outline

[NEW SYNTHESIS] *Summary of Papers 1–3.*—The first three papers of this series established that the entropy production Σ governs physics at three different scales:

TABLE I. The three scales unified by Σ.

	Paper 1	**Paper 2**	**Paper 3**
Domain	QI	Strong field	Weak field
Scale	any	$r \sim r_s$	$r \sim$ kpc
Σ	generic	r_s/r	$r_s/r + \alpha \ln r$
$\mathcal{N}$	any CPTP	grav. redshift	RG-improved grav.
Key result	$F \geq e^{-\Sigma/2}$	exp. metric	flat curves
Status	[KNOWN]	[NEW SYNTHESIS]	[NEW SYNTHESIS]

[NEW SYNTHESIS] *The pattern.*—Each paper uses the *same* mathematical structure:

1. A quantum channel $\mathcal{N}$ that processes information;
2. The QRE drop Σ under $\mathcal{N}$;
3. The Petz recovery map as the optimal retrodiction;
4. The fidelity bound $F \geq \exp(-\Sigma/2)$.

The only difference is the *channel*: a general CPTP map (Paper 1), the gravitational redshift channel (Paper 2), or the RG-improved gravitational channel (Paper 3). This suggests that all three are limits of a single, more fundamental equation.

The $F = ma$ analogy.—Newton's second law $F = ma$ is one equation. With different force laws and boundary conditions:

- Gravity + central field $\to$ Kepler orbits (planets);
- Hooke's law + spring $\to$ simple harmonic oscillation;
- Viscosity + continuum $\to$ Navier–Stokes (fluids).

Nobody says there are "three different $F = ma$'s." There is one equation with three classes of solutions. We propose the same for Σ.

THE UNIFIED EQUATION

Content outline

[SPECULATIVE] *The proposal.*—We conjecture that gravity, quantum information, and the arrow of time are all governed by

$$\boxed{\Sigma = D(\rho_{\text{spacetime}} \| \rho_{\text{matter}}).} \tag{1}$$

Interpretation.—$\rho_{\text{spacetime}}$ encodes the geometry (metric, curvature) as a density matrix. ρ_{matter} encodes the

matter content (stress-energy) as a density matrix. The quantum relative entropy between them is the entropy production Σ: the "distance" between what spacetime "is" and what matter "requires" it to be.

[KNOWN] *Formal framework: Bianconi (2025).*—Bianconi [4] provided the mathematical framework in which Eq. (1) is well defined. In the gravity-from-entropy (GfE) framework:

- The metric $g_{\mu\nu}$ is promoted to a density matrix ρ_g on the space of field configurations;
- The matter-induced metric (the metric that matter "wants" via stress-energy) is another density matrix ρ_m;
- The gravitational action is the QRE: $S_{\text{grav}} = D(\rho_g \| \rho_m)$;
- The Einstein equations arise from extremizing $D(\rho_g \| \rho_m)$;
- In the low-coupling limit, this reduces to the standard Einstein–Hilbert action.

[KNOWN] *Bianconi's key results* [4, 5]:

- Modified Einstein equations that reduce to standard GR at low coupling;
- Predicts small positive cosmological constant;
- The "G-field" (extra degree of freedom from QRE structure) is a candidate for dark matter;
- FRW cosmological solutions exist and are thermal [5].

[NEW SYNTHESIS] *Connection to our framework.*—Bianconi's $S_{\text{grav}} = D(\rho_g \| \rho_m)$ is *literally* our $\Sigma = D(\rho_{\text{spacetime}} \| \rho_{\text{matter}})$. The GfE framework provides the formal definition; our series provides the physical interpretation: Σ is the entropy production, $\tau = 1 - \exp(-\Sigma/2)$ is the arrow of time, and the Petz recovery map is the optimal retrodiction.

[SPECULATIVE] *Critical caveat: GfE vacuum solutions.*—Thattarampilly et al. [35] solved the full GfE field equations for a static, spherically symmetric vacuum and found $g_{00} = -(1 - r_s/r - \beta r_s^2/(48r^4))$: Schwarzschild with $O(r^{-4})$ perturbative corrections, *not* the exponential metric (which has $O(r^{-2})$ corrections). This means GfE in its current form does not produce the exponential metric as a vacuum solution. Our unified equation $\Sigma = D(\rho_{\text{spacetime}} \| \rho_{\text{matter}})$ should therefore be understood as a *principle*—the entropy production equals the QRE between geometry and matter—rather than being identified with any specific action functional.

[NEW SYNTHESIS] *Three routes independent of GfE.*—The identity $\Sigma_{\text{grav}} = -\ln(-g_{00})$ is supported by three independent derivations that do not depend on any specific action principle:

1. *Modular flow*: The fractional QRE loss computed via Dorau–Much modular theory [6] gives $(D_{\text{in}} - D_{\text{out}})/D_{\text{in}} = r_s/r$.
2. *Gravitational Landauer*: The Tolman-modified erasure cost [36] gives $\Sigma = -\ln(-g_{00})$.
3. *Quantum channel*: A pure-loss bosonic channel with $\eta = -g_{00}$ gives $\Sigma = -\ln \eta$; Petz recovery for this channel has been experimentally demonstrated [37].

The exponential metric follows from promoting the identity $\sqrt{-g_{00}} = e^{-\Sigma/2}$ to an exact equation, not from solving modified field equations.

STRONG-FIELD SOLUTION

Content outline

[NEW SYNTHESIS] *Boundary conditions.*—Static, spherically symmetric, $r \sim r_s$, asymptotically flat.

From Paper 2.—Under these conditions, the entropy production is [2]

$$\Sigma_{\text{strong}} = \frac{r_s}{r} = \frac{2GM}{c^2 r}, \tag{2}$$

and the Petz recovery fidelity gives the metric component

$$g_{00} = -F^2 = -\exp\left(-\frac{r_s}{r}\right). \tag{3}$$

This is the "exponential metric" that replaces Schwarzschild:

- No event horizon (the exponential never reaches zero);
- All curvature invariants remain finite;
- Testable predictions: GW echoes, neutron star redshift, QNM frequencies, shadow size, ISCO radius.

[**TO DERIVE:** Show that Eq. (2) follows from Eq. (1) under the stated boundary conditions. Specifically: solve $\delta D(\rho_g \| \rho_m) = 0$ for a static, spherically symmetric vacuum with mass M at the origin. The solution should be $g_{00} = -\exp(-r_s/r)$, reproducing Paper 2.]

[**TO DERIVE:** Resolve the tension: GfE gives Schwarzschild $+O(r^{-4})$ corrections [35], while our identity gives the exponential metric. Three possible resolutions: (a) The exponential metric is the correct vacuum geometry and GfE needs modification; (b) The true vacuum geometry is Schwarzschild + corrections, and the exponential metric is an idealization; (c) The unified equation $\Sigma = D(\rho_{\text{st}} \| \rho_{\text{m}})$ requires a different mathematical realization than GfE. This is the central open problem of the program.]

Matching condition at large r.—As $r \to \infty$, $\Sigma_{\text{strong}} \to 0$ and $g_{00} \to -1$: flat spacetime. The strong-field solution must match onto the weak-field solution (Paper 3) at intermediate radii $r \gg r_s$ but $r \ll r_c \approx 37$ kpc.

WEAK-FIELD SOLUTION

Content outline

[NEW SYNTHESIS] *Boundary conditions.*—Quasi-static, approximately axisymmetric (disk galaxy), $r \sim$ kpc, weak-field ($\Phi/c^2 \ll 1$).

From Paper 3.—Under these conditions, the RG flow of the gravitational channel introduces quantum corrections [3]:

$$\Sigma_{\text{weak}}(r) = \frac{r_s}{r} + \frac{4k_* r_s}{\pi} \ln\left(\frac{r}{r_0}\right), \tag{4}$$

where the logarithmic term arises from the marginal running ($\eta = 1$) of Newton's constant $G(k)$ at IR scales $k \ll k_*$.

Physical consequence.—The logarithmic correction produces an additional $1/r$ force that dominates at $r \gg r_c = 1/k_*$, giving flat rotation curves without dark matter particles.

[TO DERIVE: Show that Eq. (4) follows from Eq. (1) under weak-field boundary conditions with quantum corrections included. This requires: (i) solving $\delta D(\rho_g \| \rho_m) = 0$ at the semiclassical level (reproducing Dorau–Much); (ii) including one-loop corrections via the DPI/RG connection (Casini–Huerta); (iii) showing that the marginal running $\eta = 1$ emerges naturally.]

Matching to strong field.—At $r \gg r_s$ but $r \ll r_c$, the logarithmic correction is negligible and $\Sigma_{\text{weak}} \approx r_s/r$, matching Σ_{strong} smoothly. The two solutions are not disjoint—they are limits of the same equation.

QUANTUM INFORMATION SOLUTION

Content outline

[KNOWN] *Boundary conditions.*—No gravity ($G = 0$), general CPTP map $\mathcal{N}$, finite-dimensional Hilbert spaces.

From Paper 1.—With no gravity, Σ reduces to the generic entropy production of any quantum channel [1]:

$$\Sigma_{\text{QI}} = D(\rho \| \sigma) - D(\mathcal{N}(\rho) \| \mathcal{N}(\sigma)), \tag{5}$$

and the Petz recovery map $\mathcal{R}_{\sigma,\mathcal{N}}$ provides the optimal retrodiction with fidelity bound $F \geq \exp(-\Sigma/2)$.

[NEW SYNTHESIS] *The gravity $\to$ QI limit.*—When gravity is turned off, $\rho_{\text{spacetime}} \to \rho_{\text{flat}}$ (Minkowski geometry) and the QRE $D(\rho_{\text{spacetime}} \| \rho_{\text{matter}})$ reduces to the matter-sector entropy production alone. The gravitational contribution vanishes: $\Sigma_{\text{grav}} = 0$, $\tau_{\text{grav}} = 0$. Only the channel-induced entropy production remains.

Connection to quantum error correction.—From Paper 1, $\Sigma = 0 \Leftrightarrow \tau = 0 \Leftrightarrow$ perfect Petz recovery $\Leftrightarrow$ quantum Markov chain $\Leftrightarrow$ Knill–Laflamme conditions. QEC is the deliberate construction of a subsystem where $\Sigma = 0$: an artificial elimination of the arrow of time within a local code space.

MATCHING CONDITIONS AND PREDICTIONS

Content outline

[NEW SYNTHESIS] *The three regimes are connected.*—The solutions of Eq. (1) in different boundary conditions must *match* at their boundaries:

1. *Strong $\to$ weak*: At $r \gg r_s$, the exponential metric linearizes to Newton + small corrections. The logarithmic correction from Paper 3 emerges as the leading quantum correction to this classical limit.

2. *Weak $\to$ QI*: At scales where $\Phi/c^2 \to 0$ (laboratory), the gravitational Σ becomes negligible and only quantum-channel entropy production remains.

3. *Weak $\to$ cosmological*: At $r \sim r_H = c/H_0$ (Hubble radius), the de Sitter background contribution $\Sigma_{\text{dS}} \sim r/r_H$ becomes dominant, connecting galactic dynamics to cosmic expansion.

[SPECULATIVE] *Intermediate-scale predictions.*—At scales where two regimes overlap, *both* corrections matter simultaneously. Specific predictions:

- At $r \sim 100$ kpc (galaxy cluster outskirts): both running-G and de Sitter corrections contribute. This may resolve the Bullet Cluster puzzle if the two effects combine to produce the observed mass-light separation.

- At $r \sim r_s \cdot k_*/k_s$ (strong-field with quantum corrections): the exponential metric acquires logarithmic modifications. Testable via precision measurements of neutron star properties.

[SPECULATIVE] *The three-term approximation.*—In the appropriate limits, the unified Σ can be approximated as

$$\Sigma(r) \approx \underbrace{\frac{r_s}{r}}_{\text{local gravity}} + \underbrace{\alpha \ln\left(\frac{r}{r_c}\right)}_{\text{RG running}} + \underbrace{\beta \frac{r}{L_{\text{dS}}}}_{\text{de Sitter}}, \tag{6}$$

where:

- Term 1 dominates near compact objects ($r \lesssim r_s$);

- Term 2 dominates at galactic scales ($r_c \lesssim r \lesssim L_{\rm dS}$);
- Term 3 dominates at cosmological scales ($r \sim L_{\rm dS}$).

This is an approximation, not the fundamental equation. The fundamental equation is Eq. (1); the three terms emerge as limits.

THE ARROW OF TIME AS EMERGENT

Content outline

[NEW SYNTHESIS] *The central thesis.*—The arrow of time is not a separate law of nature. It *is* the entropy production Σ:

$$\tau = 1 - \exp(-\Sigma/2). \tag{7}$$

This single equation, applied across scales, gives:

TABLE II. The arrow of time at different scales.

Scale	Σ	τ	**Physical meaning**
Closed system	0	0	No arrow of time
Open (lab)	> 0	> 0	Arrow exists
Near BH	r_s/r	$\to 1$	Strong arrow
Galaxy edge	$+\alpha \ln r$	> 0	Arrow from RG
Hubble horizon	~ 1	~ 0.39	Cosmic arrow
Singularity	$\to \infty$	$= 1$	Maximal arrow

[NEW SYNTHESIS] *Three key insights.*—

1. $\tau = 0$ *(closed system): time does not exist.*—When $\Sigma = 0$, perfect Petz recovery is possible: the "future" state can be perfectly retrodicted from the "past" state, and vice versa. Past and future are operationally indistinguishable. This is not a philosophical statement but an operational one: no measurement can determine the temporal ordering of events.
2. $\tau > 0$ *(open system): time's arrow exists.*—When $\Sigma > 0$, retrodiction is imperfect: there is a preferred direction. The degree of preference is continuously quantified by $\tau \in (0, 1)$.
3. $\tau \to 1$ *(macroscopic/singularity): classical world.*—When $\tau \approx 1$, the arrow of time is absolute. Collapse is complete. Classical physics emerges. This is the everyday world.

[NEW SYNTHESIS] *The signed version.*—The full picture requires

$$\tau_{\rm signed} = \ln\left(\frac{P_{\rm fwd}}{P_{\rm rev}}\right), \tag{8}$$

which allows three cases:

- $\tau_{\rm signed} > 0$: forward arrow (normal);
- $\tau_{\rm signed} = 0$: no arrow (closed/unitary);
- $\tau_{\rm signed} < 0$: reverse arrow (non-Markovian backflow, spin echo, Wang et al. 2002 colloidal particle experiments [20]).

The second law ($\langle\Sigma\rangle \geq 0$) is a *statistical* law: $P(\Sigma < 0)/P(\Sigma > 0) = \exp(-|\Sigma|)$ (Crooks). Time reversal is not impossible, merely exponentially unlikely for macroscopic systems.

COSMOLOGICAL EXTENSION

Content outline

[SPECULATIVE] *FRW from* Σ.—In the Friedmann–Robertson–Walker (FRW) universe, there is no timelike Killing vector. The natural replacement is the Kodama vector K^a [14], which:

- Is divergence-free in any spherically symmetric spacetime;
- Gives a conserved charge = Misner–Sharp mass;
- Defines surface gravity at the apparent horizon.

The cosmological τ can be defined via [34]:

$$F_K = \frac{|K|}{|K_{\rm ref}|}, \tag{9}$$

giving $\tau = 0$ at the observer ($r = 0$) and $\tau = 1$ at the apparent horizon ($|K|^2 = 0$).

[SPECULATIVE] *Friedmann equations from QRE.*—The first law of thermodynamics at the FRW apparent horizon, $dE = TdS + WdV$, projects along the Kodama vector to give the second Friedmann equation [15, 16]. Combined with Dorau–Much and Bianconi, this suggests:

$$H^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho \quad \Leftarrow \quad \delta\, D(\rho_{\rm st}\|\rho_{\rm m}) = 0. \tag{10}$$

[**TO DERIVE:** Derive the Friedmann equations from $\Sigma = D(\rho_{\rm spacetime}\|\rho_{\rm matter})$ using Bianconi's GfE framework with FRW boundary conditions. Bianconi [5] has shown FRW solutions exist; the task is to reproduce the standard Friedmann equations as a limit.]

[SPECULATIVE] *Dark energy from finite cosmic information.*— Padmanabhan's CosMIn principle [17]: the total cosmic information $I_{\rm CosMIn}$ transferred from quantum to classical must be *finite*. This requires late-time accelerated expansion ($\Lambda > 0$). In the τ language:

$$I_{\rm CosMIn} = \int_0^\infty \tau(t)\, dt < \infty \quad \Rightarrow \quad \Lambda > 0. \tag{11}$$

[**TO WRITE:** Paragraph on the interpretation: $\Lambda > 0$ is not a "mystery" but a *necessity* for finite cosmic information processing. The universe must eventually stop producing information (entropy) at an unbounded rate. Accelerated expansion achieves this by diluting matter and radiation.]

THE ANALOGY MADE PRECISE

Content outline

[NEW SYNTHESIS] *One equation, three solutions.*—We now make the $F = ma$ analogy precise:

What the analogy captures.—

- Just as $F = ma$ does not tell you the force law (you need to specify gravity, Hooke, etc.), $\Sigma = D(\rho_{\rm st}\|\rho_{\rm m})$ does not tell you the channel (you need to specify the boundary conditions).
- Just as different solutions of $F = ma$ look completely different (ellipses vs. oscillations vs. turbulence), different solutions of the unified Σ look completely different (black holes vs. rotation curves vs. quantum channels).
- Just as $F = ma$ has a deeper formulation (Lagrangian mechanics, Hamilton's principle), the unified Σ has a deeper formulation via the Petz recovery map as the canonical retrodiction functor [10].

OPEN PROBLEMS

Content outline

[**TO WRITE:** Each open problem should be 2–3 paragraphs: statement, current status, possible approach.]

Problem 1: First-principles derivation.—Derive $\Sigma = D(\rho_{\rm spacetime}\|\rho_{\rm matter})$ from more primitive axioms. Currently, Eq. (1) is a *conjecture* motivated by Bianconi's GfE [4], Dorau–Much's QRE $\to$ Einstein equations [6], and the consistency of Papers 1–3. A derivation from axioms (e.g., von Neumann algebra axioms, or the axioms of algebraic QFT) would elevate it from conjecture to theorem.

[SPECULATIVE] Current best candidates: (i) Bianconi's entropic action principle [4], though its vacuum solutions give Schwarzschild $+O(r^{-4})$ corrections rather than the exponential metric [35], so it cannot serve as the mathematical realization in its current form; (ii) Dorau–Much's modular theory [6] extended to include backreaction; (iii) Jacobson's entanglement equilibrium [12] rewritten in QRE language. The three action-independent routes (modular flow, gravitational Landauer, quantum channel) may provide the necessary foundation without requiring a specific modified gravity action.

Problem 2: Planck scale.—What happens when spacetime itself is quantum? At the Planck scale, $\rho_{\rm spacetime}$ is not a semiclassical density matrix but a fully quantum object. The QRE $D(\rho_{\rm spacetime}\|\rho_{\rm matter})$ must be replaced by an algebraic QFT version using the Araki relative entropy [13]. This is mathematically well defined but computationally intractable with current methods.

Problem 3: Cosmological initial conditions.—Why does the universe start with $\Sigma \approx 0$ (low entropy)? In the τ framework, this is the question: why does $\tau_{\rm initial} \approx 0$? One possibility: the initial state is a "Petz-recovered" state from the final condition (a closed-universe scenario where boundary conditions at both ends enforce $\Sigma = 0$). This is highly speculative and connects to the "past hypothesis" of thermodynamics [22, 23].

Problem 4: The Born rule.—The τ framework assumes the Born rule in computing fidelity F and entropy production Σ. A complete theory should derive the Born rule, not assume it. Possible connections:

- Zurek's "envariance" derivation [21] uses environment-assisted symmetry, which connects to τ via quantum Darwinism;
- Deutsch–Wallace decision-theoretic approach relies on rational betting, which maps to Petz recovery optimality;
- Gleason's theorem constrains probability measures on Hilbert space, and the Petz map is the unique "probability measure on channels."

[SPECULATIVE] *Problem 5: Consciousness.*—We acknowledge but do not attempt this problem. The hard problem of consciousness (why subjective experience exists) is sometimes connected to the arrow of time and measurement. The τ framework provides a precise, quantitative arrow of time that does not require an observer. Whether this has implications for consciousness is a question for philosophers, not for this paper.

Problem 6: Derive $\eta = 1$.—The marginal anomalous dimension $\eta = 1$ that produces flat rotation curves (Paper 3) should follow from the properties of $D(\rho_{\rm spacetime}\|\rho_{\rm matter})$ in the IR. This is the most concrete open problem and the one most likely to be resolved in the near term. The DPI structure of QRE may constrain η to exactly 1 via a saturation argument.

Problem 7: Gravitational waves and dynamical spacetimes.— Papers 2 and 3 address static or quasi-static spacetimes. Extending to dynamical spacetimes (gravitational wave emission, binary mergers) requires:

- A time-dependent $\Sigma(t)$;
- The Kodama vector (for spherical symmetry) or its generalization [31];

TABLE III. The $F = ma$ analogy for $\Sigma = D(\rho_{\mathrm{st}}\|\rho_{\mathrm{m}})$.

	$F = ma$	$\Sigma = D(\rho_{\mathrm{st}}\|\rho_{\mathrm{m}})$	**Status**
Equation	$\mathbf{F} = m\mathbf{a}$	$\Sigma = D(\rho_{\mathrm{st}}\|\rho_{\mathrm{m}})$	[SPECULATIVE]
Solution 1	Kepler orbits (gravity, $r \gg 0$)	Exponential metric (strong field, $r \sim r_s$)	[NEW SYNTHESIS]
Solution 2	SHM (Hooke, spring)	Flat rotation curves (weak field, $r \sim$ kpc)	[NEW SYNTHESIS]
Solution 3	Navier–Stokes (continuum)	Standard QI ($G = 0$, any CPTP)	[KNOWN]
Solution 4	—	FRW cosmology ($r \sim L_H$)	[SPECULATIVE]
Matching	Virial theorem	$\Sigma_{\mathrm{strong}} \to \Sigma_{\mathrm{weak}}$ at $r \gg r_s$	**[TO DERIVE:]**
Deep structure	Lagrangian mechanics	Petz recovery map as retrodiction functor	[KNOWN]

- Raychaudhuri decomposition of $d\Sigma/ds$ along geodesic congruences.

CONCLUSION

Content outline

[TO WRITE: Concluding section synthesizing the full four-paper arc.]

Summary.—We have argued that a single equation, $\Sigma = D(\rho_{\mathrm{spacetime}}\|\rho_{\mathrm{matter}})$, unifies three scales of physics:

- Quantum information (Paper 1): $\Sigma = 0 \Leftrightarrow$ perfect recovery, no arrow of time;
- Strong-field gravity (Paper 2): $\Sigma = r_s/r \to$ exponential metric, information-theoretic "soft horizon";
- Galactic dynamics (Paper 3): $\Sigma = r_s/r + \alpha \ln r \to$ running G, flat rotation curves without dark matter particles.

The philosophical point.—

> We have not discovered a new equation. We have proposed that an existing mathematical structure—quantum relative entropy and its recovery theory—when applied consistently across scales, unifies phenomena that appear unrelated. The contribution is not a new formula but a new viewpoint that makes existing results cohere. This is the kind of contribution that, if correct, simplifies rather than complicates.

[TO WRITE: Final paragraph on falsifiability: the framework makes specific, testable predictions at each scale (Paper 2: GW echoes, shadow sizes; Paper 3: universal r_c, SPARC fits, BIG-SPARC). If these predictions fail, the framework fails. This is a feature, not a bug.]

The author thanks the quantum information, quantum gravity, astrophysics, and quantum foundations communities for the foundational works on which this synthesis builds.

* akai@fawstudio.com

[1] S.-K. Huang, "Universal recovery: The Petz map as retrodiction functor, error corrector, and entropy bound," companion paper (2026).
[2] S.-K. Huang, "Information-theoretic origin of the exponential metric: Gravitational redshift as Petz recovery fidelity," companion paper (2026).
[3] S.-K. Huang, "Flat rotation curves from quantum relative entropy: Running Newton's constant as information processing," companion paper (2026).
[4] G. Bianconi, "Gravity from entropy," Phys. Rev. D **111**, 066001 (2025); arXiv:2408.14391.
[5] G. Bianconi, "Gravity from entropy: Cosmological solutions," (2025); arXiv:2510.22545.
[6] M. Dorau and A. Much, "From quantum relative entropy to the semiclassical Einstein equations," Phys. Rev. Lett. (2025); arXiv:2510.24491.
[7] H. Casini and M. Huerta, "Relative entropy and the RG flow," J. High Energy Phys. **2017**, 044 (2017); arXiv:1611.00016.
[8] N. Kumar, "Marginal IR running of gravity as a natural explanation for dark matter," Phys. Lett. B **871**, 140008 (2025); arXiv:2509.05246.
[9] G. Gubitosi, O. F. Piattella, and R. M. Casarini, "Phenomenology of renormalization group improved gravity from the kinematics of SPARC galaxies," (2024); arXiv:2403.00531.
[10] A. J. Parzygnat and F. Buscemi, "Axioms for retrodiction: Achieving time-reversal symmetry with a prior," Quantum **7**, 1013 (2023).

[11] T. Jacobson, "Thermodynamics of spacetime: The Einstein equation of state," Phys. Rev. Lett. **75**, 1260 (1995).

[12] T. Jacobson, "Entanglement equilibrium and the Einstein equation," Phys. Rev. Lett. **116**, 201101 (2016); arXiv:1505.04753.

[13] H. Araki, "Relative entropy of states of von Neumann algebras," Publ. RIMS Kyoto **11**, 809 (1976).

[14] H. Kodama, "Conserved energy flux for the spherically symmetric system and the backreaction problem in the black hole evaporation," Prog. Theor. Phys. **63**, 1217 (1980).

[15] S. A. Hayward, "Unified first law of black-hole dynamics and relativistic thermodynamics," Class. Quantum Grav. **15**, 3147 (1998).

[16] R.-G. Cai and S. P. Kim, "First law of thermodynamics and Friedmann equations of Friedmann–Robertson–Walker universe," J. High Energy Phys. **2005**, 050 (2005).

[17] T. Padmanabhan, "Cosmic information, the cosmological constant, and the amplitude of primordial perturbations," Phys. Lett. B **773**, 81 (2017); arXiv:1703.06144.

[18] E. P. Verlinde, "Emergent gravity and the dark universe," SciPost Phys. **2**, 016 (2017); arXiv:1611.02269.

[19] L. Smolin, "MOND as a regime of quantum gravity," Phys. Rev. D **96**, 083523 (2017); arXiv:1704.00780.

[20] G. M. Wang, E. M. Sevick, E. Mittag, D. J. Searles, and D. J. Evans, "Experimental demonstration of violations of the second law of thermodynamics for small systems and short time scales," Phys. Rev. Lett. **89**, 050601 (2002).

[21] W. H. Zurek, "Probabilities from entanglement, Born's rule $p_k = |\psi_k|^2$ from envariance," Phys. Rev. A **71**, 052105 (2005).

[22] D. Z. Albert, *Time and Chance* (Harvard University Press, 2000).

[23] S. Carroll, *From Eternity to Here: The Quest for the Ultimate Theory of Time* (Dutton, 2010).

[24] M. Reuter and H. Weyer, "Running Newton constant, improved gravitational actions, and galaxy rotation curves," Phys. Rev. D **70**, 124028 (2004).

[25] F. Lelli, S. S. McGaugh, and J. M. Schombert, "SPARC: Mass models for 175 disk galaxies with Spitzer photometry and accurate rotation curves," Astron. J. **152**, 157 (2016); arXiv:1606.09251.

[26] S. S. McGaugh, F. Lelli, and J. M. Schombert, "Radial acceleration relation in rotationally supported galaxies," Phys. Rev. Lett. **117**, 201101 (2016); arXiv:1609.05917.

[27] A. Ghari and H. Haghi, "Testing emergent gravity in dwarf spheroidal galaxies," (2026); arXiv:2601.01715.

[28] M. M. Brouwer *et al.*, "The weak lensing radial acceleration relation," Astron. Astrophys. **650**, A113 (2021); arXiv:2106.11677.

[29] D. Petz, "Sufficient subalgebras and the relative entropy of states of a von Neumann algebra," Commun. Math. Phys. **105**, 123 (1986).

[30] M. L. Basso, J. Maziero, and L. C. Céleri, "Quantum detailed fluctuation theorem in curved spacetime," Phys. Rev. Lett. **134**, 050406 (2025); arXiv:2405.03902.

[31] M. Dorau and R. Verch, "Kodama-like vectors in axisymmetric spacetimes," (2024); arXiv:2402.18993.

[32] D. Clowe *et al.*, "A direct empirical proof of the existence of dark matter," Astrophys. J. Lett. **648**, L109 (2006).

[33] J. F. Donoghue, "A critique of the asymptotic safety program," Front. Phys. **8**, 56 (2020).

[34] S.-K. Huang, research notes on cosmological τ (2026).

[35] B. Thattarampilly, Y. Zheng, and K. K. Kakkat, "Spherically symmetric black hole in gravity from entropy," (2026); arXiv:2602.13694.

[36] L. Herrera, "The Gibbs paradox, the Landauer principle and the irreversibility associated with tilted observers," Entropy **22**, 340 (2020).

[37] Z. Chen, Y. Song, and V. Scarani, "Petz recovery from subsystems in bosonic quantum error correction," (2025); arXiv:2511.05941.