

Information Loss is Local: The Gravitational Refractive Index as Petz Recovery Fidelity

Sheng-Kai Huang^{1,*}

¹*Independent Researcher*

(Dated: March 11, 2026)

The temporal asymmetry parameter $\tau = 1 - F \in [0, 1]$, established in Paper I from the Petz recovery framework, quantifies the arrow of time with the universal bound $F \geq e^{-\Sigma/2}$. Defining the gravitational entropy production $\Sigma_{\text{grav}} \equiv -\ln(-g_{00}) = r_s/r$, we obtain the identification $c_{\text{eff}}/c = e^{-\Sigma_{\text{grav}}/2} = F_{\text{bound}}$: the coordinate speed of light *is* the Petz recovery fidelity bound [NEW SYNTHESIS]. We then observe that $\tau(r) = 1 - \sqrt{-g_{00}(r)}$ depends only on the *local* metric component g_{00} —no knowledge of the global causal structure is needed. This locality has a radical consequence: the operationally relevant boundary where $\tau = 1$ is the *apparent horizon*, not the event horizon. Using the Kodama vector for dynamical spacetimes, we show that τ tracks the apparent horizon in Vaidya collapse and the Hubble radius in FRW cosmology—consistent with the quantum extremal surface prescription. Since $\tau < 1$ everywhere for finite Σ_{grav} , event horizons are forbidden: information that falls in must come out. The exponential metric $g_{00} = -e^{-r_s/r}$, which saturates the Petz bound, predicts gravitational-wave echoes at $\Delta t \approx 4r_s/c$ —testable with existing LIGO–Virgo data.

INTRODUCTION

In Paper I [1], we established that the Petz recovery map [3]—the unique Bayesian retrodiction functor [6]—defines a temporal asymmetry parameter $\tau \equiv 1 - F(\rho, \mathcal{R}_{\sigma, \mathcal{N}} \circ \mathcal{N}(\rho)) \in [0, 1]$, bounded by [4, 5]

$$\tau \leq 1 - e^{-\Sigma/2}, \quad (1)$$

where Σ is the entropy production of the channel $\mathcal{N}$. $\tau = 0$ for closed (unitary) evolution; $\tau > 0$ for open systems.

General relativity admits a well-known reformulation in which gravity acts as a medium with position-dependent refractive index [7, 8]: the coordinate speed of light is $c_{\text{eff}}/c = e^{\Phi/c^2}$, where $\Phi < 0$ is the Newtonian potential. In this paper we show that

$$\frac{c_{\text{eff}}}{c} = e^{\Phi/c^2} = e^{-\Sigma_{\text{grav}}/2} = F_{\text{bound}}. \quad (2)$$

Light slowing down in a gravitational field *is* information loss; the Petz recovery fidelity *is* the refractive index.

This identification has a consequence that goes beyond translation. The quantity $\tau(r) = 1 - \sqrt{-g_{00}(r)}$ is *local*: to evaluate the degree of information loss at a spacetime point, one needs only g_{00} at that point—not the global causal structure, not the future evolution of the spacetime. In general relativity, event horizons are defined teleologically: determining whether a point lies inside an event horizon requires tracing null geodesics to future null infinity, which presupposes knowledge of the spacetime’s entire future [9]. But τ makes no reference to the future. The boundary where $\tau = 1$ is set by *local* data, and that boundary is the *apparent horizon*—a quasi-local, causal surface [41, 42].

The logical chain of this paper is:

1. $c_{\text{eff}}/c = F_{\text{bound}}$ (Sec. : triple identification);

2. τ is local (Sec. : information loss does not require the future);
3. $\tau < 1$ everywhere (Sec. : no event horizon);
4. echoes at $\Delta t \approx 4r_s/c$ (Sec. : testable consequence).

THE TRIPLE IDENTIFICATION

The coordinate speed of light

[KNOWN] In the post-Newtonian approximation of GR, the coordinate speed of light in a static gravitational field is [7, 8]

$$\frac{c_{\text{eff}}}{c} = e^{\Phi/c^2}, \quad (3)$$

where $\Phi < 0$ is the Newtonian gravitational potential. This result, first anticipated by Einstein in 1911 [7] and placed on rigorous footing by Dicke [8], is exact for the exponential metric and agrees with the Schwarzschild metric through $O(r_s/r)^2$.

Gravitational entropy production

[NEW DEFINITION] We define the dimensionless gravitational entropy production as

$$\Sigma_{\text{grav}} \equiv \frac{2|\Phi|}{c^2} = \frac{r_s}{r}, \quad (4)$$

where $r_s = 2GM/c^2$ is the Schwarzschild radius. This is an *informational* entropy production—the cost of retrodiction from infinity to radius r —not a thermodynamic

entropy in the Clausius sense. The Clausius entropy production for a photon in Tolman thermal equilibrium vanishes identically [23], because $T(\Phi)\sqrt{-g_{00}} = T_\infty$. By contrast, Σ_{grav} quantifies the informational cost of climbing the gravitational potential, which persists even in equilibrium.

Three independent arguments converge on Eq. (4):

(a) *Modular flow* [KNOWN].—In the entanglement first-law regime of AQFT on Schwarzschild, the fractional quantum relative entropy (QRE) loss between radius r and infinity is $(D_{\text{in}} - D_{\text{out}})/D_{\text{in}} = r_s/r$, exactly, via the Tolman blueshift and energy redshift [27].

(b) *Gravitational Landauer principle* [KNOWN].—The Tolman-modified cost of erasing one bit at radius r exceeds that at infinity by a factor $1/\sqrt{-g_{00}}$. The dimensionless excess erasure cost yields $\Sigma_{\text{grav}} = -\ln(-g_{00})$ [28], which equals r_s/r for the exponential metric and $-\ln(1 - r_s/r) \approx r_s/r + O((r_s/r)^2)$ for Schwarzschild.

(c) *Quantum channel model* [KNOWN].—Modeling gravitational redshift as a pure-loss bosonic channel with intensity transmissivity $\eta = -g_{00}$, the QRE decrease is $\Sigma = -\ln \eta = r_s/r$ [29].

All three routes share the common mathematical core $\Sigma_{\text{grav}} = -\ln(-g_{00})$.

The channel problem

[NEW SYNTHESIS] None of the three routes above constructs a canonical gravitational CPTP map $\mathcal{N}_{\text{grav}}$ from first principles—i.e., a map with explicit Kraus operators $\{K_i\}$ such that $\Sigma = D(\rho||\sigma) - D(\mathcal{N}_{\text{grav}}(\rho)||\mathcal{N}_{\text{grav}}(\sigma)) = r_s/r$. This remains an open problem [55]. We bypass it through a combined strategy: (i) any consistent coupling of classical gravity to quantum matter is fundamentally irreversible ($\Sigma > 0$), guaranteeing the *existence* of an irreversible CPTP map [53]; (ii) the modular flow route (a) provides the coefficient r_s/r without requiring an explicit channel [27]; (iii) the pure-loss bosonic channel (c) serves as a concrete *existence proof* that a CPTP map with $\Sigma = -\ln \eta = r_s/r$ exists when $\eta = -g_{00}$; (iv) axiomatic uniqueness: any CPTP map satisfying additivity under channel composition and monotonicity under coarse-graining yields $\Sigma = -c \ln(-g_{00})$ with $c > 0$; routes (a)–(c) fix $c = 1$. The recent modular channel construction of Trejo-Calderón [54]—a CPTP map built from Gibbs-weighted singular values of the modular operator—is the most promising candidate for a first-principles $\mathcal{N}_{\text{grav}}$, though the explicit computation of ΔD in Schwarzschild remains to be carried out.

TABLE I. Gravitational entropy production $\Sigma_{\text{grav}} = r_s/r$ and Petz recovery fidelity bound $F_{\text{bound}} = e^{-\Sigma_{\text{grav}}/2}$ for representative astrophysical objects. The last row indicates the limit of the weak-field approximation.

Object	r_s/r	F_{bound}	τ_{max}
Earth surface	1.4×10^{-9}	0.99999999930	7.0×10^{-10}
Sun surface	4.2×10^{-6}	0.9999979	2.1×10^{-6}
Neutron star	0.4	0.819	0.181
Schwarzschild $r = r_s$	1	0.607	0.393

The triple identification

[NEW SYNTHESIS] Substituting Eq. (4) into the Petz fidelity bound $F \geq e^{-\Sigma/2}$ from Paper I:

$$F_{\text{bound}} = e^{-\Sigma_{\text{grav}}/2} = e^{\Phi/c^2}. \quad (5)$$

Comparing with Eq. (3), we obtain the central result:

$$\boxed{\frac{c_{\text{eff}}}{c} = e^{-\Sigma_{\text{grav}}/2} = F_{\text{bound}}}. \quad (6)$$

The effective speed of light *is* the Petz recovery fidelity bound. Equivalently, the gravitational refractive index is the inverse fidelity:

$$n_{\text{grav}}(r) = \frac{c}{c_{\text{eff}}} = e^{+\Sigma_{\text{grav}}/2} = \frac{1}{F_{\text{bound}}}. \quad (7)$$

Table I provides numerical verification across four orders of magnitude in gravitational compactness.

Caveat.—The identification (6) is exact for the exponential metric $g_{00} = -e^{-r_s/r}$ and agrees with Schwarzschild through $O(r_s/r)$. At $O((r_s/r)^2)$, the two metrics differ: the Schwarzschild expansion gives $g_{00}^{\text{Schw}} = -1 + r_s/r - r_s^2/(2r^2) + \dots$, identical to the exponential expansion through this order. The first discrepancy appears at $O((r_s/r)^3)$. In the solar system ($r_s/r \sim 10^{-6}$), this cubic correction is of order 10^{-18} —far below current measurement capability [16].

INFORMATION LOSS IS LOCAL

τ depends only on local g_{00}

[NEW SYNTHESIS] The temporal asymmetry at a space-time point p at radius r from a mass M is

$$\tau(r) = 1 - \sqrt{-g_{00}(r)}. \quad (8)$$

This is a *local, operational* quantity: an observer at r determines τ by comparing a local clock with a reference clock at infinity. No information about the future evolution of the spacetime, no knowledge of what happens at other radii, and no global causal analysis is required.

This stands in sharp contrast with the event horizon. In general relativity, the event horizon is defined as

$$\mathcal{H}^+ = \partial J^-(\mathcal{I}^+), \quad (9)$$

the boundary of the causal past of future null infinity [9]. To determine whether a point lies inside the event horizon, one must know the *entire future* of the spacetime—including whether the black hole eventually evaporates, accretes more mass, or merges with another. The event horizon is a teleological concept: it is defined by where light rays *will eventually* reach, not by what is happening *now*.

The τ framework replaces this global, teleological criterion with a local, operational one. Information loss is not a verdict delivered by the future—it is a condition determined by the present.

The apparent horizon as $\tau = 1$ surface

[NEW SYNTHESIS] In general relativity, the *apparent horizon* is defined as the outermost marginally outer trapped surface (MOTS): the surface where the expansion of outgoing null geodesics vanishes, $\theta_+ = 0$ [41]. Unlike the event horizon, the apparent horizon is quasi-local: it depends only on the metric and its first derivatives on a single time-slice.

The τ framework naturally selects this surface. For the Schwarzschild metric, $\tau = 1$ when $g_{00} = 0$, which occurs at $r = r_s$ —both the event horizon and the apparent horizon coincide. But in dynamical spacetimes they differ, and τ tracks the apparent horizon.

This is consistent with modern quantum gravity. The quantum extremal surface (QES) prescription of Penington [32] and the island formula [33] identify the QES—not the classical event horizon—as the surface that controls the entanglement entropy of Hawking radiation. The QES is the quantum-corrected analog of the apparent horizon: it extremizes the generalized entropy $S_{\text{gen}} = A/(4G\hbar) + S_{\text{bulk}}$ on a time-slice, not at future null infinity [43]. The τ framework and the QES prescription agree: the operationally relevant boundary for information loss is local.

Dynamic extension: the Kodama vector

[NEW SYNTHESIS] The static formulation of τ requires a timelike Killing vector ξ^a . The natural dynamical extension uses the Kodama vector [44]—the unique divergence-free timelike vector field in spherically symmetric spacetimes:

$$K^a = \epsilon^{ab} \nabla_b R, \quad (10)$$

where R is the areal radius and ϵ^{ab} is the Levi-Civita tensor on the (t, r) submanifold. In the static limit, K^a

TABLE II. τ_K at the event horizon versus the apparent horizon for the self-similar Vaidya spacetime $M(v) = \lambda v$. $\tau_K(\text{AH}) = 1$ exactly for all λ ; $\tau_K(\text{EH}) < 1$. See Supplemental Material for derivations.

λ	$r_{\text{EH}}/r_{\text{AH}}$	$\tau_K(r_{\text{EH}})$	$1 - \tau_K(r_{\text{EH}})$
0.001	1.004	0.937	6.3%
0.01	1.044	0.796	20.4%
0.03	1.162	0.627	37.3%
0.05	1.382	0.474	52.6%
1/16	2.000	0.293	70.7%

reduces to the Killing vector ξ^a , and $|K| = \sqrt{-g_{00}}$. The Kodama recovery fidelity is

$$F_{\text{bound}}^K(A \rightarrow B) = \frac{|K|_A}{|K|_B}, \quad (11)$$

recovering all results of Sec. in the static limit.

Vaidya collapse.—For the ingoing Vaidya metric (mass accretion $M(v)$), $|K| = \sqrt{1 - 2M(v)/r}$ [41, 45], giving

$$\tau_K(r, v) = 1 - \sqrt{1 - \frac{2M(v)}{r}}. \quad (12)$$

This reaches $\tau = 1$ when $r = 2M(v)$ —precisely the *apparent* horizon. During gravitational collapse, $M(v)$ increases and the $\tau = 1$ surface expands outward, tracking the growing apparent horizon. During Hawking evaporation, $M(v)$ decreases and the $\tau = 1$ surface shrinks—information recovery improves, consistent with the Page curve.

Explicit demonstration: $\text{AH} \neq \text{EH}$.—The event horizon is the outermost outgoing null geodesic that fails to escape to infinity. For radial null rays in Vaidya, $dr/dv = \frac{1}{2}(1 - 2M(v)/r)$, integrated backward from $r_{\text{EH}} = 2M_0$ at $v \rightarrow \infty$. For the self-similar profile $M(v) = \lambda v$, the null geodesic equation admits the exact solution $r_{\text{EH}}(v) = C_- v$, where $C_- = (1 - \sqrt{1 - 16\lambda})/4$, while $r_{\text{AH}}(v) = 2\lambda v$. Since $C_- > 2\lambda$ for $\lambda > 0$ (see Supplemental Material [62] for proof), $r_{\text{EH}} > r_{\text{AH}}$ during accretion: the event horizon lies *outside* the apparent horizon.

Table II quantifies the discrepancy. At the event horizon, $\tau_K < 1$ —the τ framework says the EH is *not* a surface of complete information loss. At the apparent horizon, $\tau_K = 1$ exactly, for all v . During evaporation ($dM/dv < 0$), the situation reverses: $r_{\text{EH}} < r_{\text{AH}}$, and the Kodama vector becomes spacelike at the EH. In both cases, $\tau_K = 1$ at the AH—never at the EH.

For a step-function collapse ($M(v) = M_0 \theta(v)$), the event horizon originates at $r = 0$ when $v = -4M_0$ —in the *flat* region, before the shell arrives. The EH “anticipates” the collapse; the AH appears only at $v = 0$. This is the starkest illustration: τ_K is zero in the flat region

(no information loss), even though the event horizon is already present.

FRW cosmology.—In a spatially flat FRW universe with Hubble parameter H ,

$$\tau_K(R) = 1 - \sqrt{1 - H^2 R^2}, \quad (13)$$

reaching $\tau = 1$ at $R = 1/H$, the Hubble radius—which is the apparent horizon of FRW, not an event horizon [45].

Verification proposal: SYK model

[SPECULATIVE] The SYK model [49, 50] is a solvable quantum-mechanical model with a holographic dual to nearly-AdS₂ (JT) gravity, exhibiting both scrambling and black hole formation. It provides a setting where both sides of the τ equation—the QI recovery fidelity $F(t)$ and the gravitational entropy production $\Sigma_{\text{grav}}(t)$ —can be computed independently.

We propose the following test. In the SYK model with time-dependent coupling (modeling dynamical black hole formation and evaporation): (i) compute the Petz recovery fidelity $F(t)$ for an early-time operator from late-time data; (ii) compute $\Sigma_{\text{grav}}(t) = -\ln(-g_{00}^{\text{bulk}}(t))$ from the dual JT gravity; (iii) compare whether $F(t)$ tracks the apparent horizon or the event horizon in the bulk. The τ framework predicts that $F(t)$ tracks the apparent horizon. This calculation is feasible with exact diagonalization at moderate N [50].

NO EVENT HORIZON

Model-independent no-horizon theorem

[NEW SYNTHESIS] The locality of τ established in Sec. has a direct physical consequence. If gravitational redshift is a quantum channel with *finite* entropy production $\Sigma < \infty$, then the Petz bound gives $F \geq e^{-\Sigma/2} > 0$, so $\tau < 1$. An event horizon requires $F = 0$ (complete information loss), which demands $\Sigma \rightarrow \infty$ —impossible for a finite-mass object at finite distance. The Petz bound forbids event horizons model-independently [5]; neither the specific form of Σ_{grav} nor the saturation hypothesis is needed.

In the language of Sec. : because τ is local and finite, information loss is never complete. What falls in must come out.

The exponential metric

[NEW SYNTHESIS] If the Petz bound is taken as an identity— $\sqrt{-g_{00}} = e^{-\Sigma_{\text{grav}}/2}$ —the metric component is

uniquely determined. This gives the exponential metric [10, 11]:

$$ds^2 = -e^{-r_s/r} c^2 dt^2 + e^{+r_s/r} (dr^2 + r^2 d\Omega^2), \quad (14)$$

in isotropic coordinates. Since $e^{-r_s/r} > 0$ for all $r > 0$, $\tau < 1$ everywhere.

Exact solution.—The exponential metric is not a vacuum solution of the Einstein equations ($R_{\mu\nu} \neq 0$). It is an exact solution of Einstein gravity coupled to a phantom scalar field $\phi = M/r$ [17], and also arises in logarithmic $f(R)$ gravity [18] and scalar-Einstein-Gauss-Bonnet theory [19]. Makukov and Mychelkin [12] showed that three independent scalar-field families within GR all reduce to the exponential metric when the scalar charge equals the gravitational mass. Boonserm et al. [13] classified the geometry as a traversable wormhole.

The spatial metric is assumed, not derived.—The information-theoretic framework constrains only the temporal component g_{00} through $\sqrt{-g_{00}} = e^{-\Sigma_{\text{grav}}/2}$. The spatial component $g_{rr} = e^{+r_s/r}$ in Eq. (14) follows from the isometry condition $g_{00} \cdot g_{rr} = -1$ (isotropic coordinates), which is equivalent to Dicke’s scalar refractive index [8]: $n^2 = 1/(-g_{00})$ with $g_{rr} = n^2$. This is a physically motivated assumption—not a derivation from quantum information theory. The fundamental reason is that quantum channels model *temporal* processes (evolution, decoherence, loss), and the Petz recovery framework naturally constrains the temporal metric component; no analogous information-degradation process exists for the spatial direction. Consequently, we classify our predictions as *robust* (depending only on g_{00} : no-horizon theorem, refractive index, gravitational decoherence rate) or *conditional* (depending on the full metric: shadow size, ISCO radius, QNM frequencies, echo delay). The Jacobson [40] and Dorau–Much [27] programs *do* derive all metric components from information-theoretic premises, but their vacuum solution is Schwarzschild, not exponential. This tension remains an open question.

Comparison with Schwarzschild.—A persistent misconception holds that the exponential metric “predicts half the observed light bending.” This conflates the Yilmaz *field equations*—an alternative theory of gravity shown to be inconsistent by Misner [14]—with the exponential *metric* as a solution of standard GR coupled to a phantom scalar. Particles and photons follow standard geodesics of (14); only the geometry differs from Schwarzschild.

The PPN expansion in isotropic coordinates gives

$$g_{00}^{\text{exp}} = -e^{-r_s/r} = -(1 - 2U + 2U^2 - \dots), \quad (15)$$

$$g_{ij}^{\text{exp}} = e^{+r_s/r} \delta_{ij} = (1 + 2U + 2U^2 + \dots) \delta_{ij}, \quad (16)$$

where $U = r_s/(2r)$ is the Newtonian potential. Matching with the standard PPN form $g_{00} = -(1 - 2U + 2\beta U^2 + \dots)$, $g_{ij} = (1 + 2\gamma U + \dots) \delta_{ij}$ yields $\gamma = \beta = 1$ —*identical to GR*. Mychelkin et al. [15] verified explicitly

that the perihelion precession at first post-Newtonian order matches Schwarzschild: $\Delta\phi_{\text{exp}} = \Delta\phi_{\text{Schw}} = 6\pi M/p$. The first difference appears at 2PN, where the precession differs by $\sim 2.3 \times 10^{-7}$ arcsec/century for Mercury—four orders of magnitude below current precision. All solar-system tests are passed: light bending, Shapiro delay, perihelion precession, and the Cassini bound $|\gamma - 1| < 2.3 \times 10^{-5}$ [16]. *The exponential metric is observationally degenerate with Schwarzschild for every current and foreseeable weak-field test.*

Caveat on saturation.—The JRSWW bound [5] is structurally strict for $\Sigma > 0$: the proof via the Golden-Thompson inequality is an equality only when the relevant operators commute—the $\Sigma = 0$ case [3]. No known quantum channel saturates the bound for $\Sigma > 0$. The exponential metric should be understood as the geometry defined by the *identity* $\sqrt{-g_{00}} = e^{-\Sigma_{\text{grav}}/2}$; whether nature saturates this bound or admits a gap is empirical.

STRONG-FIELD PREDICTIONS

[CONDITIONAL] In the strong-field regime, the exponential and Schwarzschild metrics diverge, yielding three testable predictions. All depend on the spatial metric assumption $g_{rr} = e^{+r_s/r}$.

(1) *Shadow size.*—The exponential photon sphere at $r_{\text{ph}} = r_s$ (isotropic coordinates) yields a critical impact parameter $b_{\text{crit}} = 2em \approx 5.44m$ versus Schwarzschild’s $3\sqrt{3}m \approx 5.20m$ ($m = GM/c^2$): a 4.6% larger shadow. The EHT measurement of $42 \pm 3 \mu\text{as}$ for M87* [46] is consistent with both metrics.

(2) *Quasinormal modes.*—In the eikonal limit, $f_{\text{QNM}}^{\text{exp}}/f_{\text{QNM}}^{\text{Schw}} = 3\sqrt{3}/(2e) \approx 0.956$, a $\sim 4.4\%$ downward shift. WKB calculations find a -7.8% shift for a generalized exponential wormhole [18]. Next-generation gravitational-wave detectors (Einstein Telescope, Cosmic Explorer) will reach $\sim 1\%$ precision.

(3) *Gravitational-wave echoes.*—The most distinctive signature. The traversable wormhole structure produces echoes—repeated, damped pulses following the initial ringdown—with time delay

$$\Delta t_{\text{echo}} \approx \frac{4.2 r_s}{c}, \quad (17)$$

giving ~ 2.5 ms for a $60 M_{\odot}$ remnant. This is $\sim 90\times$ shorter than Planck-wall models [48], because the exponential metric has no horizon and its tortoise coordinate $r^* = \int e^{r_s/r} dr$ remains bounded. Current echo searches use ~ 200 ms templates; dedicated short-delay searches in existing LIGO–Virgo O3/O4 data are needed.

The echo prediction follows directly from the no-horizon theorem: since $\tau < 1$ everywhere, the gravitational potential forms a well, not a one-way membrane. Gravitational waves enter, scatter off the worm-

hole throat, and return—carrying the signature of incomplete information loss.

Honest assessment.—Since both metrics share $\gamma = \beta = 1$, weak-field data cannot distinguish them; the first differences are inherently *strong-field* effects. If nature selects $F > e^{-\Sigma_{\text{grav}}/2}$, the actual metric lies between the exponential and Schwarzschild predictions. We emphasize a two-layer structure: the *robust* predictions (no-horizon theorem, $n_{\text{grav}} = 1/F_{\text{bound}}$, decoherence bound) depend only on g_{00} and hold regardless of the spatial metric; the *conditional* predictions (shadow, QNM, echoes) depend on the isometry assumption $g_{rr} = e^{+r_s/r}$. The most distinctive signature—short-delay echoes (17)—has not yet been searched for with appropriate templates in LIGO–Virgo data.

SUPPORTING CONNECTIONS

The τ framework connects to several established results in gravitational physics. We classify each connection in Table III.

Gravitational decoherence.—[NEW SYNTHESIS] Pikovski et al. [22] showed that gravitational time dilation universally dephases composite quantum systems. In the τ framework [2],

$$\tau_{\text{grav}}(t) = 1 - \exp(-N\Gamma t^2), \quad (18)$$

where $\Gamma = (\Delta E)^2(\Delta\Phi)^2/(2\hbar^2 c^4)$. The dephasing rate is controlled by $\Delta\Phi/c^2 \sim \Sigma_{\text{grav}}/2$: the gravitational entropy production governs both the static (refractive index) and dynamic (decoherence) aspects of information loss.

Unruh effect.—[NEW SYNTHESIS] A uniformly accelerating observer at acceleration a perceives a thermal bath at $T_U = \hbar a/(2\pi c k_B)$, inducing entropy production $\Sigma_{\text{Unruh}} = 2\pi a \Delta x/(\hbar c^2)$ on any carried system [24, 25]. By the equivalence principle, this connects to Σ_{grav} .

Bekenstein bound.—[NEW SYNTHESIS] The bound $S \leq 2\pi ER/(\hbar c)$ [26] translates to $\tau \leq 1 - \exp(-\pi ER/(\hbar c))$, providing a universal upper limit on temporal asymmetry for spatially bounded systems.

Complementarity.—[NEW SYNTHESIS] The infalling observer experiences $\tau_{\text{in}} \approx 0$ (locally flat), while the external observer describes $\tau_{\text{ext}} \rightarrow 1$ [30]. Different observer foliations assign different τ values—complementarity is observer-dependent τ .

Page curve.—[NEW SYNTHESIS] The Page curve [31] translates to $\tau(t) \leq 1 - e^{-S_{\text{Page}}(t)/2}$: τ grows before the Page time (information loss worsens) and shrinks after it (recovery improves) [32, 33].

Hayden–Preskill.—[KNOWN] Cotler et al. [35] proved that the HP decoder [34] is the Petz recovery map, with fidelity satisfying the same bound (1).

TABLE III. Classification of results.

Connection	Status
$c_{\text{eff}}/c = F_{\text{bound}}$ (refractive index)	NEW SYNTHESIS
$\Sigma_{\text{grav}} = -\ln(-g_{00})$ (entropy production)	NEW DEFINITION
τ is local $\rightarrow$ apparent horizon	NEW SYNTHESIS
Kodama extension to dynamic spacetimes	NEW SYNTHESIS
No-horizon theorem	NEW SYNTHESIS
Observer-dependent τ (complementarity)	NEW SYNTHESIS
Page curve as $\tau(t)$	NEW SYNTHESIS
Gravitational decoherence as Petz recovery	NEW SYNTHESIS
HP decoder = Petz map [35]	KNOWN
Entanglement wedge via Petz [37]	KNOWN
$F^2 \geq e^{-\Delta S_{\text{gen}}}$ [32]	KNOWN
SYK verification of τ tracking AH	SPECULATIVE

Entanglement wedge.—[KNOWN] In AdS/CFT, bulk reconstruction via the Petz map [36, 37] satisfies $F^2 \geq e^{-\Delta S_{\text{gen}}}$ [35]—the same universal recovery inequality.

WHAT τ CAN AND CANNOT EXPLAIN

Can:

- Identify information loss as a local, operational quantity that does not require knowledge of the future.
- Identify the apparent horizon—not the event horizon—as the operationally relevant boundary, consistent with the QES prescription.
- Unify quantum error correction, thermodynamic irreversibility, black hole information, and gravitational decoherence under a single parameter τ .
- Provide universal bounds: any gravitational channel satisfies $F \geq e^{-\Sigma_{\text{grav}}/2}$, constraining recovery independently of the specific protocol.
- Derive the no-horizon theorem: $\tau < 1$ for any finite-mass object at finite distance (model-independent [5]).
- Extend to dynamical spacetimes via the Kodama vector, tracking apparent horizons in Vaidya and FRW.

Cannot:

- Explain dark matter: at galactic scales, $\Sigma_{\text{grav}} \sim 10^{-6}$, far too small to produce observable effects beyond Newtonian gravity.
- Explain dark energy or the cosmological constant.

- Derive g_{ab} from τ alone: the relation is one-directional ($g \rightarrow \Sigma \rightarrow \tau$); τ constrains only g_{00} . The spatial component g_{rr} follows from the Dicke isometry condition; programs that derive all components from entanglement equilibrium (Jacobson, Dorau–Much) yield Schwarzschild as the *vacuum* solution. The exponential metric requires a phantom scalar source, whose information-theoretic origin remains open.
- Construct a canonical gravitational CPTP map: the bypass strategy (Sec.) establishes $\Sigma_{\text{grav}} = r_s/r$ without an explicit channel, but the Kraus operator representation remains open [54, 55].
- Resolve the firewall paradox [38]: while $\tau < 1$ removes the event horizon premise, the framework does not address the interior structure of compact objects.

DISCUSSION AND OUTLOOK

The central result of this paper is that gravitational information loss is *local*. The temporal asymmetry $\tau = 1 - \sqrt{-g_{00}}$ depends only on the metric component at a single point. The operationally relevant boundary for information loss is therefore the apparent horizon—a quasi-local, causal surface [42]—not the event horizon. This is consistent with the quantum extremal surface prescription [32, 43] and the crossed-product algebra approach [58, 59], which makes gravitational entropy well-defined and observer-dependent [60].

The identification $c_{\text{eff}}/c = F_{\text{bound}}$ (Eq. (6)) is complementary to the Jacobson program [40]: while Jacobson derives *field equations* from thermodynamic entropy, we derive the *metric* directly from information recoverability. The Dorau–Much result [27]—deriving the semiclassical Einstein equations from quantum relative entropy—suggests that the two programs share a common informational foundation. Cirafofi [56] has established fluctuation theorems in gravitational algebras, confirming that the Crooks–Petz connection extends to de Sitter spacetime.

Several directions remain. *(i) The canonical channel:* the modular channel of Trejo-Calderón [54] is the most promising candidate; computing its QRE loss in Schwarzschild would either confirm $\Sigma = r_s/r$ from first principles or reveal a correction. *(ii) Rotation:* the Kerr analog of the exponential metric [20, 21] lacks both horizons and ergospheres; extending the framework beyond spherical symmetry requires the axisymmetric Kodama-type vector of Dorau and Verch [57]. *(iii) Entanglement-assisted recovery and $\tau < 0$:* [SPECULATIVE] when the gravitational channel is supplemented by an entanglement resource (e.g., a traversable wormhole via

the Gao–Jafferis–Wall protocol [51]), the effective entropy production becomes $\Sigma_{\text{eff}} = \Sigma_{\text{grav}} - I(L; R)$. If $I(L; R) > \Sigma_{\text{grav}}$, the recovery fidelity exceeds the geometric bound— $\tau_{\text{signed}} < 0$ —unifying the Hayden–Preskill protocol, traversable wormholes, and the Page curve transition into a single continuous parameter. This prediction is testable via SYK numerics and quantum circuit experiments [52]. *(iv) Non-Markovian gravity*: if the gravitational channel exhibits memory effects, Σ_{grav} may not be additive and the Petz bound requires modification [24]. This could serve as a signature of quantum gravity: classical gravity is necessarily Markovian, while quantum gravity may permit non-Markovian information backflow.

In Paper 1b [2], we applied the τ framework to wavefunction collapse. A future paper (Paper 3) will investigate the weak-field limit at galactic scales.

The author thanks the quantum information, quantum gravity, and quantum foundations communities for the theoretical foundations on which this work builds.

* akai@fawstudio.com

- [1] S.-K. Huang, “Universal Recovery: The Petz Map as Retrodiction Functor, Error Corrector, and Entropy Bound,” companion paper (2026).
- [2] S.-K. Huang, “Wavefunction Collapse as Retrodiction Failure: Gravitational Decoherence, Objectivity, and the τ - R Bridge,” companion paper (2026).
- [3] D. Petz, “Sufficiency of channels over von Neumann algebras,” *Q. J. Math.* **39**, 97 (1988).
- [4] O. Fawzi and R. Renner, “Quantum Conditional Mutual Information and Approximate Markov Chains,” *Commun. Math. Phys.* **340**, 575 (2015).
- [5] M. Junge, R. Renner, D. Sutter, M. M. Wilde, and A. Winter, “Universal Recovery Maps and Approximate Sufficiency of Quantum Relative Entropy,” *Ann. Henri Poincaré* **19**, 2955 (2018).
- [6] A. J. Parzygnat and F. Buscemi, “Axioms for retrodiction: Achieving time-reversal symmetry with a prior,” *Quantum* **7**, 1013 (2023).
- [7] A. Einstein, “Über den Einfluß der Schwerkraft auf die Ausbreitung des Lichtes,” *Ann. Phys. (Leipzig)* **35**, 898 (1911).
- [8] R. H. Dicke, “Gravitation without a Principle of Equivalence,” *Rev. Mod. Phys.* **29**, 363 (1957).
- [9] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, 1973).
- [10] A. Papapetrou, “Eine Theorie des Gravitationsfeldes mit einer Feldfunktion,” *Z. Phys.* **139**, 518 (1954).
- [11] H. Yilmaz, “New Approach to General Relativity,” *Phys. Rev.* **111**, 1417 (1958).
- [12] M. A. Makukov and E. G. Mychelkin, “Triple Path to the Exponential Metric,” *Found. Phys.* **50**, 1346 (2020).
- [13] P. Boonserm, T. Ngampitipan, A. Simpson, and M. Visser, “Exponential metric represents a traversable wormhole,” *Phys. Rev. D* **98**, 084048 (2018).
- [14] C. W. Misner, “Yilmaz Cancels Newton,” *Nuovo Cimento B* **114**, 1079 (1999); arXiv:gr-qc/9504050.
- [15] E. G. Mychelkin, M. A. Makukov, and G. Suliyeva, “On the weak and strong field effects in antiscalar background,” *Gen. Relativ. Gravit.* **56**, 44 (2024); arXiv:2403.11610.
- [16] B. Bertotti, L. Iess, and P. Tortora, “A test of general relativity using radio links with the Cassini spacecraft,” *Nature* **425**, 374 (2003).
- [17] K. K. Ernazarov and V. D. Ivashchuk, “Phantom scalar field solution in the exponential metric,” (2026).
- [18] S. Nath and D. Sarma, “Exponential wormhole solution in logarithmic $f(R)$ gravity,” (2025).
- [19] K. K. Ernazarov, “The Yilmaz-Rosen and Janis-Newman-Winicour metric solutions in the scalar-Einstein-Gauss-Bonnet 4d gravitational model,” arXiv:2510.21625 (2025).
- [20] J. R. Graber, “Rotating Yilmaz metric,” arXiv:1708.05645 (2017).
- [21] M. A. Makukov and E. G. Mychelkin, “Rotating exponential metric via Newman–Janis algorithm,” arXiv:2301.08118 (2023).
- [22] I. Pikovski, M. Zych, F. Costa, and Č. Brukner, “Universal decoherence due to gravitational time dilation,” *Nat. Phys.* **11**, 668 (2015).
- [23] R. C. Tolman, “On the Weight of Heat and Thermal Equilibrium in General Relativity,” *Phys. Rev.* **35**, 904 (1930).
- [24] G. T. Landi and M. Paternostro, “Irreversible entropy production: From classical to quantum,” *Rev. Mod. Phys.* **93**, 035008 (2021).
- [25] W. G. Unruh, “Notes on black-hole evaporation,” *Phys. Rev. D* **14**, 870 (1976).
- [26] J. D. Bekenstein, “Universal upper bound on the entropy-to-energy ratio for bounded systems,” *Phys. Rev. D* **23**, 287 (1981).
- [27] L. Dorau and A. Much, “From Quantum Relative Entropy to the Semiclassical Einstein Equations,” *Phys. Rev. Lett.* (2025), arXiv:2510.24491.
- [28] L. Herrera, “Landauer Principle and General Relativity,” *Entropy* **22**, 340 (2020).
- [29] M. Ahmadi, D. E. Bruschi, C. Sabin, G. Adesso, and I. Fuentes, “Relativistic Quantum Metrology,” *Sci. Rep.* **4**, 4996 (2014).
- [30] L. Susskind, L. Thorlacius, and J. Uglum, “The Stretched Horizon and Black Hole Complementarity,” *Phys. Rev. D* **48**, 3743 (1993).
- [31] D. N. Page, “Average entropy of a subsystem,” *Phys. Rev. Lett.* **71**, 1291 (1993).
- [32] G. Penington, “Entanglement Wedge Reconstruction and the Information Problem,” *J. High Energy Phys.* **2020**(09), 002 (2020).
- [33] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian, and A. Tajdini, “The entropy of Hawking radiation,” *Rev. Mod. Phys.* **93**, 035002 (2021).
- [34] P. Hayden and J. Preskill, “Black holes as mirrors: quantum information in random subsystems,” *J. High Energy Phys.* **2007**(09), 120 (2007).
- [35] J. Cotler, P. Hayden, G. Penington, G. Salton, B. Swingle, and M. Walter, “Entanglement Wedge Reconstruction via Universal Recovery Channels,” *Phys. Rev. X* **9**, 031011 (2019).
- [36] A. Almheiri, X. Dong, and D. Harlow, “Bulk Locality and Quantum Error Correction in AdS/CFT,” *J. High Energy Phys.* **2015**(04), 163 (2015).

- [37] H. Chen, G. Penington, and G. Salton, “Entanglement Wedge Reconstruction using the Petz Map,” *J. High Energy Phys.* **2020**(01), 168 (2020).
- [38] A. Almheiri, D. Marolf, J. Polchinski, and J. Sully, “Black Holes: Complementarity vs. Firewalls,” *J. High Energy Phys.* **2013**(02), 062 (2013).
- [39] J. Maldacena and L. Susskind, “Cool horizons for entangled black holes,” *Fortschr. Phys.* **61**, 781 (2013).
- [40] T. Jacobson, “Thermodynamics of Spacetime: The Einstein Equation of State,” *Phys. Rev. Lett.* **75**, 1260 (1995).
- [41] S. A. Hayward, “General laws of black-hole dynamics,” *Phys. Rev. D* **49**, 6467 (1994).
- [42] A. Ashtekar and B. Krishnan, “Isolated and dynamical horizons and their applications,” *Living Rev. Relativ.* **7**, 10 (2004).
- [43] N. Engelhardt and A. C. Wall, “Quantum extremal surfaces: holographic entanglement entropy beyond the classical regime,” *J. High Energy Phys.* **2015**(01), 073 (2015).
- [44] H. Kodama, “Conserved energy flux for the spherically symmetric system and the backreaction problem in the black hole evaporation,” *Prog. Theor. Phys.* **63**, 1217 (1980).
- [45] G. Abreu and M. Visser, “Kodama time: geometrically preferred foliations of GR spacetimes,” *Phys. Rev. D* **82**, 044027 (2010).
- [46] Event Horizon Telescope Collaboration, “First M87 Event Horizon Telescope Results. I,” *Astrophys. J. Lett.* **875**, L1 (2019).
- [47] M. C. Miller *et al.*, “PSR J0030+0451 Mass and Radius from NICER Data,” *Astrophys. J. Lett.* **887**, L24 (2019).
- [48] V. Cardoso, E. Franzato, and P. Pani, “Is the Gravitational-Wave Ringdown a Probe of the Event Horizon?,” *Phys. Rev. Lett.* **116**, 171101 (2016).
- [49] S. Sachdev and J. Ye, “Gapless spin-fluid ground state in a random quantum Heisenberg magnet,” *Phys. Rev. Lett.* **70**, 3339 (1993).
- [50] J. Maldacena and D. Stanford, “Remarks on the Sachdev-Ye-Kitaev model,” *Phys. Rev. D* **94**, 106002 (2016).
- [51] P. Gao, D. L. Jafferis, and A. C. Wall, “Traversable wormholes via a double trace deformation,” *J. High Energy Phys.* **2017**(12), 151 (2017).
- [52] D. Jafferis, A. Zlokapa, J. D. Lykken, D. K. Kolchmeyer, S. I. Davis, N. Lauk, H. Neven, and M. Spiropulu, “Traversable wormhole dynamics on a quantum processor,” *Nature* **612**, 51 (2022).
- [53] T. D. Galley, F. Giacomini, and J. O. E. Selby, “Any consistent coupling between classical gravity and quantum matter is fundamentally irreversible,” *Quantum* **7**, 1142 (2023).
- [54] J. A. Trejo-Calderón, “Modular Channels, Thermal Filtering and the Spectral Emergence of Spacetime,” *arXiv:2504.20457* (2025).
- [55] M. Kasprzak and E. Tjoa, “Transmission of quantum information through quantum fields in curved spacetimes,” *arXiv:2408.00518* (2024).
- [56] M. Cirafici, “Fluctuation theorems, quantum channels and gravitational algebras,” *arXiv:2408.04219* (2024).
- [57] L. Dorau and R. Verch, “Modular flow and entropy for axisymmetric spacetimes,” *arXiv:2402.18993* (2024).
- [58] E. Witten, “Gravity and the crossed product,” *J. High Energy Phys.* **2022**(10), 008 (2022).
- [59] V. Chandrasekaran, R. Longo, G. Penington, and E. Witten, “An algebra of observables for de Sitter space,” *J. High Energy Phys.* **2023**(02), 082 (2023).
- [60] J. Kudler-Flam, S. Leutheusser, and G. Satishchandran, “Generalized black hole entropy is von Neumann entropy,” *arXiv:2309.15897* (2025).
- [61] S. V. Moreira and L. C. Celeri, “Decoherence and entropy production due to quantum fluctuations of spacetime,” *arXiv:2603.02034* (2026).
- [62] See Supplemental Material for detailed Kodama vector derivation, event horizon calculations in Vaidya spacetime, QES connection, and echo delay derivation.
- [63] W. A. Hiscock, “Models of evaporating black holes,” *Phys. Rev. D* **23**, 2823 (1981).