

Warp Field Theory in the Σ Framework: From No-Free-Warp Theorem to Subluminal Bubble Engineering

Sheng-Kai Huang^{1,*}

¹*Independent Researcher*

(Dated: March 29, 2026)

We develop a complete theory of warp drive physics using the information field $\Sigma = -\ln(-g_{00})$. We prove a *No-Free-Warp Theorem*: no warp bubble satisfying asymptotic flatness and a non-trivial interior can have $\nabla^2 \Sigma = 0$ everywhere, so every bubble costs strictly positive energy. The optimal wall shape minimizing Fisher information is the harmonic profile $h(r) = A + B/r$, which reduces energy by 33% compared to the Alcubierre sigmoid. We derive the master energy formula $E_{\text{warp}} = c^4 \Sigma_0^2 \mathcal{G}/(8G)$, where $\mathcal{G}$ depends only on bubble geometry, and show that $E_{\text{warp}} \propto v_s^4/c^4$ makes subluminal warp exponentially cheaper than superluminal. A 1-meter bubble at $v_s = 44$ m/s requires ~ 1 kg of mass-energy—extraordinary but finite. We prove that all known warp metrics (Alcubierre, Natário, Lentz) share the same Σ dependence on the shape function; the exponential warp metric is the unique horizon-free completion. We analyze Casimir energy as a source, finding a tentative sweet spot for bacterium-scale bubbles at walking speed (~ 10 kJ), though the required geometry is non-trivial. We propose a 4-phase experimental roadmap from current Casimir experiments to a photon-scale warp prototype. Superluminal warp requires null energy condition (NEC) violation. For subluminal warp, the situation is metric-dependent: the Alcubierre shift-vector construction can satisfy the NEC [?], while the isotropic exponential metric requires NEC violation even at subluminal speeds (a structural consequence of $G^r_r = -e^{-\Sigma} \Sigma'^2/4 < 0$). This distinction—between shift-vector and lapse-function approaches—is a key open question.

I. INTRODUCTION

In 1994, Alcubierre demonstrated that general relativity admits a spacetime geometry in which a “warp bubble” transports a flat-space interior at arbitrary coordinate velocity [1]. The line element is

$$ds^2 = -c^2 dt^2 + (dx - v_s f(r_s) dt)^2 + dy^2 + dz^2, \quad (1)$$

where $v_s(t)$ is the velocity of the bubble center, $r_s = [(x - x_s(t))^2 + y^2 + z^2]^{1/2}$ is the distance from the center, and $f(r_s)$ is a shape function with $f(0) = 1$ and $f \rightarrow 0$ at infinity.

This geometry requires violation of the null energy condition (NEC) [1, 2], with early estimates placing the total negative energy at $\sim M_\odot c^2 \sim 10^{47}$ J [2]. Subsequent work has progressively clarified the situation: Van Den Broeck’s volume reduction [3], Natário’s zero-expansion drive [4], Bobrick and Martire’s classification framework [5], Lentz’s positive-energy soliton (whose validity has been debated [7, 8]), and White’s Casimir nanostructure proposal [9].

Our contribution is to reformulate warp physics entirely in terms of the information field $\Sigma \equiv -\ln(-g_{00})$, introduced in our companion papers [20–22]. The Σ framework unifies all warp metrics through a single scalar field, proves sharp optimality theorems for bubble geometry, and reveals that subluminal warp at modest speeds is exponentially cheaper than superluminal warp (energy $\propto v^4/c^4$). Our numerical simulation of the isotropic exponential metric shows that NEC violation persists even

at subluminal speeds ($G^r_r < 0$ structurally); however, this is specific to the lapse-only approach and does not apply to shift-vector constructions [?].

We adopt the convention that c appears explicitly; G is Newton’s constant, $\hbar$ is the reduced Planck constant.

II. WARP DRIVES IN Σ LANGUAGE

A. The Alcubierre metric

Expanding Eq. (1), the lapse-squared component reads

$$g_{00} = -(1 - v_s^2 f^2/c^2), \quad (2)$$

giving the information field

$$\Sigma_{\text{Alc}} = -\ln(1 - v_s^2 f^2/c^2). \quad (3)$$

Three regimes emerge:

(i) Subluminal interior ($v_s f < c$): $\Sigma > 0$ and finite. At the bubble center ($f = 1$), $\Sigma = \ln \gamma^2$ with $\gamma = (1 - v_s^2/c^2)^{-1/2}$.

(ii) Bubble wall ($v_s f \rightarrow c$): $\Sigma \rightarrow +\infty$. This divergence is *structurally identical* to a Schwarzschild horizon, where $g_{00} = -(1 - r_s^2/r^2) \rightarrow 0$.

(iii) Superluminal region ($v_s f > c$): $g_{00} > 0$ —a signature change. Σ becomes complex, signaling a causally pathological region.

B. The exponential warp metric

In the companion paper [21], we showed that the exponential metric $g_{00} = -e^{-\Sigma}$ arises from saturating the

* akai@fawstudio.com

Petz recovery bound and never permits $g_{00} = 0$. Applied to the warp bubble:

$$g_{00}^{(\text{exp})} = -\exp(-v_s^2 f^2 / c^2), \quad (4)$$

so that

$$\Sigma_{\text{exp}} = v_s^2 f^2 / c^2. \quad (5)$$

For any finite v_s and $0 \leq f \leq 1$, we have $0 \leq \Sigma_{\text{exp}} \leq v_s^2 / c^2$, which is *everywhere finite*. Even at $v_s = c$, $f = 1$: $g_{00}^{(\text{exp})} = -e^{-1} \approx -0.368$ —no horizon forms.

In ADM decomposition, this corresponds to a modified lapse

$$N_{\text{exp}}^2 = \exp(-v_s^2 f^2 / c^2) + v_s^2 f^2 / c^2, \quad (6)$$

which is finite and nondegenerate for all v_s .

C. Universality of the Σ profile

Proposition 1 (Warp metric universality). *All warp metrics with a given shape function $f(r_s)$ produce the same Σ profile up to the choice of $g_{00}(\Sigma)$. Specifically, the Alcubierre, Natário, and Lentz metrics differ in the shift vector β^i but share the same functional dependence $\Sigma = \Sigma(f)$.*

Proof. For any ADM metric with shift β^i and spatial metric $\gamma_{ij} = \delta_{ij}$, the effective g_{00} is $g_{00} = -N^2 + \gamma_{ij}\beta^i\beta^j = -N^2 + |\beta|^2$. In the Alcubierre class, $|\beta|^2 = v_s^2 f^2$ and $N = 1$, so $g_{00} = -(1 - v_s^2 f^2 / c^2)$ regardless of the *direction* of β^i . The Natário metric modifies β^i to achieve zero volume expansion but with the same $|\beta|^2$ dependence on f . Lentz’s shift-only soliton likewise enters Σ only through $|\beta|^2$. The exponential metric is the unique completion that makes Σ finite for all f . $\square$

III. NO-FREE-WARP THEOREM

Theorem 1 (No-Free-Warp). *Let $\Sigma : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a smooth function satisfying:*

- (i) $\Sigma = \Sigma_0 > 0$ in a bounded region Ω (the bubble interior);
- (ii) $\Sigma = 0$ outside a larger bounded region $\Omega' \supset \Omega$ (asymptotic flatness).

Then $\nabla^2 \Sigma \not\equiv 0$, and the Fisher information

$$\mathcal{F}[\Sigma] = \int_{\mathbb{R}^3} |\nabla \Sigma|^2 d^3x > 0. \quad (7)$$

Proof. Suppose for contradiction that $\nabla^2 \Sigma = 0$ throughout $\mathbb{R}^3$. Then Σ is a harmonic function. By the strong maximum principle for harmonic functions [23], if Σ attains its supremum Σ_0 at any interior point of a connected open set, then Σ is constant on that set. Since

$\Sigma = \Sigma_0$ in Ω and $\Sigma = 0$ on $\partial\Omega'$, the connected component of $\{\Sigma \leq \Sigma_0\}$ containing the boundary forces $\Sigma = \Sigma_0$ everywhere in the connected domain—but this contradicts $\Sigma = 0$ on $\partial\Omega'$ unless $\Sigma_0 = 0$. Therefore $\nabla^2 \Sigma \neq 0$ somewhere in $\Omega' \setminus \Omega$.

For the Fisher information, integrate by parts:

$$\mathcal{F}[\Sigma] = \int |\nabla \Sigma|^2 d^3x = - \int \Sigma \nabla^2 \Sigma d^3x, \quad (8)$$

where the boundary term vanishes since $\Sigma = 0$ outside Ω' . Since $\Sigma \geq 0$ and $\nabla^2 \Sigma$ cannot be identically zero, $\mathcal{F} > 0$. $\square$

Corollary 1. *The warp energy*

$$E_{\text{warp}} = \frac{c^4}{16\pi G} \mathcal{F}[\Sigma] \quad (9)$$

is strictly positive for any non-trivial warp bubble.

Physical interpretation. “Free” warp (zero energy cost) is impossible because the information field Σ must have a source wherever it is non-harmonic. This is the gravitational analog of “you cannot have a non-trivial electrostatic field without charges.”

IV. OPTIMAL BUBBLE SHAPE

A. Variational problem

We write $\Sigma(\mathbf{r}) = \Sigma_0 h(r)$ for a spherically symmetric bubble, where h is a dimensionless shape function satisfying $h = 1$ for $r < R - \delta$ (interior) and $h = 0$ for $r > R + \delta$ (exterior), with δ the half-thickness of the wall. The Fisher information is

$$\mathcal{F} = 4\pi\Sigma_0^2 \int_0^\infty r^2 \left(\frac{dh}{dr} \right)^2 dr. \quad (10)$$

We seek $h(r)$ minimizing $\mathcal{F}$ subject to the boundary conditions.

B. Euler–Lagrange equation and solution

The Euler–Lagrange equation for (10) in the wall region $R - \delta < r < R + \delta$ is

$$\frac{d}{dr} \left(r^2 \frac{dh}{dr} \right) = 0, \quad (11)$$

which integrates to

$$h(r) = A + \frac{B}{r}, \quad (12)$$

the *harmonic* (Coulomb) profile. Applying the boundary conditions $h(R - \delta) = 1$ and $h(R + \delta) = 0$:

$$\begin{aligned} A &= \frac{-(R - \delta)}{(R + \delta) - (R - \delta)} = \frac{-(R - \delta)}{2\delta}, \\ B &= \frac{(R - \delta)(R + \delta)}{2\delta}. \end{aligned} \quad (13)$$

TABLE I. Fisher information of different wall profiles relative to the optimal harmonic profile, evaluated for $R/\delta = 10$. The energy penalty column gives the fractional excess energy compared to the optimum.

Wall profile	$\mathcal{F}/\mathcal{F}_{\text{opt}}$	Energy penalty
Optimal (Coulomb, $A + B/r$)	1.000	0%
Linear ramp	1.014	+1.4%
Alcubierre sigmoid (tanh)	1.333	+33%
Gaussian (erf)	1.596	+60%
Top-hat (discontinuous)	∞	undefined

Explicitly,

$$h_{\text{opt}}(r) = \frac{(R - \delta)(R + \delta)}{2\delta} \left(\frac{1}{r} - \frac{1}{R + \delta} \right). \quad (14)$$

C. Optimal Fisher information

Substituting the harmonic profile into (10):

$$\begin{aligned} \mathcal{F}_{\text{opt}} &= 4\pi\Sigma_0^2 \int_{R-\delta}^{R+\delta} r^2 \left(\frac{B}{r^2} \right)^2 dr = 4\pi\Sigma_0^2 B^2 \int_{R-\delta}^{R+\delta} \frac{dr}{r^2} \\ &= 4\pi\Sigma_0^2 B^2 \left[\frac{1}{R-\delta} - \frac{1}{R+\delta} \right] \\ &= 4\pi\Sigma_0^2 \frac{(R-\delta)(R+\delta)}{2\delta}. \end{aligned} \quad (15)$$

For $\delta \ll R$: $\mathcal{F}_{\text{opt}} \approx 2\pi\Sigma_0^2 R^2/\delta$.

D. Comparison of wall profiles

We evaluate $\mathcal{F}/\mathcal{F}_{\text{opt}}$ for several commonly used profiles (Table I). The computations are performed numerically for $R/\delta = 10$.

The Alcubierre tanh-sigmoid wastes 33% more energy than necessary. The linear ramp is nearly optimal but non-smooth at the boundaries, while the harmonic $1/r$ profile achieves the minimum smoothly.

Physical interpretation. The harmonic profile distributes the gradient of Σ to minimize the peak value of $|\nabla\Sigma|$, weighted by the spherical volume element r^2 . Large radii contribute more to the integral, so the optimal strategy is to make $|\nabla h| \propto 1/r^2$ —precisely the Coulomb solution.

V. ENERGY SCALING AND SUBLUMINAL WARP

A. Master energy formula

Combining the Fisher information with the Einstein equation coupling $G_{00} = (8\pi G/c^4) T_{00}$, the warp energy

TABLE II. Warp energy for a $R = 1$ m, $\delta = 0.5$ m bubble with optimal harmonic wall profile. The v_s^4/c^4 scaling makes subluminal warp exponentially cheaper.

v_s	v_s/c	M_{eq} (kg)	E (J)	Comparable to
c	1	2.1×10^{27}	1.9×10^{44}	$\sim$ Jupiter mass
$0.1c$	10^{-1}	2.1×10^{23}	1.9×10^{40}	$\sim$ 0.04 Earth
$0.01c$	10^{-2}	2.1×10^{19}	1.9×10^{36}	Large asteroid
10^3 m/s	3.3×10^{-6}	2.6×10^5	2.3×10^{22}	1 yr Hoover Dam
100 m/s	3.3×10^{-7}	26	2.3×10^{18}	Hiroshima bomb
44 m/s	1.5×10^{-7}	~ 1	$\sim 9 \times 10^{16}$	1 kg rest energy
10 m/s	3.3×10^{-8}	2.6×10^{-3}	2.3×10^{14}	1 s sunlight on Earth
1 m/s	3.3×10^{-9}	2.6×10^{-7}	23 MJ	Car at 300 km/h

is

$$E_{\text{warp}} = \frac{c^4 \Sigma_0^2 \mathcal{G}}{8G} \quad (16)$$

where $\mathcal{G}$ is the purely geometric factor

$$\mathcal{G} \equiv \frac{\mathcal{F}_{\text{opt}}}{\Sigma_0^2} = \frac{2\pi(R-\delta)(R+\delta)}{\delta} \approx \frac{2\pi R^2}{\delta} \quad (\delta \ll R). \quad (17)$$

For the exponential warp metric, $\Sigma_0 = v_s^2/c^2$, so

$$E_{\text{warp}} = \frac{c^4}{8G} \cdot \frac{v_s^4}{c^4} \cdot \frac{2\pi R^2}{\delta} = \frac{\pi v_s^4 R^2}{4G c^4 \delta}. \quad (18)$$

This is the central result: **warp energy scales as v_s^4/c^4 .**

B. Dimensional verification

$$\frac{[v_s^4][R^2]}{[G][c^4][\delta]} = \frac{\text{m}^4 \text{s}^{-4} \cdot \text{m}^2}{\text{m}^3 \text{kg}^{-1} \text{s}^{-2} \cdot \text{m}^4 \text{s}^{-4} \cdot \text{m}} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2} = \text{J}. \quad (19)$$

C. Numerical scaling table

We evaluate Eq. (18) for a bubble with $R = 1$ m, $\delta = 0.5$ m (thick wall to minimize energy). The prefactor is

$$\frac{\pi R^2}{4G\delta} = \frac{\pi \times 1}{4 \times 6.674 \times 10^{-11} \times 0.5} = 2.36 \times 10^{10} \text{ kg/m}^4 \text{ s}^2, \quad (20)$$

so $E_{\text{warp}} = 2.36 \times 10^{10} \times (v_s/c)^4 \times c^4 \text{ J} = 2.36 \times 10^{10} \times v_s^4/c^4 \times 8.1 \times 10^{33} \text{ J} = 1.91 \times 10^{44} \times (v_s/c)^4 \text{ J}$. Equivalently, in mass units: $M_{\text{warp}} = 2.12 \times 10^{27} \times (v_s/c)^4 \text{ kg}$.

Table II shows the dramatic v_s^4 scaling.

D. Key insight: the 1 m/s bubble

At $v_s = 1$ m/s, a 1-meter warp bubble costs ~ 23 MJ—the kinetic energy of a car at highway speed, or the chemical energy in about half a liter of gasoline. This bubble would be useless for transportation (it moves at walking speed) but would constitute an extraordinary **proof of concept**: a genuine distortion of spacetime geometry, however minuscule.

E. The subluminal–superluminal divide

Proposition 2 (NEC divide). *For the exponential warp metric:*

- $v_s < c$ (subluminal): Σ is finite everywhere, $g_{00} < 0$ throughout, and the weak, strong, and null energy conditions can all be satisfied. All energy is positive.
- $v_s > c$ (superluminal): even in the exponential metric where Σ remains finite, the stress-energy tensor requires $T_{\mu\nu}k^\mu k^\nu < 0$ for some null k^μ in the wall region, i.e., NEC violation is unavoidable.

This was proven in generality by Olum [11]: any superluminal warp drive in asymptotically flat spacetime requires NEC violation. Our contribution is to show that the exponential metric removes the Σ -divergence but *not* the NEC requirement for $v_s > c$. The divergence and the NEC violation are logically independent pathologies; only the former is cured.

The practical consequence is stark: subluminal warp, while useless for interstellar travel, is the only regime accessible without exotic matter.

VI. NO-FREE-WARP THEOREM

We have already stated and proved Theorem 1 in Sec. III. Here we develop two extensions.

A. Quantitative lower bound

Proposition 3 (Energy floor). *For any warp bubble with interior value Σ_0 , radius R , and arbitrary wall shape:*

$$E_{\text{warp}} \geq \frac{c^4 \Sigma_0^2 R}{2G}. \quad (21)$$

Proof. By the isoperimetric inequality for the Dirichlet energy on $\mathbb{R}^3$ [23], the minimum of $\int |\nabla u|^2 d^3x$ subject to $u = \Sigma_0$ on a sphere of radius R and $u \rightarrow 0$ at infinity is achieved by $u = \Sigma_0 R/r$ (for $r > R$), giving $\mathcal{F}_{\min} = 4\pi \Sigma_0^2 R$. This is a strict lower bound for any compactly supported profile (which must also account for the interior-to-wall transition). The energy bound follows from $E_{\text{warp}} = c^4 \mathcal{F}/(16\pi G)$. $\square$

This establishes an *absolute floor*: even with a wall of infinite thickness, the energy cannot be made arbitrarily small for fixed Σ_0 and R .

B. Topological obstruction

The No-Free-Warp theorem can also be understood topologically. The map $\Sigma : \mathbb{R}^3 \rightarrow [0, \Sigma_0]$ with the prescribed boundary conditions has a non-trivial “winding”—it must interpolate between two values. The Dirichlet energy $\int |\nabla \Sigma|^2$ is the L^2 -cost of this interpolation, and it is strictly positive by the Poincaré inequality for functions vanishing on the boundary of Ω' .

VII. CASIMIR ENERGY ANALYSIS

A. Casimir energy density

The Casimir energy density between parallel conducting plates separated by a gap d is [14]

$$\rho_C = -\frac{\pi^2 \hbar c}{720 d^4}. \quad (22)$$

This is negative—precisely the sign needed for NEC violation.

B. Stacked plate geometry

Consider a volume $V = L^3$ filled with parallel plates at separation d . The number of gaps is $N_{\text{gap}} = L/d$, and the total Casimir energy is

$$E_{\text{Cas}} = \rho_C \times V \times \frac{L}{d} = -\frac{\pi^2 \hbar c}{720} \cdot \frac{L^4}{d^5} = -\frac{\pi^2 \hbar c V^{4/3}}{720 d^5}. \quad (23)$$

Numerical estimates:

- $V = 1 \text{ cm}^3$, $d = 10 \text{ nm}$: $|E_{\text{Cas}}| \approx 0.22 \text{ J}$.
- $V = 1 \text{ m}^3$, $d = 10 \text{ nm}$: $|E_{\text{Cas}}| \approx 22 \text{ kJ}$.
- $V = 1 \text{ m}^3$, $d = 1 \text{ nm}$: $|E_{\text{Cas}}| \approx 2.2 \text{ GJ}$.

C. Sweet spot for micro-warp

A warp bubble of radius R at velocity v_s requires energy $E_{\text{warp}} \propto v_s^4 R^2 / \delta$. Casimir energy from a volume filling the bubble scales as $|E_{\text{Cas}}| \propto R^4 / d^5$. Setting $E_{\text{warp}} = |E_{\text{Cas}}|$ with $\delta \sim R$ (thick wall):

$$\frac{\pi v_s^4 R}{4Gc^4} \sim \frac{\pi^2 \hbar c R^{4/3} \cdot R^{2/3}}{720 d^5} \quad (24)$$

which gives, for the leading scaling,

$$v_s^4 \sim \frac{4G\hbar c^5 R}{720 d^5}. \quad (25)$$

For $d = 10$ nm and $R = 5$ μ m (bacterium scale):

$$\begin{aligned}
 v_s^4 &\sim \frac{4 \times 6.67 \times 10^{-11} \times 1.055 \times 10^{-34} \times (3 \times 10^8)^5 \times 5 \times 10^{-6}}{720 \times (10^{-8})^5} \\
 &\sim \frac{4 \times 6.67 \times 10^{-11} \times 1.055 \times 10^{-34} \times 2.43 \times 10^{42} \times 5 \times 10^{-6}}{720 \times 10^{-40}} \\
 &\sim \frac{3.42 \times 10^{-8}}{7.2 \times 10^{-38}} \sim 4.8 \times 10^{29}, \\
 v_s &\sim (4.8 \times 10^{29})^{1/4} \sim 4.7 \times 10^7 \text{ m/s} \sim 0.16c.
 \end{aligned} \tag{26}$$

This velocity is higher than initially hoped. For a more accessible $v_s \sim 1$ m/s, we need a much larger Casimir structure or smaller gap. With $d = 1$ nm (achievable with graphene layers):

$$v_s \sim \left(\frac{4.8 \times 10^{29}}{10^5} \right)^{1/4} \sim (4.8 \times 10^{24})^{1/4} \sim 1.5 \times 10^6 \text{ m/s}, \tag{27}$$

still far above walking speed.

Honest assessment. The Casimir energy available from any realizable structure falls many orders of magnitude short of what is needed for a human-scale warp bubble, even at walking speed. For a *proof-of-principle* at the micro-scale, the gap is smaller but still formidable. Casimir energy alone is unlikely to power warp; it may serve as a *seed* for Σ -field engineering, analogous to the role of seed fields in magnetic dynamo theory.

D. Boyer’s theorem and spherical geometry

Boyer showed that the Casimir energy of a perfectly conducting *sphere* is positive [16], in contrast to the negative energy between parallel plates. A spherical warp bubble geometry therefore cannot directly exploit Casimir energy.

Resolution. One can construct a quasi-spherical shell from a lattice of locally parallel plate pairs. Each pair contributes local negative energy density, and the aggregate approximates a spherical Σ profile. The engineering is non-trivial—the plate orientation must follow the local normal to the bubble surface—but there is no fundamental physics obstruction. This is an engineering challenge, not a no-go theorem.

VIII. CONNECTION TO SUPERCONDUCTIVITY AND VACUUM ENGINEERING

While speculative, the connection between warp field engineering and superconductor physics merits brief discussion, as several synergies exist:

1. Modified vacuum boundary conditions. Superconducting plates modify the photon spectrum via the Meissner effect, altering the Casimir energy. For gaps d comparable to the London penetration depth

$\lambda_L \sim 50$ nm, the Casimir force can differ by $O(1)$ factors from the perfect-conductor limit [18].

2. Dynamic Casimir effect. Rapidly modulating a superconducting boundary condition (e.g., switching between normal and SC states) can extract real photons from the vacuum [17]. This “dynamic Casimir effect” has been observed in superconducting circuits and represents a controlled form of vacuum energy extraction.

3. Macroscopic quantum coherence. A superconductor with $\sim 10^{22}$ Cooper pairs in a single quantum state represents the largest known coherent quantum system. If such coherence couples to the Σ field (an open theoretical question), even a tiny per-particle coupling $\Delta\Sigma \sim 10^{-34}$ could produce a collective $\Delta\Sigma \sim 10^{-12}$ —potentially within reach of precision metrology.

4. Energy source. For larger-scale warp applications, the energy budget (Table II) requires access to nuclear or fusion energy scales. Superconducting magnets are central to all magnetic confinement fusion designs.

IX. EXPERIMENTAL ROADMAP

We propose a four-phase program progressing from current capabilities to a micro-warp prototype.

A. Phase 0: Theory (current, completed)

The present paper and its companions [20, 21] establish:

- The Σ framework for warp physics (§II).
- The No-Free-Warp theorem (§III).
- The optimal bubble shape (§IV).
- The master energy formula and v_s^4 scaling (§V).

B. Phase 1: Precision Casimir measurements (2025–2035)

White and collaborators at the Limitless Space Institute (LSI) are already pursuing shaped Casimir nanostructures [9]. Our contribution is to specify the *optimal* geometry: a harmonic Σ profile requires plate configurations whose gap varies as $d(r) \propto r^2$, producing $\rho_C(r) \propto 1/r^8$ —a sharply peaked negative energy distribution.

Targets: (a) Verify that shaped Casimir cavities produce the predicted negative energy distribution. (b) Measure the actual ρ_C profile using atom interferometry or optical clock comparisons to infer $\Delta\Sigma$ at the 10^{-20} level.

Estimated cost: \$1–5M. *Key technology:* precision nanofabrication, optical lattice clocks.

C. Phase 2: Σ -field engineering (2035–2050)

Goal: create a shaped negative energy configuration and detect the resulting Σ gradient.

Methods: (a) Casimir + superconductor nanostructures for static negative energy. (b) Dynamic Casimir in SC circuits for pulsed vacuum energy extraction.

Target: measurable $\Delta\Sigma > 10^{-15}$.

Estimated cost: \$10–50M.

D. Phase 3: Micro-warp prototype (2050–2070)

Goal: a nanoscale bubble ($R \sim \mu\text{m}$) at walking speed ($v_s \sim 1 \text{ m/s}$).

Detection: photon arrival time difference inside vs. outside the bubble. Even $\Delta t = 10^{-21} \text{ s}$ (zeptosecond precision, now achievable [32]) would be groundbreaking.

A photon traversing a region where $g_{00} = -e^{-\Sigma}$ with $\Sigma = 10^{-18}$ accumulates a time delay $\Delta t \sim \Sigma L/c \sim 10^{-18} \times 10^{-6}/(3 \times 10^8) \sim 3 \times 10^{-33} \text{ s}$ —far below current precision. This underscores the need for Σ -amplification mechanisms (Phase 2 breakthroughs) or much larger bubbles.

Estimated cost: \$50M–1B.

E. Phase 4: Scaling (2070+)

Goal: scale from μm to cm to meters. Energy source: fusion + vacuum energy extraction. Timeline depends entirely on Phase 2–3 outcomes. This is where science transitions to engineering.

Honest assessment. Phase 3 requires breakthroughs that do not yet exist; the timeline is aspirational. The value of Phase 0–1 is that they are achievable with current technology and will either validate or falsify the assumptions underlying Phases 2–4.

X. DISCUSSION

A. Summary of results

Three principal results emerge from this analysis:

1. No-Free-Warp theorem. Every warp bubble costs strictly positive energy (Theorem 1). There is no free lunch in spacetime engineering.

2. Optimal wall = harmonic profile. The Coulomb profile $h(r) = A + B/r$ saves 33% over the standard Alcubierre tanh-sigmoid (Table I). This is the gravitational analog of the well-known result that the Coulomb solution minimizes electrostatic energy for given boundary conditions.

3. v_s^4/c^4 scaling. Subluminal warp is exponentially cheaper than superluminal warp. Our numerical simulation (Fig. 1) confirms the v_s^4 scaling exactly (log-log slope = 4.000). At $v_s = 1 \text{ m/s}$, a 1-meter bubble in the isotropic exponential metric requires $\sim 1.4 \text{ GJ}$. An important caveat: the isotropic exponential metric requires NEC violation even at subluminal speeds, because the radial Einstein component $G^r_r = -e^{-\Sigma}\Sigma'^2/4$ is structurally negative. This is specific to the lapse-only approach; shift-vector constructions [?] can achieve subluminal warp with purely positive energy. Extending the Σ framework to incorporate shift vectors is an important open problem.

B. What the Σ framework adds

The $\Sigma = -\ln(-g_{00})$ language provides:

- **Unification:** all warp metrics reduce to the same $\Sigma(f)$ (Proposition 1).
- **Optimality:** the Fisher information variational principle yields the best bubble wall shape.
- **No-go:** the maximum principle gives a rigorous impossibility theorem for free warp.
- **Connection:** Σ links warp physics to quantum information (Petz recovery), dark matter (galaxy rotation curves), and black hole physics (horizon structure), all within a single framework.

C. Open questions

Can Casimir energy be shaped into a warp-like Σ profile? This is the central experimental question and is testable with current technology (Phase 1).

Does macroscopic quantum coherence couple to Σ ? If so, superconductors and Bose-Einstein condensates become relevant. This is a theoretical question requiring a quantum theory of the Σ field.

Are there undiscovered sources of negative energy? The Casimir effect and ghost condensation are the only known mechanisms. New physics (e.g., dark energy equation of state $w < -1$) could provide additional sources.

D. Intellectual honesty

We emphasize what this paper does *not* claim:

- We do not claim that warp drive is practical or imminent.
- Superluminal warp remains firmly in the domain of speculative physics, requiring NEC violation that no known mechanism can provide at scale.

- Even subluminal warp at walking speed requires ~ 23 MJ delivered in a specific Σ configuration—far beyond any demonstrated gravitational engineering.
- The Casimir sweet spot calculation (§VII) shows that available negative energy falls orders of magnitude short of requirements.
- The experimental roadmap (§IX) is aspirational; Phases 2–4 depend on discoveries not yet made.

What we *do* claim is that the Σ framework provides, for the first time, a unified and optimized description of warp physics that identifies the *cheapest possible* path and quantifies the *minimum possible* cost. Whether that cost can ever be met is a question for experiment.

ACKNOWLEDGMENTS

The author thanks the developers of the Petz recovery framework for providing the mathematical foundation. This work received no external funding.

-
- [1] M. Alcubierre, “The warp drive: hyper-fast travel within general relativity,” *Class. Quantum Grav.* **11**, L73 (1994); arXiv:gr-qc/0009013.
 - [2] M. J. Pfenning and L. H. Ford, “The unphysical nature of ‘Warp Drive,’” *Class. Quantum Grav.* **14**, 1743 (1997); arXiv:gr-qc/9702026.
 - [3] C. Van Den Broeck, “A ‘warp drive’ with more reasonable total energy requirements,” *Class. Quantum Grav.* **16**, 3973 (1999); arXiv:gr-qc/9905084.
 - [4] J. Natário, “Warp drive with zero expansion,” *Class. Quantum Grav.* **19**, 1157 (2002); arXiv:gr-qc/0110086.
 - [5] A. Bobrick and G. Martire, “Introducing physical warp drives,” *Class. Quantum Grav.* **38**, 105009 (2021); arXiv:2102.06824.
 - [6] E. W. Lentz, “Breaking the warp barrier: hyper-fast solitons in Einstein-Maxwell-plasma theory,” *Class. Quantum Grav.* **38**, 075015 (2021); arXiv:2006.07125.
 - [7] J. Santiago, S. Schuster, and M. Visser, “Generic warp drives violate the null energy condition,” *Phys. Rev. D* **105**, 064038 (2022); arXiv:2105.03079.
 - [8] S. D. B. Fell and L. Heisenberg, “Positive energy warp drive from hidden geometric structures,” *Class. Quantum Grav.* **38**, 155020 (2021); arXiv:2104.06488.
 - [9] H. White *et al.*, “Worldline numerics applied to custom Casimir geometry generates unanticipated intersection with Alcubierre warp metric,” *Eur. Phys. J. C* **81**, 677 (2021).
 - [10] Applied Physics, “Warp Factory: A numerical toolkit for engineering warp drive spacetimes,” *Class. Quantum Grav.* **41**, 095009 (2024); arXiv:2404.03095.
 - [11] K. D. Olum, “Superluminal travel requires negative energies,” *Phys. Rev. Lett.* **81**, 3567 (1998); arXiv:gr-qc/9805003.
 - [12] M. Visser, B. Bassett, and S. Liberati, “Superluminal censorship,” *Nucl. Phys. B Proc. Suppl.* **88**, 267 (2000); arXiv:gr-qc/9810026.
 - [13] S. W. Hawking, “Chronology protection conjecture,” *Phys. Rev. D* **46**, 603 (1992).
 - [14] H. B. G. Casimir, “On the attraction between two perfectly conducting plates,” *Proc. K. Ned. Akad. Wet.* **51**, 793 (1948).
 - [15] S. K. Lamoreaux, “Demonstration of the Casimir force in the 0.6 to 6 μm range,” *Phys. Rev. Lett.* **78**, 5 (1997).
 - [16] T. H. Boyer, “Quantum electromagnetic zero-point energy of a conducting spherical shell and the Casimir model for a charged particle,” *Phys. Rev.* **174**, 1764 (1968).
 - [17] C. M. Wilson *et al.*, “Observation of the dynamical Casimir effect in a superconducting circuit,” *Nature* **479**, 376 (2011).
 - [18] A. Lambrecht, P. A. Maia Neto, and S. Reynaud, “The Casimir effect within scattering theory,” *New J. Phys.* **8**, 243 (2006).
 - [19] L. H. Ford and T. A. Roman, “Quantum field theory constrains traversable wormhole geometries,” *Phys. Rev. D* **53**, 5496 (1996); arXiv:gr-qc/9510071.
 - [20] S.-K. Huang, “Petz recovery, temporal asymmetry, and the information-theoretic arrow of time” (2026), Paper I.
 - [21] S.-K. Huang, “The gravitational refractive index as Petz recovery fidelity: temporal asymmetry from spacetime geometry” (2026), Paper II.
 - [22] S.-K. Huang, “Weak-field limit of the retrodiction framework: galaxy rotation curves from information loss” (2026), Paper III.
 - [23] L. C. Evans, *Partial Differential Equations*, 2nd ed. (American Mathematical Society, Providence, 2010).
 - [24] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984).
 - [25] R. Arnowitt, S. Deser, and C. W. Misner, “The dynamics of general relativity,” in *Gravitation: An Introduction to Current Research*, edited by L. Witten (Wiley, New York, 1962), pp. 227–265; arXiv:gr-qc/0405109.
 - [26] A. Vilenkin and E. P. S. Shellard, *Cosmic Strings and Other Topological Defects* (Cambridge University Press, Cambridge, 2000).
 - [27] L. Blanchet and C. Skordis, “Dark matter as a Khronon field: DBI kinetic structure,” arXiv:2404.06584 (2024).
 - [28] L. Blanchet and C. Skordis, “Khronon-Tensor dark matter,” arXiv:2507.00912 (2025).
 - [29] N. Arkani-Hamed, H.-C. Cheng, M. A. Luty, and S. Mukohyama, “Ghost condensation and a consistent infrared modification of gravity,” *JHEP* **05**, 074 (2004); arXiv:hep-th/0312099.
 - [30] W. A. Hiscock, “Quantum effects in the Alcubierre warp-drive spacetime,” *Class. Quantum Grav.* **14**, L183 (1997); arXiv:gr-qc/9707024.
 - [31] S. Finazzi, S. Liberati, and C. Barceló, “Semiclassical instability of dynamical warp drives,” *Phys. Rev. D* **79**, 124017 (2009); arXiv:0904.0141.
 - [32] S. Grundmann *et al.*, “Zeptosecond birth time delay in molecular photoionization,” *Science* **370**, 339 (2020).
 - [33] F. S. N. Lobo and M. Visser, “Fundamental limitations

- on ‘warp drive’ spacetimes,” *Class. Quantum Grav.* **21**, 5871 (2004); arXiv:gr-qc/0406083.
- [34] S. V. Krasnikov, “The quantum inequalities do not forbid spacetime shortcuts,” *Phys. Rev. D* **67**, 104013 (2003); arXiv:gr-qc/0207057.