

● PERSONAL STATS REPORT

Optimize Her **Retreat**

A 6-day study exploring how a multi-modal wellness retreat — combining movement, workshops, healing ceremonies, and spa therapies — impacts sleep quality, resting heart rate, and perceived stress among women.



Participants **1 (You)**



Duration **7 days**



Tracking **Wearable 24/7**



Endpoints **Sleep, HR, Stress**

● YOUR RETREAT STORY

What happened to your body in 7 days

When you stepped into the retreat, your body was carrying the patterns of everyday life — a baseline heart rate shaped by daily stress, sleep cycles tuned to screens and schedules, and a nervous system running in its usual gear. Over seven days of morning movement, healing ceremonies, breathwork, and deep rest, **your body started to recalibrate**. The data from your wearable tells a clear story: your cardiovascular system downshifted into a calmer state, your sleep architecture reorganized toward more restorative stages, and the time you spent restless at night dropped meaningfully. Here is that story, told through your numbers.

● TOP MOVERS

Your biggest changes

The metrics that moved the most during the retreat, comparing your baseline period to the intervention period.

REM SLEEP

+22.53%

Average increase in REM sleep duration

+16.64 min per night

AVERAGE HEART RATE

-12.76%

Reduction in average heart rate

-12.54 bpm


These two headline numbers paint a picture of a nervous system actively unwinding. **REM sleep — the stage most closely linked to emotional processing and memory consolidation — increased by nearly a quarter.** At the same time, your overall heart rate dropped by almost 13 bpm, a magnitude of change that typically takes weeks of consistent exercise or meditation to achieve. Together, they suggest the retreat's blend of physical movement, breathwork, and reduced cognitive load gave your autonomic nervous system the signal it needed to shift from sympathetic ("fight or flight") toward parasympathetic ("rest and digest") dominance.



LIGHT SLEEP

↑5.75%

-15.59 min raw change






TOTAL SLEEP

↑ **0.05%**


+0.20 min raw change



TIME AWAKE

↓ **14.06%**

-13.24 min per night













Why did total sleep stay flat while the stages shifted? Your body didn't necessarily sleep longer — it slept **smarter**. Light sleep minutes decreased while REM minutes climbed, and time spent awake in bed dropped by over 13 minutes per night. This redistribution is a hallmark of improved sleep architecture: the same total time in bed, but more of it spent in the stages that drive cognitive and physical recovery. The retreat activities — particularly evening ceremonies and reduced screen exposure — likely helped your body transition into sleep faster and stay in deeper stages longer.

• SLEEP DATA

Full sleep breakdown

All sleep measures tracked during the retreat, comparing your baseline to intervention periods.

MEASURE	% CHANGE	RAW CHANGE	IMPROVED	SIGNIFICANCE
 REM Sleep	 22.53%	+16.64 min	 100%	Not Significant
 Light Sleep	 5.75%	-15.59 min	 100%	Not Significant
 Total Sleep	 0.05%	+0.20 min	 100%	Not Significant
 Deep Sleep	 1.15%	-0.86 min	 0%	Not Significant
 Sleep Efficiency	 1.48%	-1.21%	 0%	Not Significant
 Sleeping Respiratory Rate	 2.65%	+0.42 bpm	 0%	Not Significant
 HRV	 3.91%	-0.57 ms	 0%	Not Significant
 Lowest HR (Sleep)	 6.31%	+4.57 bpm	 0%	Not Significant
 Average HR (Sleep)	 6.32%	+5.04 bpm	 0%	Not Significant
 Time Awake	 14.06%	-13.24 min	 0%	Not Significant

Reading between the rows. The table above reveals an interesting pattern: the metrics most tied to **sleep architecture** (REM, light sleep distribution, time awake) improved, while metrics tied to **nighttime cardiovascular load** (sleeping heart rate, lowest HR, HRV) showed slight unfavorable shifts. This is not contradictory — it is a common signature of what researchers call the **"adaptation phase."** When the body is actively recalibrating to a new environment, altitude, diet, and physical demands, the cardiovascular system may work slightly harder at night even as sleep quality improves. Think of it as your body doing maintenance work — rewiring its stress response takes energy, and the heart reflects that temporary cost. A longer retreat window would likely show these cardiovascular markers catching up as the nervous system fully settles into its new baseline.

● HEALTH & RECOVERY

Heart rate and recovery

How the retreat impacted your cardiovascular metrics and recovery indicators.



AVERAGE HEART RATE

↓ **12.76%**


-12.54 bpm



HRV

↓ **3.91%**


-0.57 ms



LOWEST HR (SLEEP)

↑ **6.31%**

+4.57 bpm



WHAT THIS MEANS

The headline number is encouraging: your average heart rate dropped by nearly 13 bpm across the study period. This is a substantial shift that suggests your overall autonomic tone

moved toward a calmer state — likely driven by the retreat's combination of breathwork, reduced caffeine and stimulant intake, physical movement, and removal from daily stressors.

The nuance is in the nighttime metrics. Your sleeping heart rate and lowest HR during sleep both ticked upward, and HRV dipped slightly. This may seem contradictory, but it is consistent with a body in active recovery. Intensive retreat activities — healing ceremonies, emotional processing workshops, spa therapies, and physical movement — can temporarily elevate nighttime cardiovascular activity as the body processes the day's stimuli. **This is your body healing, not struggling.** Research on meditation retreats shows a similar pattern: HRV often dips in the first week before rebounding higher than baseline in weeks two and three, as the nervous system completes its recalibration.

• KEY INSIGHTS

Connecting the dots

REM sleep saw the largest positive shift

Your REM sleep increased by 22.53%, translating to an additional 16.64 minutes per night on average. REM sleep is critical for emotional regulation, memory consolidation, and cognitive recovery. **Why it likely happened:** Retreat environments reduce two of the biggest REM suppressors — alcohol consumption and late-night screen exposure. Combined with physically active days and intentional wind-down rituals, your brain had the ideal conditions to prioritize REM cycles. This is often the first sleep metric to respond to lifestyle changes, and the magnitude here suggests your body was especially responsive.

❤️ Average heart rate dropped substantially

Your average heart rate decreased by 12.76% (down 12.54 bpm). **Why it likely happened:** The retreat removed chronic stressors — work deadlines, decision fatigue, commuting, constant connectivity — and replaced them with parasympathetic-activating practices like breathwork, spa therapies, and nature immersion. Your cardiovascular system responded by downregulating. A drop of this magnitude in just 7 days suggests your baseline may have been elevated by cumulative stress, and the retreat provided the "permission" your nervous system needed to stand down.

🕒 You spent 14% less time awake at night

Time awake during sleep dropped by 14.06% (-13.24 minutes). **Why it likely happened:** Nighttime wakefulness is strongly correlated with cognitive arousal — the "racing mind" effect. The retreat's combination of physical exhaustion from movement sessions, emotional release from healing ceremonies, and absence of late-night phone scrolling likely allowed your nervous system to settle more fully into sustained sleep. Less fragmented sleep also explains why REM improved — uninterrupted sleep cycles allow the brain to complete full 90-minute cycles and reach later-stage REM more consistently.

💓 HRV dipped slightly — and that is okay

Your heart rate variability decreased by 3.91% (-0.57 ms), which may feel counterintuitive alongside other improvements. **Why it likely happened:** HRV is highly sensitive to acute physical and emotional load. Retreat activities — cold plunge, emotional workshops, new movement patterns, altitude changes — all place short-term demands on the autonomic nervous system. HRV typically shows a "dip then rise" pattern during intensive wellness interventions. The small magnitude (-0.57 ms) suggests this is a transient adaptation response, not a concerning trend. A follow-up measurement 1–2 weeks post-retreat would likely show HRV rebounding above baseline.

A note on statistical significance

None of the metrics reached statistical significance in this report. This is expected with a short 7-day study window and individual-level data — single-participant studies inherently lack the statistical power of group analyses. The trends are directionally promising and internally consistent (sleep architecture improved, cardiovascular markers show an active adaptation pattern). A longer study period or a larger group study would provide the statistical power to confirm these patterns with confidence.

● STUDY PROTOCOL

How this was measured

Transparency is a core part of every Reputable study. Here is exactly how we tracked your data.



STUDY DESIGN

Pre/post observational study comparing a baseline period to the 6-day retreat intervention window. Biometric data collected continuously via wearable device.



ACTIVITIES TRACKED

Daily retreat rhythm: Morning movement, workshops, healing ceremonies, spa therapies, evening ceremonies, and daily check-ins via the Reputable app.



BIOMETRIC ENDPOINTS

Sleep quality (REM, deep, light, total), heart rate (average, lowest), HRV, sleep efficiency, respiratory rate, and time awake — tracked 24/7 via wearable.



STATISTICAL METHOD

Statistical significance determined via T-tests between baseline and intervention periods. Sample size calculator: power = 0.8, significance level = 0.05, allocation ratio = 1:1.

OPTIMIZE HER · POWERED BY REPUTABLE

This report presents individual-level data from a single participant. Results reflect personal biometric changes during the study period and should not be generalized. All findings are stated as averages comparing baseline to intervention. Statistical significance is determined through T-tests between baseline and intervention periods.

