

# SML

January 2020

## Assignment 1

Q1. Consider the following decision rule for a two-category one-dimensional problem:

Decide  $\omega_1$  if  $x > \theta$ ; otherwise decide  $\omega_2$ .

(a) Show that the probability of error for this rule is given by

$$P(\text{error}) = P(\omega_1) \int_{-\inf}^{\theta} p(x|\omega_1) dx + P(\omega_2) \int_{\theta}^{\inf} p(x|\omega_2) dx \quad (1)$$

(b) By differentiating, show that a necessary condition to minimize  $P(\text{error})$  is that  $\theta$  satisfy  $p(\theta|\omega_1)P(\omega_1) = p(\theta|\omega_2)P(\omega_2)$

Q2. Let the conditional densities for a two-category one-dimensional problem be given by the Cauchy distribution

$$p(x|\omega_i) = \frac{1}{\pi b} \frac{1}{1 + \left(\frac{x-a_i}{b}\right)^2}, \quad i = 1, 2 \quad (2)$$

Assuming  $P(\omega_1) = P(\omega_2)$ , show that  $P(\omega_1|x) = P(\omega_2|x)$  if  $x = (a_1 + a_2)/2$ , i.e., the minimum error decision boundary is a point midway between the peaks of the two distributions, regardless of  $b$ .

Q3. Suppose we have three equi-probable categories in two dimensions with the following underlying distributions:

$$\begin{aligned} p(x|\omega_1) &\sim N(0, I) \\ p(x|\omega_2) &\sim N\left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}, I\right) \\ p(x|\omega_3) &\sim 0.5N\left(\begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix}, I\right) + 0.5N\left(\begin{bmatrix} -0.5 \\ 0.5 \end{bmatrix}, I\right) \end{aligned}$$

By explicit calculation of posterior probabilities, classify the point  $\mathbf{x} = \begin{bmatrix} 0.3 \\ 0.3 \end{bmatrix}$  for minimum probability of error.

Q4. a. Write a procedure to generate random samples according to a normal distribution  $N(\mu, \Sigma)$  in  $d$  dimensions.

b. Write a procedure to calculate the discriminant function for a given normal distribution with  $\Sigma = \sigma^2 I$  and prior probability  $P(\omega_i)$ .

c. Compare the discriminant function's values for two different distributions  $N(\mu_1 = \begin{bmatrix} 0.5 \\ 0.5 \end{bmatrix}, \Sigma_1 = \sigma^2 I)$  and  $N(\mu_2 = \begin{bmatrix} -0.5 \\ 0.5 \end{bmatrix}, \Sigma_2 = \sigma^2 I)$  in  $d = 2$  dimensions.

Assume the test sample to be  $X = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  and  $P(\omega_1) = 1/3$  and  $P(\omega_2) = 2/3$ .

In a general process, you would be given several samples from two (or more) classes. Counting each class' frequency will give the priors. With these samples as  $d$  dimensional vectors, you can estimate mean and covariance using MLE or other techniques, which is a part of later lecture. This computed info is sufficient for computing discriminants and thereby classifying the sample into one of the classes.