

Equations over Groups

Christian Sattler

University of Nottingham

June 2013


An Elementary Problem

An Elementary Problem

Consider an n -gon with edges oriented arbitrarily and corners uniquely labelled.

An Elementary Problem


Consider an n -gon with edges oriented arbitrarily and corners uniquely labelled.



An Elementary Problem

Consider an n -gon with edges oriented arbitrarily and corners uniquely labelled.



Consider also its mirror image.



An Elementary Problem

Consider an n -gon with edges oriented arbitrarily and corners uniquely labelled.



Consider also its mirror image.



An Elementary Problem

Consider an n -gon with edges oriented arbitrarily and corners uniquely labelled.

Consider also its mirror image.





Let a topological tiling of the sphere with copies of these tiles be given.

An Elementary Problem


Consider an n -gon with edges oriented arbitrarily and corners uniquely labelled.

Consider also its mirror image.



Let a topological tiling of the sphere with copies of these tiles be given.

Edge orientations must match!




An Elementary Problem


Can all corners be grouped in pairs of the form (u, u^{-1}) and connected by non-intersecting lines?

An Elementary Problem

Can all corners be grouped in pairs of the form (u, u^{-1}) and connected by non-intersecting lines?

Conjecture: If positively oriented edges equal negatively oriented edges in number, this will always be possible.





An Abstract Problem

An Abstract Problem

(Recall: free product $G * H$ is coproduct in category of groups.)

An Abstract Problem

(Recall: free product $G * H$ is coproduct in category of groups.)

(Recall: S^H denotes normal closure of set S in H .)

An Abstract Problem

Group G , word $r \in G * \mathbb{Z}$

An Abstract Problem

Group G , word $r \in G * \mathbb{Z}$

Is the composition

$$G \hookrightarrow G * \mathbb{Z} \rightarrow (G * \mathbb{Z}) / \langle r^{G * \mathbb{Z}} \rangle$$

monic?

An Abstract Problem

Group G , word $r \in G * \mathbb{Z}$

Is the composition

$$G \hookrightarrow G * \mathbb{Z} \rightarrow (G * \mathbb{Z}) / \langle r^{G * \mathbb{Z}} \rangle$$

monic?

Conjecture: Define $s : G * \mathbb{Z} \rightarrow \mathbb{Z}$ by $s = [\text{const } 0, \text{id}]$.
If $s(r) \neq 0$, the above holds.

An Important Problem

An Important Problem

Group G , “unknown” t , equation $a_1 t^{d_1} \dots a_n t^{d_n} = e$ over G

An Important Problem

Group G , “unknown” t , equation $a_1 t^{d_1} \dots a_n t^{d_n} = e$ over G

Does the equation have a solution in a *group extension* of G ?

An Important Problem

Group G , “unknown” t , equation $a_1 t^{d_1} \dots a_n t^{d_n} = e$ over G

Does the equation have a solution in a *group extension* of G ?

Kervaire-Laudenbach Conjecture:

If $d_1 + \dots + d_n \neq 0$, the answer is affirmative.

An Important Problem

Group G , “unknown” t , equation $a_1 t^{d_1} \dots a_n t^{d_n} = e$ over G

Does the equation have a solution in a *group extension* of G ?

Kervaire-Laudenbach Conjecture:

If $d_1 + \dots + d_n \neq 0$, the answer is affirmative.

For fields, the equivalent problem of synthesizing roots of polynomials in a field extension is treated in every algebra course.

An Important Problem

Group G , “unknown” t , equation $a_1 t^{d_1} \dots a_n t^{d_n} = e$ over G

Does the equation have a solution in a *group extension* of G ?

Kervaire-Laudenbach Conjecture:

If $d_1 + \dots + d_n \neq 0$, the answer is affirmative.

For fields, the equivalent problem of synthesizing roots of polynomials in a field extension is treated in every algebra course.

For groups, it turns out to be surprisingly difficult.

An Important Problem

Group G , “unknown” t , equation $a_1 t^{d_1} \dots a_n t^{d_n} = e$ over G

Does the equation have a solution in a *group extension* of G ?

Kervaire-Laudenbach Conjecture:

If $d_1 + \dots + d_n \neq 0$, the answer is affirmative.

For fields, the equivalent problem of synthesizing roots of polynomials in a field extension is treated in every algebra course.

For groups, it turns out to be surprisingly difficult.

Groups are perverse objects.

Revelations

Revelations

All these problems are actually the same!

An Important Problem: Progress

$$x^n = g \text{ (Neumann, 1949)}$$

An Important Problem: Progress

$$x^n = g \text{ (Neumann, 1949)}$$

$$d_1, \dots, d_n \geq 0 \text{ (Levin, 1962)}$$

An Important Problem: Progress

$$x^n = g \text{ (Neumann, 1949)}$$

$$d_1, \dots, d_n \geq 0 \text{ (Levin, 1962)}$$

G locally residually finite (Rothaus, 1977)

An Important Problem: Progress

$$x^n = g \text{ (Neumann, 1949)}$$

$$d_1, \dots, d_n \geq 0 \text{ (Levin, 1962)}$$

G locally residually finite (Rothaus, 1977)

$$|d_1| + \dots + |d_n| \leq 3 \text{ (Howie, 1983)}$$

An Important Problem: Progress

$$x^n = g \text{ (Neumann, 1949)}$$

$$d_1, \dots, d_n \geq 0 \text{ (Levin, 1962)}$$

$$G \text{ locally residually finite (Rothaus, 1977)}$$

$$|d_1| + \dots + |d_n| \leq 3 \text{ (Howie, 1983)}$$

$$|d_1| + \dots + |d_n| \leq 4 \text{ (Edjvet and Howie, 1991)}$$

An Important Problem: Progress

$$x^n = g \text{ (Neumann, 1949)}$$

$$d_1, \dots, d_n \geq 0 \text{ (Levin, 1962)}$$

$$G \text{ locally residually finite (Rothaus, 1977)}$$

$$|d_1| + \dots + |d_n| \leq 3 \text{ (Howie, 1983)}$$

$$|d_1| + \dots + |d_n| \leq 4 \text{ (Edjvet and Howie, 1991)}$$

$$|d_1| + \dots + |d_n| \leq 5 \text{ (Evangelidou, 2003)}$$

An Important Problem: Problems

Case distinction trees grow exponentially in equation length.

An Important Problem: Problems

Case distinction trees grow exponentially in equation length.

Every case needs to be treated differently.

An Important Problem: Problems

Case distinction trees grow exponentially in equation length.

Every case needs to be treated differently.

Several complementary techniques were developed by Edjvet et.al.

An Important Problem: Problems

Case distinction trees grow exponentially in equation length.

Every case needs to be treated differently.

Several complementary techniques were developed by Edjvet et.al.

Ideal case study of using dependent type systems for real-life proofs.

An Important Problem: Problems

Case distinction trees grow exponentially in equation length.

Every case needs to be treated differently.

Several complementary techniques were developed by Edjvet et.al.

Ideal case study of using dependent type systems for real-life proofs.

$|d_1| + \dots + |d_n| \leq 6$ (Agda and Coq, 2014)?

An Important Problem: Problems

Case distinction trees grow exponentially in equation length.

Every case needs to be treated differently.

Several complementary techniques were developed by Edjvet et.al.

Ideal case study of using dependent type systems for real-life proofs.

$|d_1| + \dots + |d_n| \leq 6 + x$ (Agda and Coq, 2014)?